



Moderate Grazing During the Off-Season (Fall-Winter) Reduces Exotic Annual Grasses in Sagebrush-Bunchgrass Steppe☆

Authors: Davies, Kirk W., Boyd, Chad S., Copeland, Stella M., and Bates, Jon D.

Source: Rangeland Ecology and Management, 82(1) : 51-57

Published By: Society for Range Management


URL: <https://doi.org/10.1016/j.rama.2022.02.003>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.



Original Research

Moderate Grazing During the Off-Season (Fall-Winter) Reduces Exotic Annual Grasses in Sagebrush-Bunchgrass Steppe[☆]

Kirk W. Davies*, Chad S. Boyd, Stella M. Copeland, Jon D. Bates

US Department of Agriculture–Agricultural Research Service, Eastern Oregon Agricultural Research Center, Burns, OR, 97720



ARTICLE INFO

Article history:

Received 15 October 2021

Revised 6 January 2022

Accepted 14 February 2022

Key Words:

Cheatgrass

Dormant-season grazing

Grazing management

Invasive annual grass

Selective grazing

ABSTRACT

Exotic annual grass invasion and dominance of sagebrush-bunchgrass steppe is a concern because it decreases biodiversity and promotes frequent wildfires. Management is needed to reduce exotic annual grasses to prevent sagebrush-bunchgrass communities from transitioning to annual grasslands. Grazing during the off season (fall-winter) has shown promise at reducing exotic annual grasses, but it has not been evaluated in plant communities dominated by sagebrush and native bunchgrasses. We compared moderate grazing during the off season with not grazing in five Wyoming big sagebrush (*Artemisia tridentata* ssp. *wyomingensis* [Beetle & A. Young] S.L. Welsh)–bunchgrass communities in the northern Great Basin. Treatments were applied annually for 10 yr (2009–2010 through 2018–2019). Plant community characteristics were measured after treatments had been applied from 6 to 10 yr. Off-season grazing reduced exotic annual grass density and cover. After a decade, annual grass cover was twofold greater in ungrazed areas. Sandberg bluegrass (*Poa secunda* J. Presl) density increased with off-season grazing, but large bunchgrass density was similar between off-season grazed and ungrazed areas. Perennial and annual forb density and cover were similar between off-season grazed and ungrazed treatments. Biological soil crust cover was also similar between off-season grazed and ungrazed areas. The results of this study provide strong evidence that off-season grazing has application for managing exotic annual grasses in sagebrush-bunchgrass steppe. Considering the vast scope of the exotic annual grass problem, properly applied grazing may be the most cost-efficient tool to mediate the impacts of annual grass invasion.

Published by Elsevier Inc. on behalf of The Society for Range Management.

This is an open access article under the CC BY-NC-ND license

(<http://creativecommons.org/licenses/by-nc-nd/4.0/>)

Introduction

Invasion of sagebrush steppe communities by exotic annual grasses is a major threat to ecological sustainability and rural economies. Millions of hectares of sagebrush communities are now exotic annual grasslands, and the area dominated by annual grasses is increasing at a rate of almost 200 000 ha annually (Smith et al. 2022). Annual grass invasion exponentially decreases biodiversity (Davies 2011) and degrades habitat for sagebrush associated wildlife (Knick et al. 2003; Crawford et al. 2004). Exotic annual grasses are highly competitive, especially with bunchgrass seedlings (Nasri and Doescher 1995; Rafferty and Young 2002), and

can deplete soil moisture and other resources, resulting in the exclusion of native plant species (Melgoza et al. 1990; Humphrey and Schupp 2004). Annual grass invasion is particularly devastating as it increases wildfire frequency (D'Antonio and Vitousek 1992; Balch et al. 2013). Annual grasses increase fine fuel amount and continuity and dry out earlier than native vegetation, thus elevating the probability of fire (Brooks et al. 2004; Davies and Nafus 2013). More frequent fire associated with annual grass invasion prevents reestablishment of fire intolerant native vegetation, resulting in annual grass monocultures and an exotic grass-fire cycle (D'Antonio and Vitousek 1992; Brooks et al. 2004; Eiswerth et al. 2009). Most of these annual grasslands are permanent community transitions because efforts to restore them to native perennial dominance are expensive and generally futile (Davies et al. 2021a).

A different management approach is needed to address the exotic annual grass problem with management intervention occurring before annual grass dominance. Many sagebrush-bunchgrass communities already have a noticeable presence of exotic annual grasses. Management of these plant communities needs to focus

[☆] Mention of a proprietary product does not constitute a guarantee or warranty of the product by the USDA or the authors and does not imply its approval to the exclusion of other products. The USDA is an equal opportunity provider and employer.

* Correspondence: Kirk Davies, EOARC, 67826-A Hwy 205, Burns, OR 97720, USA.
E-mail address: kirk.davies@usda.gov (K.W. Davies).

<https://doi.org/10.1016/j.rama.2022.02.003>

1550-7424/Published by Elsevier Inc. on behalf of The Society for Range Management. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>)

on reducing the competitive advantage of exotic annual grasses over native perennial vegetation (Perryman et al. 2018). An understory dominated by perennial bunchgrasses is probably the single most important assurance against exotic annual grass dominance (Chambers et al. 2007; Davies 2008; Davies and Johnson 2017). Therefore, management approaches should strive to limit exotic annual grasses with neutral to positive effects on perennial bunchgrasses. Prior efforts have focused on limiting disturbances, most notably excluding or greatly reducing grazing, in these communities. However, this has largely been ineffective, unless grazing management was improper (e.g., heavy, repeated use during the growing season). Thus, land management needs an approach that recognizes the complexity of ecosystem problems such as exotic annual grasses.

Land management needs to apply actions that reduce exotic annual grass density and cover, but these management actions need to be feasible to apply across the vast rangelands already invaded. Preemergent herbicides have shown great promise at reducing annual grasses and not negatively impacting established perennial bunchgrasses (Kyser et al. 2007; Sheley et al. 2007; Davies and Sheley 2011). Application of preemergent herbicides is expensive and, therefore, not feasible to apply to the extensive areas in need of exotic annual grass management (Stohlgren and Schnase 2006). In contrast, strategic grazing may be feasibly applied to expansive landscapes. Grazing by cattle can be an efficient treatment to reduce fuels across vast rangelands (Davies et al. 2015), so grazing may be a practical treatment to manage exotic annual grasses.

However, there is disagreement over the ability of grazing to reduce exotic annual grasses without negatively impacting native perennial vegetation. Heavy, repeated grazing during the growing season has promoted exotic annual grasses by decreasing competition from native perennial bunchgrasses (Daubenmire 1970 Mack 1981; Knapp 1996). Subsequently, some have recommended reducing livestock grazing to decrease exotic annual grasses (e.g., Reisner et al. 2013; Williamson et al. 2020), while others have found that grazing can reduce annual grasses and promote native perennials (Davies et al. 2020; Porensky et al. 2021). Reductions in grazing have also been suggested to be a contributing factor to the recent expansion of exotic annual grasslands (Perryman et al. 2018). Some of these contrasting results are likely tied to differences in grazing management, and thus, specific, experimental applied grazing needs to be investigated.

Off-season (fall–winter) grazing is increasingly becoming a common grazing management alternative to growing season use in Wyoming big sagebrush communities, in part to reduce accumulations of fine fuels to decrease fire probability and also to decrease the cost associated with feeding hay. Moderate grazing in the off-season may decrease exotic annual grasses and favor native perennial vegetation but is an understudied grazing management strategy. Off-season grazing may decrease annual grasses by removing ground litter (usually annual grass litter). Litter is critical to annual grass success because it creates a favorable microenvironment for its emergence and growth (Evans and Young 1970; Facelli and Pickett 1991; Newingham et al. 2007; Wolkovich et al. 2009). Off-season grazing may also result in cattle selectively grazing exotic annual grasses because they can initiate growth earlier than native perennial grasses (Smith et al. 2012; Davies et al. 2021b). After a substantial postfire invasion of annual grasses, off-season grazing reduced annual grasses and increased Sandberg bluegrass (*Poa secunda* J. Presl), a native bunchgrass (Davies et al. 2021b). In a case study, off-season grazing reduced annual grasses and increased an introduced bunchgrass in an exotic annual grass–introduced bunchgrass community (Schmelzer et al. 2014). However, the effects of moderate grazing during the off season on unburned Wyoming big sagebrush communities with native bunchgrass-dominated understories are unknown.

The objective of this study was to investigate the effects of moderate grazing during the off-season on exotic annual grasses and native vegetation in Wyoming big sagebrush–bunchgrass communities that are moderately invaded by annual grasses. To accomplish this objective, we compared areas that for the past 6–10 yr were off-season grazed or not grazed. We hypothesized that 1) moderate grazing in the off-season would reduce exotic annual grass cover and density and, subsequently, 2) increase the density of native bunchgrasses compared with the ungrazed treatment. Native bunchgrass cover was not expected to be greater in grazed areas because off-season grazing would remove prior years' growth, which contributes substantially to bunchgrass cover.

Methods

Study area

This study was conducted in Wyoming sagebrush–bunchgrass steppe 65–70 km south and southeast of Burns, Oregon (43°04'N, 118°40'W). These sagebrush–bunchgrass communities have low resilience and resistance to annual grass invasion (Chambers et al. 2014a, 2014b). The understory in these communities was dominated by native perennial bunchgrasses, either Thurber's needlegrass (*Achnatherum thurberianum* [Piper] Barkworth) or bluebunch wheatgrass (*Pseudoroegneria spicata* [Pursh] A. Löve). Other common bunchgrasses included Sandberg bluegrass and bottlebrush squirreltail (*Elymus elymoides* [Raf.] Swezey). Exotic annual grasses were observed across the study sites before study initiation. Average (1981–2010) annual precipitation across the study area was 280 mm (PRISM 2020). Precipitation patterns are representative of the northern Great Basin with most events occurring in the winter and spring, and summers are hot and dry. Study sites were generally flat, and elevation was approximately 1 450 m. Study sites were on Sandy Loam 10–12 PZ (R023XY213OR) and Droughty Loam 11–13 PZ (R023XY316OR) Ecological Sites. Historical fire occurrence for these steppe communities was infrequent with fire return intervals estimated to be 50–100+ yr (Wright and Bailey 1982; Mensing et al. 2006). We could find no evidence or record of recent (~50 yr) fire at the study sites.

Experimental design and measurements

We used a randomized complete block design with two treatments: grazed (moderate, off-season grazing) and ungrazed (not grazed since 2009). Treatments were applied to five blocks (sites) that varied in site and vegetation characteristics and randomly assigned to plots in each block. Vegetation characteristics were generally similar between treatments within blocks before treatment initiation (Davies et al. 2016a). The study was initiated in the fall of 2009 by building 60 × 100 m barbed wire enclosures to exclude grazing from the ungrazed treatment. The grazed and ungrazed treatments were separated by a 20-m buffer. Grazing by cattle was applied as part of the normal management (initiated at the start of the study) of the pastures containing enclosures between November and early April each year. Pastures were ~800–1 000 ha and located adjacent to one another. Cattle consumed 40–60% of the available forage based on biomass determined by the method described in Anderson and Curreir (1973). Cattle were provided a protein supplement to ensure their nutritional needs were met during the off-season. Sampling was not initiated until 2015 to allow vegetation time to respond to treatments. Treatments were applied annually for the duration of the study.

Vegetation characteristics were measured in mid to late June of each yr from 2015 through 2019. Four, parallel 45-m transects spaced at 5-m intervals were established in each treatment replicate to measure vegetation characteristics. Herbaceous vegetation




Figure 1. Exotic annual grass, Sandberg bluegrass, and large perennial bunchgrass cover and density (mean \pm standard of error) in moderate off-season grazed (grazed) and ungrazed (ungrazed) treatments from 2015 to 2019.

cover was visually estimated by species in 0.2-m² quadrats located at 3-m intervals on each 45-m transect, starting at the 3-m mark and ending at the 45-m mark (15 quadrats·transect⁻¹ and 60 quadrats·treatment replicate⁻¹). Bare ground, biological soil crust, and litter cover were also estimated in the 0.2-m² quadrats. Herbaceous vegetation density was measured by species by counting each plant rooted in the 0.2-m² quadrats. Sagebrush cover was measured using the line-intercept method along each 45-m transect. Sagebrush density was measured by placing a 2 \times 45 m belt transect over each 45-m transect and counting all individuals rooted in it.

Statistical analyses

To compare treatments, we used a repeated-measures analysis of variance using the mixed model procedure (SAS v. 9.4, SAS Institute, Cary, NC). Block and block-by-treatment interactions were

treated as random variables in analyses with year being the repeated variable. Treatment was treated as a fixed variable in analyses. The Akaike information criterion was used to select the appropriate covariance structure for analyses (Littell et al. 1996). Data that violated assumptions of analysis of variance were log-transformed. For analyses, herbaceous vegetation was separated into plant functional groups: exotic annual grasses, Sandberg bluegrass, large perennial bunchgrasses, perennial forbs, and annual forbs. The exotic annual grass group consisted primarily of cheatgrass (*Bromus tectorum* L.) with some medusahead (*Taeniatherum caput-medusae* [L.] Nevski). Sandberg bluegrass was analyzed independent of the other bunchgrasses because it generally develops phenologically earlier, is smaller in stature, and responds differently to grazing and fire (McLean and Tisdale 1972; Yensen et al. 1992; Davies et al. 2021b). The perennial bunchgrass and perennial forb groups were composed solely of native species. The annual forb group consisted of non-native and native species. All fig-

ures and text present nontransformed data. Means were reported with standard errors in the text and figures with statistical significance set at $P \leq 0.05$. Year-by-treatment interactions were included in all models, but only significant treatment-by-year interactions were reported in the text.

Results

Exotic annual grass cover was reduced with grazing and varied among years (Fig. 1A; $P < 0.001$ and < 0.001). At the final sampling, annual grass cover was twofold greater in the ungrazed compared with the grazed treatment. Density of exotic annual grasses was relatively similar the first sampling year but thereafter was substantially greater in ungrazed areas (Fig. 1B; $P < 0.001$). On average, annual grass density was 1.6-fold greater in ungrazed compared with grazed areas. Sandberg bluegrass cover was similar between treatments (Fig. 1C; $P=0.372$) but varied among years ($P < 0.001$). Sandberg bluegrass density was greater in the grazed compared with the ungrazed treatment (Fig. 1D; $P=0.002$) and varied among years ($P=0.008$). Averaged across sample years, Sandberg bluegrass density was 1.4-fold greater in the grazed compared with the ungrazed treatment. Large perennial bunchgrass cover was reduced with grazing (Fig. 1E; $P=0.005$) and varied among years ($P < 0.001$). In the final sampling year, perennial bunchgrass cover was 1.6-fold greater in the ungrazed compared with the grazed treatment. In contrast, large perennial bunchgrass density was similar between grazed and ungrazed areas (see Fig. 1F; $P=0.943$). Large perennial bunchgrass density did not vary among years ($P=0.137$). Perennial forb cover and density did not differ between grazed and ungrazed treatments (Fig. 2A and 2B; $P=0.612$ and 0.993) but varied among years ($P < 0.001$). Annual forb cover and density were also similar between treatments (Fig. 2C and 2D; $P=0.954$ and $=0.79$) but varied among years ($P < 0.001$). Sagebrush cover did not differ between treatments (Fig. 2E; $P=0.215$) but varied among years ($P=0.046$). Sagebrush cover appeared to generally increase with time in both grazed and ungrazed areas. Sagebrush density was similar in the grazed and ungrazed treatment (Fig. 2F; $P=0.785$) and did not vary among years ($P=0.141$). Litter cover on the soil surface was similar between the grazed and ungrazed treatment (Fig. 3A; $P=0.260$) but varied among years ($P < 0.001$). Bare ground was greater in the grazed compared with the ungrazed treatment (Fig. 3B; $P=0.036$). At the conclusion of the study, bare ground was 1.3-fold greater in grazed compared with ungrazed areas. Bare ground varied among years ($P < 0.001$), but no clear pattern emerged. Biological soil crust cover did not differ between grazed ($1.2\% \pm 0.2\%$) and ungrazed ($1.5\% \pm 0.2\%$) areas ($P=0.538$). Biological soil crust cover varied among years ($P < 0.001$), but no trend was apparent.

Discussion

Our hypotheses were largely supported by the response of Wyoming big sagebrush-bunchgrass communities to 6–10 yr of moderate grazing in the off-season compared with not being grazed. Off-season grazing reduced the density and cover of exotic annual grasses in Wyoming big sagebrush-bunchgrass communities in support of our first hypothesis. Partially supporting our second hypothesis that off-season grazing would increase native bunchgrass density, Sandberg bluegrass density was greater in grazed compared with ungrazed areas. However, large native bunchgrasses density was similar between grazed and ungrazed treatments. The response of annual grasses and Sandberg bluegrass aligned with our hypotheses, but the large bunchgrass response was not as expected.

Exotic annual grass cover and density were likely reduced with off-season grazing because cattle may selectively graze it during

this time period and grazing may reduce safe sites for its emergence and growth. Annual grass may grow in the fall or late winter/early spring before native perennial grasses (Hironaka 1961; Harris 1977). This early growth of annual grasses may result in livestock selectively grazing it because other vegetation may be dormant or producing little accessible new growth (Smith et al. 2012; Davies et al. 2021b). Annual grass success is also expected to be less in grazed areas because grazing decreases safe sites for its emergence and growth by reducing herbaceous vegetation and litter cover (Davies et al. 2021b). In our current study, off-season grazing reduced large perennial bunchgrass cover, which may have contributed to the decline in annual grasses. Exotic annual grass success declined with reduced vegetation cover in a study in Colorado, likely because vegetation cover, similar to litter, creates a favorable microenvironment for the establishment of annual grass (Adair et al. 2008). Litter covering the soil surface improves annual grass emergence and growth because of its effects on microsites (Evans and Young 1970; Facelli and Pickett 1991; Newingham et al. 2007; Wolkovich et al. 2009) and nutrient cycling (Booth et al. 2003; Sperry et al. 2006). However, we did not find reduced litter cover with off-season grazing. Potentially, grazing reduced the litter biomass and thickness and, thus, negatively impacted annual grasses (Davies et al. 2021b). However, we did not investigate if litter biomass or thickness was influenced by off-season grazing. Bare ground was greater with off-season grazing, suggesting that favorable microsites for exotic annual grasses were reduced with this treatment, even without finding evidence that litter cover was reduced with grazing. Thus, we speculate that off-season grazing reduced annual grass density and cover through the physical effects of grazing it, as well as modifying the microenvironment for its emergence and growth.

The results from our current study provide additional evidence that off-season grazing may be a tool for reducing exotic annual grasses. Similar to our results, annual grass standing crop was reduced with fall grazing in an exotic annual grass–introduced bunchgrass community in Nevada (Schmelzer et al. 2014). Fall grazing in this community also reduced the annual grass seed bank (Schmelzer et al. 2014; Perryman et al. 2020). Off-season grazing also reduced exotic annual grass density and cover in sagebrush-bunchgrass steppe communities substantially invaded by annual grass after wildfire removed sagebrush (Davies et al. 2021b). Our study is the first to find that off-season grazing compared with not grazing can decrease exotic annual grass density and cover in sagebrush-bunchgrass–dominated plant communities. These results combined with prior studies (Schmelzer et al. 2014; Perryman et al. 2020; Davies et al. 2021b) suggest that off-season grazing can be used to reduce exotic annual grasses in plant communities with varying species and plant functional group composition.

The increased density of the native bunchgrass, Sandberg bluegrass, suggests that off-season grazing may be opening safe sites for native perennials by reducing exotic annual grasses. Similarly, Sandberg bluegrass increased with reductions in exotic annual grass with off-season (Davies et al. 2021b) and spring grazing (Davies et al. 2020) in recently burned sagebrush communities. Reductions in annual grasses likely favored Sandberg bluegrass because annual grasses are highly competitive and can exclude native vegetation (Melgoza et al. 1990; Nasri and Doescher 1995; Rafferty and Young 2002). Sandberg bluegrass may also be negatively impacted by shading from annual grasses because of its relatively short stature. Therefore, when grazing reduces exotic annual grasses, other vegetation may increase because of less competition and shading.

Large perennial bunchgrass density did not increase with reductions in annual grasses with off-season grazing. This outcome is similar to a prior study where off-season grazing applied after fire in an exotic annual grass–native bunchgrass community did




Figure 2. Annual forb, perennial forb, and Wyoming big sagebrush cover and density (mean \pm standard of error) in moderate off-season grazed (grazed) and ungrazed (ungrazed) treatments from 2015 to 2019.

not increase large bunchgrass density (Davies et al. 2021b) but dissimilar to a study in Nevada where bunchgrass, largely composed of an introduced species, standing crop increased with fall grazing (Schmelzer et al. 2014). We did not measure large perennial bunchgrass standing crop in our current study but doubt it increased with grazing since large perennial bunchgrass cover was less in the grazed treatment. This was expected as grazing would reduce prior years' growth, a major contributor to both cover and standing crop of large perennial bunchgrasses (Davies et al. 2009, 2016b). There are a couple of possible reasons why large perennial bunchgrass density did not increase with the grazing-induced reduction in exotic annual grasses. First, recruitment of large native bunchgrasses is infrequent in arid and semiarid environments (Holmes and Rice 1996; James et al. 2009; Svejcar et al. 2017) and would need to be greater than mortality to see an increase in density. Second, the sagebrush steppe–bunchgrass communities included in this study may be near capacity for large perennial bunchgrass

density. The off-season grazed and ungrazed treatments averaged 9.0 ± 0.87 and 8.9 ± 0.42 large perennial bunchgrasses·m⁻², respectively. The average density for large perennial bunchgrasses in intact Wyoming big sagebrush–bunchgrass steppe in this region is ~ 10 individuals·m⁻² (Davies and Bates 2010; Bates and Davies 2019). Thus, large perennial bunchgrass density may not yet be affected by the limited exotic annual grass invasion in these communities. Regardless, off-season grazing for 6–10 yr did not increase large perennial bunchgrass density in sagebrush–bunchgrass communities with already high abundance of large bunchgrasses. Importantly, large bunchgrass density didn't decrease with grazing, suggesting that off-season grazing is compatible with bunchgrass conservation goals.

The differing response of Sandberg bluegrass and large bunchgrasses to off-season grazing is likely a product of their differences in stature and response to disturbances. Sandberg bluegrass may be especially sensitive to annual grass competition because it




Figure 3. Litter and bare ground cover (mean \pm standard of error) in moderate off-season grazed (grazed) and ungrazed (ungrazed) treatments from 2015 to 2019.

usually grows in the interspaces between large perennial bunchgrasses, the location where annual grasses proliferate during invasion (Reisner et al. 2013; Rayburn et al. 2014). In the current study, where annual grasses were not dominant and large bunchgrasses plentiful, reductions in annual grass abundance likely opened safe sites for Sandberg bluegrass, but not necessarily for large bunchgrasses. Sandberg bluegrass may also increase with grazing as it can be difficult for cattle to graze its short leaves (Davies et al. 2021b) and it is considered grazing tolerant because it will increase with heavy spring grazing when other native bunchgrasses decline (McLean and Tisdale 1972). Thus, the effects of off-season grazing likely differ by community characteristics, such as level of invasion, and possibly vary among species and plant functional groups.

Management Implications

Off-season grazing can be used to reduce exotic annual grass density and cover in Wyoming big sagebrush–bunchgrass communities without compromising the native bunchgrass component. This reduction in annual grasses can facilitate increases in Sandberg bluegrass density. However, off-season grazing does not appear to increase the density of other native bunchgrasses. This treatment may not benefit large bunchgrasses, or these communities may be at or near capacity for large bunchgrass abundance. Regardless, this study demonstrates that increases in the density of large bunchgrasses should not be expected with off-season grazing in Wyoming big sagebrush–bunchgrass communities with abundant large bunchgrasses. Prior research has shown that off-season grazing can be used to reduce exotic annual grasses in annual grass–native bunchgrass and annual grass–introduced bunchgrass communities (Schmelzer et al. 2014; Perryman et al. 2020; Davies et al. 2021b). The results of these prior studies and our current study suggest that off-season grazing by cattle may be a valuable tool for managing the expanding exotic annual grass problem. Further research evaluating the effects of off-season grazing is warranted. Investigating the effects of off-season grazing on restoration efforts may prove fruitful as success is frequently hindered by exotic annual grasses (Svejcar et al. 2017). Research into cattle performance, supplementation strategies, economics, and management contingencies for adverse weather are also needed to improve the application of off-season grazing. Finally, evaluations of the effects of off-season grazing on wildlife are also needed to ensure off-season grazing is applied in a manner that is compatible with species of conservation concern and to develop wildlife friendly management practices when and where they are needed.

Declaration of Competing Interest

We do not have any conflict of interest to declare.

Acknowledgments

We are grateful to the private landowners for allowing this research project to occur on their lands and applying the grazing treatment. We also thank Aleta Nafus, Urban Strachan, and numerous summer students for setting up the experiment and data collection. We appreciate thoughtful reviews of earlier versions of this manuscript by Barry Perryman and Sergio Arispe.

References

- Adair, E.C., Burke, I.C., Lauenroth, W.K., 2008. Contrasting effect so of resource availability and plant mortality on plant community invasion by *Bromus tectorum* L. Plant and Soil 304, 103–115.
- Anderson, E.W., Curreir, W.F., 1973. Evaluating zones of utilization. Journal of Range Management 26, 87–91.
- Balch, J.K., Bradley, B.A., D'Antonio, C.M., 2013. Introduced annual grass increases regional fire activity across the arid western USA (1980–2009). Global Change Biology 19, 173–183.
- Bates, J.D., Davies, K.W., 2019. Characteristics of intact Wyoming big sagebrush associations in southeastern Oregon. Rangeland Ecology & Management 72, 36–46.
- Booth, M.S., Caldwell, M.M., Stark, J.M., 2003. Overlapping resource use in three Great Basin species: implications for community invasibility and vegetation dynamics. Journal of Ecology 91, 36–48.
- Brooks, M.L., D'Antonio, C.M., Richardson, D.M., Grace, J.B., Keeley, J.E., DiTomaso, J.M., Hobbs, R.J., Pellant, M., Pyke, D., 2004. Effects of invasive alien plants on fire regimes. Bioscience 54, 677–688.
- Chambers, J.C., Bradley, B.A., Brown, C.S., D'Antonio, C., Germino, M.J., Grace, J.B., Hardegree, S.P., Miller, R.F., Pyke, D.A., 2014a. Resilience to stress and disturbance and resistance to *Bromus tectorum* L. invasion in cold desert shrublands of western North America. Ecosystems 17, 360–375.
- Chambers, J.C., Miller, R.F., Board, D.I., Pyke, D.A., Rund, B.A., Brace, J.B., Schupp, E.W., Tausch, R.J., 2014b. Resilience and resistance of sagebrush ecosystems: implications for state and transition models and management treatments. Rangeland Ecology & Management 67, 440–454.
- Chambers, J.C., Roundy, B.A., Blank, R.R., Meyer, S.E., Whittaker, A., 2007. What makes Great Basin sagebrush ecosystems invulnerable by *Bromus tectorum*? Ecological Monographs 77, 117–145.
- Crawford, J.A., Olson, R.A., West, N.E., Mosley, J.C., Schroeder, M.A., Whitson, T.D., Miller, R.F., Gregg, M.A., Boyd, C.S., 2004. Ecology and management of sage-grouse and sage-grouse habitat. Journal of Range Management 57, 2–19.
- D'Antonio, C.M., Vitousek, P.M., 1992. Biological invasions by exotic grasses, the grass/fire cycle, and global change. Annual Review of Ecology and Systematics 23, 63–87.
- Daubenmire, R., 1970. Steppe vegetation of Washington, 62. Washington Agricultural Experiment Station Technical Bulletin, Pullman, WA, USA, p. 131.
- Davies, K.W., 2008. Medusahead dispersal and establishment in sagebrush steppe plant communities. Rangeland Ecology & Management 61, 110–115.
- Davies, K.W., 2011. Plant community diversity and native plant abundance decline with increasing abundance of an exotic annual grass. Oecologia 167, 481–491.

- Davies, K.W., Bates, J.D., 2010. Vegetation characteristics of mountain and Wyoming big sagebrush plant communities in the northern Great Basin. *Rangeland Ecology & Management* 63, 461–466.
- Davies, K.W., Bates, J.D., Boyd, C.S., 2016b. Effects of intermediate-term grazing rest on sagebrush communities with depleted understories: evidence of a threshold. *Rangeland Ecology & Management* 69, 173–178.
- Davies, K.W., Bates, J.D., Boyd, C.S., 2020. Response of planted sagebrush seedlings to cattle grazing applied to reduce fine fuels to decrease fire probability. *Rangeland Ecology & Management* 73, 629–635.
- Davies, K.W., Bates, J.D., Perryman, B., Arispe, S., 2021b. Fall-winter grazing after fire in annual grass-invaded sagebrush steppe reduced annuals and increased a native bunchgrass. *Rangeland Ecology & Management* 77, 1–8.
- Davies, K.W., Boyd, C.S., Bates, J.D., Hulet, A., 2015. Dormant-season grazing may decrease wildfire probability by increasing fuel moisture and reducing fuel amount and continuity. *International Journal of Wildland Fire* 24, 849–856.
- Davies, K.W., Johnson, D.D., 2017. Established perennial vegetation provides high resistance to reinvasion by exotic annual grasses. *Rangeland Ecology & Management* 70, 748–754.
- Davies, K.W., Leger, E.A., Boyd, C.S., Hallett, L.M., 2021a. Living with exotic annual grasses in the sagebrush ecosystem. *Journal of Environmental Management* 288, 112417.
- Davies, K.W., Nafus, A.M., 2013. Exotic annual grass invasion alters fuel amounts, continuity and moisture content. *International Journal of Wildland Fire* 22, 353–358.
- Davies, K.W., Nafus, A.M., Boyd, C.S., Hulet, A., Bates, J.D., 2016a. Effects of using winter grazing as a fuel treatment on Wyoming big sagebrush plant communities. *Rangeland Ecology & Management* 69, 179–184.
- Davies, K.W., Sheley, R.L., 2011. Promoting native vegetation and diversity in exotic annual grass infestations. *Restoration Ecology* 19, 159–165.
- Davies, K.W., Svejcar, T.J., Bates, J.D., 2009. Interaction of historical and nonhistorical disturbances maintains native plant communities. *Ecological Applications* 19, 1536–1545.
- Eiswerth, M.E., Krauter, K., Swanson, S.R., Zielinski, M., 2009. Post-fire seeding on Wyoming big sagebrush ecological sites: regression analyses of seeded non-native and native species densities. *Journal of Environmental Management* 90, 1320–1325.
- Evans, R.A., Young, J.A., 1970. Plant litter and establishment of alien annual weed species in rangeland communities. *Weed Science* 18, 697–703.
- Facelli, J.M., Pickett, S.T.A., 1991. Plant litter: its dynamics and effects on plant community structure. *Botanical Review* 57, 1–32.
- Harris, G.A., 1977. Root phenology as a factor of competition among grass seedlings. *Journal of Range Management* 30, 172–177.
- Hironaka, M., 1961. The relative rate of root development of cheatgrass and medusa-head. *Journal of Range Management* 14, 263–267.
- Holmes, T.H., Rice, K.J., 1996. Patterns of growth and soil-water utilization in some exotic annuals and native perennial bunchgrasses of California. *Annals of Botany* 78, 233–243.
- Humphrey, L.D., Schupp, E.W., 2004. Competition as a barrier to establishment of a native perennial grass (*Elmus elymoides*) in alien annual grass (*Bromus tectorum*) communities. *Journal of Arid Environments* 58, 405–422.
- James, J.J., Mangold, J.M., Sheley, R.L., Svejcar, T., 2009. Root plasticity of native and invasive Great Basin species in response to soil nitrogen heterogeneity. *Plant Ecology* 202, 211–220.
- Knick, S.T., Dobkins, D.S., Rothenberry, J.T., Schroeder, M.A., Vander Haegen, W.M., Van Riper, C., 2003. III. Teetering on the edge or too late: conservation and research issues for avifauna of sagebrush habitats. *Condor* 105, 611–634.
- Knapp, P.A., 1996. Cheatgrass (*Bromus tectorum* L.) dominance in the Great Basin Desert: history, persistence, and influences to human activities. *Global Environmental Change* 6, 37–52.
- Kyser, G.B., DiTomaso, J.M., Doran, M.P., Orloff, S.B., Wilson, R.G., Lancaster, D.L., Lile, D.F., Porath, M.L., 2007. Control of medusahead (*Taeniatherum caput-medusae*) and other annual grasses with imazapic. *Weed Technology* 21, 66–75.
- Littell, R.C., Milliken, G.A., Stroup, W.W., Wolfinger, R.D., 1996. SAS System for Mixed Models. SAS Institute Inc. 633 p, Cary, NC, USA.
- Mack, R.N., 1981. Invasion of *Bromus tectorum* L. into western North America: an ecological chronicle. *Agro-Ecosystems* 7, 145–165.
- McLean, A., Tisdale, E.W., 1972. Recovery rate of depleted range sites under protection from grazing. *Journal of Range Management* 25, 178–184.
- Melgoza, G., Nowak, R.S., Tausch, R.J., 1990. Soil-water exploitation after fire—competition between *Bromus tectorum* (cheatgrass) and 2 native species. *Oecologia* 83, 7–13.
- Mensing, S., Livingston, S., Barker, P., 2006. Long-term fire history in Great Basin sagebrush reconstructed from macroscopic charcoal in spring sediments, Newark Valley, Nevada. *Western North American Naturalist* 66, 64–77.
- Nasri, M., Doescher, P.S., 1995. Effect of competition by cheatgrass on shoot growth of Idaho fescue. *Journal of Range Management* 48, 402–405.
- Newingham, B.A., Vidiella, P., Belnap, J., 2007. Do soil characteristics or microhabitat determine field emergence and success of *Bromus tectorum*? *Journal of Arid Environments* 70, 389–402.
- Perryman, B.L., Schultz, B.W., McAdoo, J.K., Alvert, R.L., Cervantes, J.C., Foster, S., McCuin, G., Swanson, S., 2018. Viewpoint: an alternative management paradigm for plant communities affected by invasive annual grass in the Intermountain West. *Rangelands* 40, 77–82.
- Perryman, B.L., Schultz, B.W., Burrows, M., Shenkoru, T., Wilker, J., 2020. Fall-grazing and grazing-exclusion effects on cheatgrass (*Bromus tectorum*) seed bank assays in Nevada, United States. *Rangeland Ecology & Management* 73, 343–347.
- Porensky, L.M., Baughman, O., Williamson, M.A., Perryman, B.L., Madsen, M.D., Leger, E.A., 2021. Using native grass seeding and targeted spring grazing to reduce low-level *Bromus tectorum* invasion on the Colorado Plateau. *Biological Invasions* 23, 705–722.
- PRISM. 2020. PRISM Climatic Group. Available at: <http://prism.nacse.org/explorer/>. Accessed April 17, 2020.
- Rafferty, D.L., Young, J.A., 2002. Cheatgrass competition and establishment of desert needlegrass seedlings. *Journal of Range Management* 55, 70–72.
- Rayburn, A.P., Schupp, E.W., Kay, S., 2014. Effects of perennial semi-arid bunchgrass spatial patterns on performance of the invasive annual cheatgrass (*Bromus tectorum* L.). *Plant Ecology* 215, 247–251.
- Reisner, M.D., Grace, J.B., Pyke, D.A., Doescher, P.S., 2013. Conditions favouring *Bromus tectorum* dominance of endangered sagebrush steppe ecosystems. *Journal of Applied Ecology* 50, 1039–1049.
- Schmelzer, L., Perryman, B., Bruce, B., Schultz, B., McAdoo, K., McCuin, G., Swanson, S., Wilker, J., Conley, K., 2014. Case study: reducing cheatgrass (*Bromus tectorum* L.) fuel loads using fall cattle grazing. *Professional Animal Scientist* 30, 270–278.
- Sheley, R.L., Carpinelli, M.F., Morghan, K.J.R., 2007. Effects of imazapic on target and nontarget vegetation during revegetation. *Weed Technology* 21, 1071–1081.
- Smith, B., Sheley, R., Svejcar, T., 2012. Grazing invasive annual grass: the green and brown guide. Eastern Oregon Agricultural Research Center, Burns, OR, USA, p. 36.
- Smith, J.T., Allred, B.W., Boyd, C.S., Davies, K.W., Jones, M.O., Maestas, J.D., Morford, S.L., Naugle, D.E., 2022. The elevation ascent and spread of exotic annual grasslands in the Great Basin, USA. *Diversity and Distributions* 28, 83–96.
- Sperry, L.J., Belnap, J., Evans, R.D., 2006. *Bromus tectorum* invasion alters nitrogen dynamics in an undisturbed arid grassland ecosystem. *Ecology* 87, 603–615.
- Stohlgren, T.J., Schnase, J.L., 2006. Risk analysis for biological hazards; what we need to know about invasive species. *Risk Analysis* 26, 163–173.
- Svejcar, T., Boyd, C., Davies, K., Hamerlynck, E., Svejcar, L., 2017. Challenges and limitations to native species restoration in the Great Basin, USA. *Plant Ecology* 218, 81–94.
- Williamson, M.A., Fleishman, E., MacNally, R.C., Chambers, J.C., Bradley, B.A., Dobkin, D.S., Board, D.I., Fogarty, F.A., Horning, N., Leu, M., Zillig, M.W., 2020. Fire, livestock grazing, topography, and precipitation affect occurrence and prevalence of cheatgrass (*Bromus tectorum*) in the central Great Basin, USA. *Biological Invasions* 22, 663–680.
- Wolkovich, E.M., Bolger, D.T., Cottingham, K.L., 2009. Invasive grass litter facilitates native shrubs through abiotic effects. *Journal of Vegetation Science* 20, 1121–1132.
- Wright, H.A., Bailey, A.W., 1982. Fire ecology: United States and southern Canada. John Wiley & Sons, Inc. 496 p, New York, NY, USA.
- Yensen, E., Quinney, D.L., Johnson, K., Timmerman, K., Steenhof, K., 1992. Fire, vegetation changes, and population fluctuations of Townsend's ground squirrels. *American Midland Naturalist* 128, 299–312.