

## **When using glyphosate plus dicamba, 2,4-D, halauxifen or pyraflufen/2,4-D for glyphosate-resistant horseweed (*Erigeron canadensis*) control in soybean, which third mix partner is better, saflufenacil or metribuzin?**

Authors: Dilllott, Meghan, Soltani, Nader, Hooker, David C., Robinson, Darren E., and Sikkema, Peter H.

Source: Weed Technology, 36(2) : 295-302

Published By: Weed Science Society of America

URL: <https://doi.org/10.1017/wet.2022.18>

---

BioOne Complete ([complete.BioOne.org](https://complete.bioone.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

## Research Article

**Cite this article:** Dilllotti M, Soltani N, Hooker DC, Robinson DE, Sikkema PH (2022) When using glyphosate plus dicamba, 2,4-D, halauxifen or pyraflufen/2,4-D for glyphosate-resistant horseweed (*Erigeron canadensis*) control in soybean, which third mix partner is better, saflufenacil or metribuzin? Weed Technol. **36**: 295–302. doi: [10.1017/wet.2022.18](https://doi.org/10.1017/wet.2022.18)

Received: 29 October 2021

Revised: 8 February 2022

Accepted: 25 February 2022

First published online: 15 March 2022

**Associate Editor:**

Vipin Kumar, Kansas State University

**Nomenclature:**

2,4-D ester; dicamba; glyphosate; halauxifen-methyl; pyraflufen-ethyl/2,4-D; metribuzin; saflufenacil; horseweed; *Erigeron canadensis* L.; soybean; *Glycine max* (L.) Merr.



**Keywords:**

Additive; antagonistic; biomass; density; synergy; weed control; grain yield

**Author for correspondence:**

Nader Soltani, Department of Plant Agriculture, University of Guelph, Ridgetown Campus, 120 Main Street East, Ridgetown, ON, Canada N0P 2C0. Email: [soltani@uoguelph.ca](mailto:soltani@uoguelph.ca)

# When using glyphosate plus dicamba, 2,4-D, halauxifen or pyraflufen/2,4-D for glyphosate-resistant horseweed (*Erigeron canadensis*) control in soybean, which third mix partner is better, saflufenacil or metribuzin?

Meghan Dilllotti<sup>1</sup> , Nader Soltani<sup>2</sup> , David C. Hooker<sup>3</sup>, Darren E. Robinson<sup>4</sup> and Peter H. Sikkema<sup>5</sup>

<sup>1</sup>Graduate Student, Department of Plant Agriculture, University of Guelph, Ridgetown, ON, Canada; <sup>2</sup>Adjunct Professor, Department of Plant Agriculture, University of Guelph, Ridgetown, ON, Canada; <sup>3</sup>Associate Professor, Department of Plant Agriculture, University of Guelph, Ridgetown, ON, Canada; <sup>4</sup>Professor, Department of Plant Agriculture, University of Guelph, Ridgetown, ON, Canada and <sup>5</sup>Professor, Department of Plant Agriculture, University of Guelph, Ridgetown, ON, Canada

**Abstract**

Glyphosate-resistant (GR) horseweed interference in soybean can reduce soybean yield up to 93%. Glyphosate plus dicamba, 2,4-D ester, halauxifen-methyl or pyraflufen-ethyl/2,4-D applied preplant (PP) provide variable GR horseweed control in soybean. The objective of this study was to determine if the addition of saflufenacil or metribuzin to glyphosate plus dicamba, 2,4-D ester, halauxifen-methyl, or pyraflufen-ethyl/2,4-D will improve the level and consistency of GR horseweed control. Four trials were conducted over the 2020 and 2021 field seasons in fields with GR horseweed populations. Glyphosate plus dicamba, 2,4-D ester, halauxifen-methyl, or pyraflufen-ethyl/2,4-D controlled GR horseweed 96%, 77%, 71%, and 52%, respectively, at 8 wk after application (WAA). When saflufenacil or metribuzin was added to glyphosate plus dicamba or 2,4-D ester, GR horseweed control was not improved at 8 WAA. When saflufenacil or metribuzin was added to glyphosate plus halauxifen-methyl, GR horseweed control improved by 27% and 25%, respectively, at 8 WAA. When saflufenacil or metribuzin was added to glyphosate plus pyraflufen-ethyl/2,4-D, GR horseweed control was improved by 47% and 37%, respectively, at 8 WAA. The consistency of GR horseweed control was improved when saflufenacil or metribuzin was added to glyphosate plus dicamba, 2,4-D ester, halauxifen-methyl, or pyraflufen-ethyl/2,4-D compared to each herbicide applied alone. Synergism was observed when metribuzin was added to glyphosate plus halauxifen-methyl and when saflufenacil or metribuzin was added to glyphosate plus pyraflufen-ethyl/2,4-D at 8 WAA. Though GR horseweed control was improved with the addition of saflufenacil or metribuzin to glyphosate plus halauxifen-methyl or pyraflufen-ethyl/2,4-D, all treatments including saflufenacil resulted in the highest level and most consistent control.

**Introduction**

Horseweed is a broadleaf weed from the Asteraceae family (Weaver 2001). Horseweed has been found in the southern parts of all Canadian provinces excluding Newfoundland (Weaver 2001). It is now considered a cosmopolitan weed that is found most commonly in the north temperate geographical zone (Holm et al. 1997). The first glyphosate-resistant (GR) horseweed biotype was discovered in a soybean field in the US state of Delaware in 2000 (VanGessel 2001). In Canada, the first report of GR horseweed was in Essex County, ON, in 2010 (Byker et al. 2013d). As of 2015, 30 Ontario counties have been confirmed with GR horseweed (Budd et al. 2017).

Horseweed is a facultative winter annual with nondormant seeds that can emerge in the fall or spring (Weaver 2001). Fall-emerged horseweed first establishes a basal rosette, and the stem elongates in the spring (Loux et al. 2006; Weaver 2001). One study observed that an individual GR horseweed plant produced nearly 200,000 seeds in a noncompetitive environment (Bhowmik and Bekech 1993). More recently, Davis et al. (2009) observed a multiple-resistant horseweed plant that produced 1 million seeds in a no-till field, following a 2,4-D application. The seeds are about 1 to 2 mm long with a pappus approximately 3 to 5 mm long (Frankton and Mulligan 1987). The pappus resembles a parachute, allowing the seed to undergo wind dispersal (Frankton and Mulligan 1987). Seeds have been collected hundreds of kilometers from the mother plant (Shields et al. 2006); however, most seeds fall within 100 m. Horseweed has a low outcrossing rate of 4% despite producing approximately 95,000 pollen grains per day

© The Author(s), 2022. Published by Cambridge University Press on behalf of the Weed Science Society of America. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.



(Smisek 1995). This substantial release of pollen could also contribute to the spread of resistant traits (Ye et al. 2016).

If no control strategies are implemented, GR horseweed interference can decrease cotton (*Gossypium hirsutum* L.), corn (*Zea mays* L.), and soybean yield up to 46%, 69%, and 93%, respectively (Byker et al. 2013b; Ford et al. 2014; Steckel and Gwathmey 2009). In conventional tillage systems, small plants can be controlled using aggressive fall and/or spring tillage (Kapusta 1979). In no-tillage crop production systems, the use of herbicides is imperative to manage GR horseweed (Bruce and Kells 1990). Generally, only preplant (PP) and preemergence (PRE) herbicides are effective for providing control of GR horseweed in soybean, because postemergence herbicides provide limited control (Byker et al. 2013c). Using PP or PRE herbicide mixes that consist of multiple effective modes of action can broaden the range of weeds controlled and slow the onset of herbicide resistance (Green and Owen 2011; Loux et al. 2006).

Variable GR horseweed control has been reported with glyphosate plus dicamba, 2,4-D ester, halauxifen-methyl, or pyraflufen-ethyl/2,4-D applied PP in soybean. Byker et al. (2013b) observed that glyphosate (900 g ae ha<sup>-1</sup>) plus dicamba (300 g ae ha<sup>-1</sup>) and glyphosate (900 g ae ha<sup>-1</sup>) plus 2,4-D ester (560 g ae ha<sup>-1</sup>) applied PP to soybean controlled GR horseweed 68% to 100% and 73% to 95%, respectively, at 8 WAA. Soltani et al. (2020a) and Zimmer et al. (2018) reported 71% and 87% GR horseweed control, respectively, with glyphosate (900 g ae ha<sup>-1</sup>) plus halauxifen-methyl (5 g ai ha<sup>-1</sup>) applied PP to soybean. Soltani et al. (2020a) reported that glyphosate (900 g ae ha<sup>-1</sup>) plus pyraflufen-ethyl/2,4-D (532 g ai ha<sup>-1</sup>) applied PP controlled GR horseweed 60% at 8 WAA. The aforementioned studies confirm variable GR horseweed control when glyphosate plus dicamba, 2,4-D ester, halauxifen-methyl, or pyraflufen-ethyl/2,4-D is applied PP to soybean.

The addition of an effective third herbicide mix partner such as saflufenacil or metribuzin may improve the level and consistency of GR horseweed control (Mellendorf et al. 2013). In addition, three-way mixes that include different effective modes of action can be useful to delay the evolution of herbicide resistance (Monks et al. 1993; Scott et al. 1998). Glyphosate (900 g ae ha<sup>-1</sup>) plus saflufenacil (25 g ai ha<sup>-1</sup>) plus dicamba (600 g ae ha<sup>-1</sup>) or 2,4-D ester (500 g ae ha<sup>-1</sup>) controlled GR horseweed 98% and 95%, respectively, at 8 WAA (Budd et al. 2016). Similarly, Zimmer et al. (2018) observed that glyphosate (1,120 g ae ha<sup>-1</sup>) plus 2,4-D (560 g ae ha<sup>-1</sup>) plus either metribuzin (210 g ai ha<sup>-1</sup>) or saflufenacil (37 g ai ha<sup>-1</sup>) controlled GR horseweed 82% and 97%, respectively, at 5 WAA. Glyphosate (900 g ae ha<sup>-1</sup>) plus metribuzin (400 g ai ha<sup>-1</sup>) plus pyraflufen-ethyl/2,4-D (532 g ai ha<sup>-1</sup>), applied PP, controlled GR horseweed 96% at 8 WAA (Soltani et al. 2020a). GR horseweed control with three-way mixes including saflufenacil or metribuzin appear to improve GR horseweed control, but further investigation is needed to see which provide better and more consistent GR horseweed control.

Variable GR horseweed control has been reported with glyphosate plus dicamba, 2,4-D ester, halauxifen-methyl, or pyraflufen-ethyl/2,4-D in soybean. With the aforementioned two-way mixes, it is not known whether the addition of saflufenacil or metribuzin would be the better third mix partner. The objective of this research was to ascertain if saflufenacil or metribuzin is a better mix partner with glyphosate plus dicamba, 2,4-D ester, halauxifen-methyl or pyraflufen-ethyl/2,4-D, applied PP, for GR horseweed control in soybean. Our hypothesis is that the addition of saflufenacil or metribuzin to glyphosate plus dicamba, 2,4-D ester, halauxifen-

methyl, or pyraflufen-ethyl/2,4-D, applied PP, will provide better and more consistent GR horseweed control in soybean.

## Materials and Methods

### Experimental Methods

Field research trials were completed over a 2-yr period at four different site locations in southwestern Ontario, Canada. Two trials were conducted in 2020, and two were conducted in 2021. The site, year, the nearest town to the site location, location coordinates, soil information, weather at the treatment application, treatment spray date, and soybean seeding and emergence are presented in Table 1. The horseweed resistance profile for each location was confirmed in greenhouse screenings. Horseweed seed was collected randomly from multiple plants at each location. Transplanting flats (25 cm × 25 cm × 5 cm) were filled with potting mix (Berger Growing Media with sphagnum peat moss, perlite, wetting agent, dolomitic and calcitic limestone) and were watered until the soil was completely saturated. Horseweed seeds (approximately 300) were sprinkled onto the soil surface. Approximately 1 mm of the potting mix was used to cover the seed on the soil surface. The trays remained in the greenhouse (16-h photoperiod with 26 C day and 17 C night temperatures) and were watered daily with approximately 20 ml of water. Once the seedlings had at least four leaves, 100 horseweed plants from each population were transplanted into individual circular pots measuring 10 cm diam. Once the horseweed reached 10 cm diam, 40 horseweed plants from each population were sprayed with glyphosate (900 g ae ha<sup>-1</sup>), and another 40 were sprayed with cloransulam-methyl (17.5 g ai ha<sup>-1</sup>). The horseweed was sprayed in a spray chamber equipped with flat-fan nozzles calibrated to deliver 205 L ha<sup>-1</sup> at 2.6 km h<sup>-1</sup> and 280 kPa. Two untreated checks for every 10 horseweed plants were used as comparisons to conduct the visible-control ratings. Visible-control ratings were completed 1, 3, and 5 WAA with a 0% to 100% scale; 0% represented no control, 100% represented complete necrosis (Canadian Weed Science Society 2018). The values in Table 2 represent the percent of horseweed resistant to glyphosate and cloransulam-methyl at each location at 5 WAA. Seed was not collected at the Bothwell site, and therefore resistance screening was not conducted for this location.

The study was a 3-by-5 factorial structured as a randomized complete block design with four replications. Factor One was the Group 4 herbicides control, dicamba, 2,4-D ester, halauxifen-methyl, and pyraflufen-ethyl/2,4-D, and Factor Two was control, saflufenacil, and metribuzin. Glyphosate (900 g ae ha<sup>-1</sup>) was included in each herbicide treatment to remove the confounding effects of glyphosate-susceptible horseweed and other weed species. There were a total of 14 treatments plus one weedy control. The plot width was 2.25 m, encompassing three soybean rows with 0.75-m spacings. The plot length was 8 m, and a 2-m alleyway separated each replicate. A CO<sub>2</sub>-pressurized backpack sprayer calibrated to deliver 200 L ha<sup>-1</sup> at 240 kPa was used to make the PP herbicide applications when horseweed approached 10 cm in height or diameter. The handheld boom measured 1.5 m in length equipped with 4 ULD 120-02 nozzles spaced 50 cm apart. The spray width was 2 m. Roundup Ready 2 Xtend<sup>®</sup> soybean cultivar (DKB12-16) was seeded to a 3.8-cm depth at about 416,000 seeds ha<sup>-1</sup> following the PP herbicide applications. Glyphosate (450 g ae ha<sup>-1</sup>) was applied postemergence to the entire experimental area to control glyphosate-susceptible horseweed

**Table 1.** The site, year, the nearest town to the site location, location coordinates, soil characteristics, weather at the treatment application, treatment spray date, and soybean seeding and emergence dates for four field trials conducted in southwestern Ontario, Canada in 2020 and 2021.

| Site | Year | Nearest town | Location coordinates | Soil characteristics |      |      |      |                 | Weather at treatment application |                 |                   | Agronomic information      |                      |                      |                        |
|------|------|--------------|----------------------|----------------------|------|------|------|-----------------|----------------------------------|-----------------|-------------------|----------------------------|----------------------|----------------------|------------------------|
|      |      |              |                      | Texture              | Sand | Silt | Clay | OM <sup>a</sup> | pH                               | Air temperature | Relative humidity | Wind speed                 | Treatment spray date | Soybean seeding date | Soybean emergence date |
| S1   | 2020 | Ridgetown    | 42.46° N, 81.85° W   | Sandy loam           | 75   | 17   | 7    | 1.9             | 7.1                              | 25              | 60                | 7.5–8.8 km h <sup>-1</sup> | May 28               | June 5               | June 11                |
| S2   | 2020 | Zone Centre  | 42.64° N, 81.94° W   | Loamy sand           | 85   | 9    | 5    | 2.9             | 6.5                              | 25              | 26                | 3.6–4.8                    | June 17              | June 24              | July 3                 |
| S3   | 2021 | Kintyre      | 42.56° N, 81.77° W   | Sandy loam           | 53   | 29   | 18   | 4.4             | 6.9                              | 21              | 50                | 0.5–1.2                    | May 19               | May 20               | May 25                 |
| S4   | 2021 | Bothwell     | 42.62° N, 81.91° W   | Loamy Sand           | 85   | 11   | 4    | 3.3             | 6.8                              | 20              | 62                | 1.5–5.6                    | May 27               | June 12              | June 18                |

<sup>a</sup>Abbreviations: OM, organic matter.**Table 2.** The site, year, the nearest town to the site location, glyphosate-resistant (GR) horseweed size and density at the time of the treatment application, and resistance profile for four field trials in southwestern Ontario, Canada in 2020 and 2021.

| Site | Year | Nearest town | GR horseweed |                        | Resistance profile |                    |
|------|------|--------------|--------------|------------------------|--------------------|--------------------|
|      |      |              | Size         | Density                | Glyphosate         | Cloransulam-methyl |
|      |      |              | cm           | plants m <sup>-2</sup> | %                  |                    |
| S1   | 2020 | Ridgetown    | 6            | 384                    | 100                | 99                 |
| S2   | 2020 | Zone Centre  | 10           | 57                     | 79                 | 100                |
| S3   | 2021 | Kintyre      | 8            | 147                    | 98                 | 85                 |
| S4   | 2021 | Bothwell     | 9            | 46                     | –                  | –                  |

**Table 3.** Herbicides and surfactants used in four field trials examining the control of glyphosate-resistant horseweed using herbicide mixes in southwestern Ontario, Canada in 2020 and 2021.

| Common name                   | Trade name      | Manufacturer                             |
|-------------------------------|-----------------|--|
| Herbicides                    |                 |  |
| Metribuzin                    | Sencor 480SC    | Bayer CropScience Inc., Calgary, AB      |
| Saflufenacil <sup>a</sup>     | Eragon LQ       | BASF Canada Inc., Mississauga, ON        |
| Pyraflufen-ethyl/2,4-D        | Blackhawk       | Nufarm Canada, Calgary, AB               |
| Halaxifen-methyl <sup>b</sup> | Elevore         | Dow AgroSciences Canada Inc. Calgary, AB |
| 2,4-D ester                   | Ester 700       | Nufarm Canada                            |
| Dicamba                       | Xtendimax       | Bayer CropScience Inc.                   |
| Surfactants                   |                 |  |
| Nonionic surfactant           | Merge           | BASF Canada Inc.                         |
| Methylated seed oil           | MSO concentrate | Dow AgroSciences Canada Inc.             |

<sup>a</sup>The surfactant Merge (1 L ha<sup>-1</sup>) was included in treatments with saflufenacil.<sup>b</sup>The surfactant MSO (1% v/v) was included in all treatments with halaxifen-methyl.

plus other weed species. The postemergence application was applied when soybean was at the V1 to V4 growth stages at all locations. Horseweed size and density at the time of the PP treatment application are included in Table 2. The herbicides used in this study are included in Tables 3 and 4.

A visual rating (%) was used to assess horseweed control 2, 4, and 8 WAA using a rating scale of 0 to 100; 0 indicated no control, and 100 indicated plant death (Canadian Weed Science Society 2018). Horseweed density was collected 8 WAA. Quadrats measuring 0.25 m<sup>2</sup> were placed randomly in the plot—one quadrat in the front and one in the back. Plants were counted in each quadrat to determine density. Plant biomass was assessed 8 WAA by cutting all plants within both quadrats close to the soil surface, putting all samples in bags, and drying the samples in a kiln at 60 C for approximately 2 wk or until the samples reached stable moisture and the biomass was recorded. Soybean injury (%) was visually evaluated 2 and 4 wk after soybean emergence using a rating scale of 0 to 100; 0 indicated no soybean injury and 100 indicated soybean death (Canadian Weed Science Society 2018). A small-plot harvester combined two soybean rows per plot and recorded soybean moisture and weight. Before analysis, soybean grain yield was corrected to 13.5% moisture content.

### Statistical Analysis

PROC GLIMMIX in SAS version 9.4 (Statistical Analysis Systems 2020) was used for the analyses. The study was a mixed model with



**Table 4.** Herbicide treatments and rates used in the present study for glyphosate-resistant horseweed control in southwestern Ontario, Canada in 2020 and 2021.

| Treatment <sup>a</sup>                | Rate                     |
|---------------------------------------|--------------------------|
|                                       | g ai/ae ha <sup>-1</sup> |
| Saflufenacil <sup>b</sup>             | 25                       |
| Metribuzin                            | 400                      |
| Dicamba                               | 600                      |
| Dicamba + saflufenacil                | 600 + 25                 |
| Dicamba + metribuzin                  | 600 + 400                |
| 2,4-D ester                           | 528                      |
| 2,4-D ester + saflufenacil            | 528 + 25                 |
| 2,4-D ester + metribuzin              | 528 + 400                |
| Halauxifen-methyl <sup>c</sup>        | 5                        |
| Halauxifen-methyl + saflufenacil      | 5 + 25                   |
| Halauxifen-methyl + metribuzin        | 5 + 400                  |
| Pyraflufen-ethyl/2,4-D                | 532                      |
| Pyraflufen-ethyl/2,4-D + saflufenacil | 532 + 25                 |
| Pyraflufen-ethyl/2,4-D + metribuzin   | 532 + 400                |

<sup>a</sup>Glyphosate (900 g ae ha<sup>-1</sup>) was included in all herbicide treatments.

<sup>b</sup>The surfactant Merge (1 L ha<sup>-1</sup>) was included in all treatments with saflufenacil.

<sup>c</sup>The surfactant MSO (1 % v/v) was included in all treatments with halauxifen-methyl.

a fixed effect (saflufenacil or metribuzin, Group 4 herbicide, saflufenacil or metribuzin by Group 4 herbicide) and random effects (location, block within the location, location-by-saflufenacil or metribuzin, location-by-Group 4 herbicide). Normality was confirmed after conducting the Shapiro-Wilk test and reviewing studentized residual plots. An arcsine square-root back-transformation was used for control 2, 4, and 8 WAA. A log-normal distribution was used for density and biomass. Soybean yield was analyzed using a normal distribution. The least-square means were analyzed in the analysis format then converted back to the data format using the ilink option except when log-normal was specified, which used the omega procedure to back-transform means. Tukey-Kramer's multiple-range test ( $\alpha = 0.05$ ) was used to compare the least-square means. Letter codes were assigned in Tables 5 and 6 to indicate significant data.

A common method to investigate herbicide interactions is with Colby's equation (Colby 1967). Because glyphosate (900 g ae ha<sup>-1</sup>) is not effective on GR horseweed and was only included in the tank to control glyphosate-susceptible horseweed and other weed species, only two-way interactions between the Group 4 herbicides and saflufenacil or metribuzin were evaluated. Previous literature used the two-way Colby's equation to determine the expected control of a three-way mix when one of the herbicides was not effective on the weed of interest (Meyer and Norsworthy 2019). The expected control mean was calculated using the observed control means for *A* (Group 4 herbicide) and *B* (Saflufenacil or metribuzin) in the Colby's equation.

$$\text{Expected} = (A + B) - \left( \frac{A \times B}{100} \right) \quad (1)$$

The expected density and biomass data were calculated using an adjusted Colby's equation. The nontreated control mean was represented as *W* in the equation.

$$\text{Expected} = \left( \frac{A \times B}{W} \right) \quad (2)$$

A t-test was used to compare the observed and expected values. Significance was noted when  $P < 0.05$ . If the control values differ

from one another (i.e., if the observed value is greater than or less than the expected value), then the interaction is considered synergistic or antagonistic, respectively. If the observed value is the same as expected, the interaction is considered additive (Colby 1967). In contrast, if the density or biomass values differ from one another (i.e., if the observed value is greater than or less than the expected value), then the interaction is considered antagonistic or synergistic, respectively (Colby 1967).

The coefficient of variation (CV) was calculated to determine the consistency of GR horseweed control for each least-square mean estimate. When comparing treatments, a lower CV would indicate greater consistency in control, whereas a higher CV would indicate less consistency in control (Shechtman 2013).

## Results and Discussion

### Soybean Injury

There was minimal soybean injury ( $\leq 5\%$ ) at all sites across both years (data not shown).

### Glyphosate-Resistant Horseweed Control, Density, and Biomass

There was a significant interaction between Factor One (control, dicamba, 2,4-D ester, halauxifen-methyl, and pyraflufen-ethyl/2,4-D) and Factor Two (control, saflufenacil, and metribuzin) for GR horseweed control 2, 4, and 8 WAA, density, and biomass, so the simple effects are presented (Table 6).

Glyphosate plus dicamba controlled GR horseweed 48%, 87%, and 96% at 2, 4, and 8 WAA, respectively, and decreased density and biomass 93% and 98%, respectively (Table 6). Byker et al. (2013b) observed that glyphosate (900 g ae ha<sup>-1</sup>) plus dicamba (300 g ae ha<sup>-1</sup>) provided a minimum of 68% GR horseweed control at 8 WAA. The decreases in GR horseweed density and biomass reported in this study are similar to those reported by Eubank et al. (2008) and Byker et al. (2013b), respectively. When saflufenacil was added to glyphosate plus dicamba, GR horseweed control improved by 43% at 2 WAA; control did not improve at 4 or 8 WAA, and there was no decrease in density or biomass. Similarly, Hedges et al. (2018) observed a 43% increase in GR horseweed control at 2 WAA and no reduction in density or biomass when saflufenacil (25 g ai ha<sup>-1</sup>) was added to glyphosate/dicamba (1,800 g ae ha<sup>-1</sup>). Based on Colby's equation, saflufenacil plus glyphosate plus dicamba was additive for control 2 and 4 WAA and antagonistic for control 8 WAA (Equation 1). The observed density and biomass were greater than the expected, indicating an antagonistic interaction (Equation 2). GR horseweed control was not improved when metribuzin was added to glyphosate plus dicamba 2, 4, or 8 WAA, and there was no reduction in density or biomass. The addition of metribuzin to glyphosate plus dicamba was additive for control 2, 4, and 8 WAA and for biomass reduction. The observed densities were greater than expected indicating an antagonistic interaction.

Glyphosate plus 2,4-D ester controlled GR horseweed 50%, 79%, and 77% at 2, 4, and 8 WAA, respectively; there was no decrease in density or biomass (Table 6). Similar control was observed by Byker et al. (2013a) at two site locations at 4 WAA, whereas Soltani et al. (2020b) observed only 53% GR horseweed control with glyphosate (900 g ae ha<sup>-1</sup>) plus 2,4-D ester (500 g ae ha<sup>-1</sup>) at 8 WAA. When saflufenacil was added to glyphosate plus 2,4-D ester, GR horseweed control increased by 43% at 2 WAA, and biomass was reduced 85%; however, there was no

**Table 5.** Main effects and interaction for glyphosate-resistant (GR) horseweed control 2, 4, and 8 wk after application (WAA), density, biomass, and soybean yield with metribuzin or saflufenacil-based mixes with Group 4 herbicides from four factorial trials conducted in southwestern Ontario, Canada in 2020 and 2021.<sup>a,b</sup>

| Main effects  | Rate                     | GR horseweed control |       |       | Density <sup>c</sup>       | Biomass <sup>c</sup> | Soybean yield <sup>d</sup> |
|---|--------------------------|----------------------|-------|-------|----------------------------|----------------------|----------------------------|
|   |                          | 2 WAA                | 4 WAA | 8 WAA |                            |                      |                            |
|   | g ai/ae ha <sup>-1</sup> | —                    | —     | —     | No. plants m <sup>-2</sup> | g m <sup>-2</sup>    | kg ha <sup>-1</sup>        |
| <b>Group 4 herbicide</b>                              |                          | **                   | **    | **    | *                          | **                   | *                          |
| None  | —                        | 41                   | 48    | 46    | 46                         | 172                  | 1,840 b                    |
| Pyraflufen-ethyl/2,4-D                                | 532                      | 80                   | 85    | 85    | 27                         | 68                   | 2,340 ab                   |
| Halauxifen-methyl <sup>e</sup>                        | 5                        | 77                   | 89    | 91    | 21                         | 19                   | 2,380 a                    |
| 2,4-D ester   | 528                      | 73                   | 87    | 88    | 22                         | 33                   | 2,370 a                    |
| Dicamba   | 600                      | 73                   | 93    | 97    | 9                          | 4                    | 2,480 a                    |
| SE <sup>b</sup>                                       |                          | 2                    | 2     | 2     | 5                          | 5                    | 0                          |
| <b>Metribuzin or saflufenacil</b>                     |                          | **                   | **    | **    | **                         | **                   | NS                         |
| None  | —                        | 34                   | 55    | 59    | 67                         | 87                   | 2,080                      |
| Metribuzin  | 400                      | 71                   | 84    | 86    | 13                         | 46                   | 2,370                      |
| Saflufenacil <sup>f</sup>                             | 25                       | 94                   | 97    | 98    | 5                          | 7                    | 2,390                      |
| SE  |                          | 2                    | 2     | 2     | 5                          | 5                    | 0                          |
| <b>Group 4 herbicide × metribuzin or saflufenacil</b> |                          | **                   | **    | **    | **                         | **                   | NS                         |

<sup>a</sup>\*\*\* Statistically significant when  $P < 0.01$ ; \* significant when  $P < 0.05$ ; NS, non-significant.

<sup>b</sup>Abbreviation: SE, standard error of the mean.

<sup>c</sup>Density and biomass were collected 8 WAA.

<sup>d</sup>Means accompanied by a different letter in a column (a–b) significantly differ based on Tukey-Kramer's LSD ( $\alpha = 0.05$ ).

<sup>e</sup>Treatments with halauxifen-methyl included the surfactant MSO (1% v/v).

<sup>f</sup>Treatments with saflufenacil included the surfactant Merge (1 L ha<sup>-1</sup>).

improvement in control at 4 and 8 WAA and no decrease in density. Mahoney et al. (2016) reported similar results with a 98% decrease in GR horseweed biomass when saflufenacil (25 g ai ha<sup>-1</sup>) was added to glyphosate (900 g ae ha<sup>-1</sup>) plus 2,4-D ester (500 g ae ha<sup>-1</sup>). Based on Colby's equation, the addition of saflufenacil to glyphosate plus 2,4-D ester was additive for control at 2 WAA (Equation 1) and for the reduction of density and biomass (Equation 2). The expected control values were greater than the observed control values at 4 and 8 WAA, indicating an antagonistic interaction. When metribuzin was added to glyphosate plus 2,4-D ester, GR horseweed control did not improve and there was no reduction in density and biomass. In contrast, Soltani et al. (2020b) observed a 32% and 36% improvement in GR horseweed control at 4 and 8 WAA, respectively, and a 90% and 88% decrease in GR horseweed density and biomass, respectively, when metribuzin (400 g ai ha<sup>-1</sup>) was added to glyphosate (900 g ae ha<sup>-1</sup>) plus 2,4-D ester (500 g ae ha<sup>-1</sup>). Based on Colby's equation, metribuzin plus glyphosate plus 2,4-D ester was additive for control at 2 and 8 WAA and for the reduction in density and biomass. The observed control values were less than the expected control values at 4 WAA, indicating an antagonistic interaction.

Glyphosate plus halauxifen-methyl controlled GR horseweed 40%, 63%, and 71% at 2, 4, and 8 WAA, respectively, and did not reduce density and biomass (Table 6). Similar control was observed by Soltani et al. (2020a) at 8 WAA, but contrasts with Zimmer et al. (2018), who observed 87% GR horseweed control using glyphosate (560 g ae ha<sup>-1</sup>) plus halauxifen-methyl (5 g ai ha<sup>-1</sup>) at 5 WAA. In the present study, when saflufenacil was added to glyphosate plus halauxifen-methyl, GR horseweed control improved by 57%, 36%, and 27% at 2, 4, and 8 WAA, respectively, and decreased GR horseweed density and biomass by 95% and 92%, respectively (Table 6). Similarly, Quinn et al. (2021) observed 91% GR horseweed control using glyphosate (900 g ae ha<sup>-1</sup>) plus halauxifen-methyl (5 g ai ha<sup>-1</sup>) plus saflufenacil (25 g ai ha<sup>-1</sup>) when applied PP in soybean at 8 WAA and a 97% and 98% decrease in density and biomass, respectively. Based on Colby's equation, the addition of saflufenacil to glyphosate plus

halauxifen-methyl was additive for control 4 and 8 WAA and synergistic at 2 WAA; the interaction for density and biomass reduction was additive (Equations 1 and 2). When metribuzin was added to glyphosate plus halauxifen-methyl, GR horseweed control improved by 43%, 31%, and 25% at 2, 4, and 8 WAA, respectively, and decreased density and biomass 94% and 86%, respectively. Similar results were reported by Quinn et al. (2021), who observed 93% GR horseweed control in soybean at 8 WAA and a 98% and 99% reduction in density and biomass, respectively, with glyphosate (900 g ae ha<sup>-1</sup>) plus halauxifen-methyl (5 g ai ha<sup>-1</sup>) plus metribuzin (400 g ai ha<sup>-1</sup>). Based on Colby's equation, the improvement in GR horseweed control 2, 4, and 8 WAA and the decrease in density and biomass was synergistic when metribuzin was added to glyphosate plus halauxifen-methyl (Equations 1 and 2).

Glyphosate plus pyraflufen-ethyl/2,4-D controlled GR horseweed 48%, 55%, and 52% at 2, 4, and 8 WAA, respectively; there was no reduction in density and biomass (Table 6). When saflufenacil was added to glyphosate plus pyraflufen-ethyl/2,4-D, GR horseweed control improved by 49%, 44%, and 47% at 2, 4, and 8 WAA, respectively, and decreased density and biomass by 97% and 95%, respectively. Based on Colby's equation, the improvement in GR horseweed control at 2 and 8 WAA from the addition of saflufenacil to glyphosate plus pyraflufen-ethyl/2,4-D was synergistic; the interactions for control at 4 WAA and density and biomass were additive. When metribuzin was added to glyphosate plus pyraflufen-ethyl/2,4-D, GR horseweed control improved by 36%, 34%, and 37% at 2, 4, and 8 WAA, respectively, and decreased density and biomass by 89% and 93%, respectively. Similarly, Soltani et al. (2020a) observed 95% GR horseweed control and a 98% and 97% density and biomass reduction, respectively, with glyphosate (900 g ae ha<sup>-1</sup>) plus pyraflufen-ethyl/2,4-D (532 g ai ha<sup>-1</sup>) plus metribuzin (400 g ai ha<sup>-1</sup>) applied PP in soybean at 8 WAA. Based on the Colby's equation, the improvement in GR horseweed control 2, 4, and 8 WAA and the decrease in biomass was synergistic when metribuzin was added to glyphosate plus pyraflufen-ethyl/2,4-D; the interaction for density was additive.

**Table 6.** The level of glyphosate-resistant horseweed control 2, 4, and 8 wk after application (WAA), density, and biomass from four factorial trials in southwestern Ontario, Canada in 2020 and 2021.<sup>a,b,c,d</sup>

|                                | Metribuzin or saflufenacil |              |                           |    |
|--------------------------------|----------------------------|--------------|---------------------------|----|
| Group 4 herbicide              | None                       | Metribuzin   | Saflufenacil <sup>e</sup> | SE |
| <b>Weed control 2 WAA</b>      | %                          |              |                           |    |
| None                           | 0b X                       | 43b Y        | 91a Z                     | 6  |
| Pyraflufen-ethyl/2,4-D         | 48a Y                      | 84a Z (69)** | 97a Z (88)*               | 4  |
| Halauxifen-methyl <sup>c</sup> | 40a Y                      | 83a Z (65)*  | 97a Z (87)*               | 4  |
| 2,4-D ester                    | 50a Y                      | 69a Y (70)   | 93a Z (89)                | 4  |
| Dicamba                        | 48a Y                      | 70a Y (68)   | 94a Z (88)                | 4  |
| SE                             | 3                          | 3            | 2                         |    |
| <b>Weed control 4 WAA</b>      |                            |              |                           |    |
| None                           | 0c X                       | 51b Y        | 96a Z                     | 6  |
| Pyraflufen-ethyl/2,4-D         | 55b Y                      | 89a Z (74)*  | 99a Z (96)                | 4  |
| Halauxifen-methyl <sup>f</sup> | 63b Y                      | 94a Z (79)*  | 99a Z (96)                | 3  |
| 2,4-D ester                    | 79ab Z                     | 85a Z (89)*  | 96a Z (98)**              | 2  |
| Dicamba                        | 87a Z                      | 91a Z (90)   | 98a Z (98)                | 2  |
| SE                             | 4                          | 3            | 1                         |    |
| <b>Weed control 8 WAA</b>      |                            |              |                           |    |
| None                           | 0c X                       | 46b Y        | 95a Z                     | 6  |
| Pyraflufen-ethyl/2,4-D         | 52b Y                      | 89a Z (73)*  | 99a Z (94)*               | 4  |
| Halauxifen-methyl <sup>f</sup> | 71b Y                      | 96a Z (86)** | 98a Z (97)                | 2  |
| 2,4-D ester                    | 77b Z                      | 88ab Z (86)  | 96a Z (97)*               | 2  |
| Dicamba                        | 96a Z                      | 96ab Z (98)  | 98a Z (100)**             | 1  |
| SE                             | 4                          | 3            | 1                         |    |
| <b>Density</b>                 | No. plants m <sup>-2</sup> |              |                           |    |
| None                           | 130b Y                     | 28a ZY       | 8a Z                      | 17 |
| Pyraflufen-ethyl/2,4-D         | 96b Y                      | 11a ZY (43)  | 3a Z (7)                  | 12 |
| Halauxifen-methyl <sup>f</sup> | 77b Y                      | 5a ZY (23)*  | 4a Z (10)                 | 10 |
| 2,4-D ester                    | 63ab Z                     | 13a Z (16)   | 6a Z (6)                  | 10 |
| Dicamba                        | 9a Z                       | 15a Z (5)*   | 6a Z (1)*                 | 2  |
| SE                             | 13                         | 2            | 1                         |    |
| <b>Biomass</b>                 | g m <sup>-2</sup>          |              |                           |    |
| None                           | 188b Z                     | 206b Z       | 25a Y                     | 16 |
| Pyraflufen-ethyl/2,4-D         | 114b Z                     | 8a Y (97)*   | 6a Y (19)                 | 10 |
| Halauxifen-methyl <sup>f</sup> | 50b Z                      | 7a Y (30)*   | 4a Y (15)                 | 5  |
| 2,4-D ester                    | 53b Z                      | 38a Z (51)   | 8a Z (12)                 | 5  |
| Dicamba                        | 3a Z                       | 8a Z (2)     | 2a Z (1)*                 | 1  |
| SE                             | 9                          | 10           | 3                         |    |

<sup>a</sup>Abbreviations: SE, standard error of the mean.<sup>b</sup>Means accompanied with a different letter in a column (a–c) or row (X–Z) within each section significantly differ based on Tukey-Kramer's LSD ( $\alpha = 0.05$ ).<sup>c</sup>\*\* Significant at  $P < 0.01$ ; \* significant at  $P < 0.05$  based on a t-test comparing observed and expected values.<sup>d</sup>Values in parentheses represent the expected values from Colby's analysis.<sup>e</sup>Treatments with saflufenacil included the surfactant Merge (1 L ha<sup>-1</sup>).<sup>f</sup>Treatments with halaxifen-methyl included the surfactant MSO (1% v/v).

Glyphosate plus saflufenacil controlled GR horseweed 91%, 96%, and 95% at 2, 4, and 8 WAA, respectively, and decreased density and biomass 94% and 87%, respectively (Table 6). Eubank et al. (2013) reported similar findings at 3 WAA, but the results from this study are in contrast to Ikley (2012), who observed 57% GR horseweed control with glyphosate (874 g ae ha<sup>-1</sup>) plus saflufenacil (25 g ai ha<sup>-1</sup>) at 5 WAA. There was no improvement in GR horseweed control when dicamba, 2,4-D ester, halaxifen-methyl, or pyraflufen-ethyl/2,4-D were added to glyphosate plus saflufenacil 2, 4, or 8 WAA, and there was no decrease in density or biomass. Similarly, Budd et al. (2016) observed no improvement in GR horseweed control at 4 and 8 WAA and no reduction in density when dicamba (600 g ae ha<sup>-1</sup>) or 2,4-D ester (500 g ae ha<sup>-1</sup>) were added to glyphosate (900 g ae ha<sup>-1</sup>) plus saflufenacil (25 g ai ha<sup>-1</sup>). In the present study, all mixes that included saflufenacil provided the fastest GR horseweed control, with greater than 90% control at 2 and 4 WAA.

Glyphosate plus metribuzin controlled GR horseweed 43%, 51%, and 46% at 2, 4, and 8 WAA, respectively; density and

**Table 7.** The consistency of glyphosate-resistant horseweed control 2, 4, and 8 wk after application (WAA) from four factorial trials in southwestern Ontario, Canada in 2020 and 2021.

| Group 4 herbicide                   | Saflufenacil or metribuzin |            |                           |
|-------------------------------------|----------------------------|------------|---------------------------|
|                                     | None                       | Metribuzin | Saflufenacil <sup>a</sup> |
| <b>Consistency of control 2 WAA</b> |                            |            |                           |
| None                                | —                          | 87.1       | 49.6                      |
| Pyraflufen-ethyl/2,4-D              | 81.6                       | 54.1       | 44.9                      |
| Halaxifen-methyl <sup>b</sup>       | 90.6                       | 54.7       | 45.1                      |
| 2,4-D ester                         | 80.0                       | 63.4       | 47.9                      |
| Dicamba                             | 81.5                       | 62.8       | 47.4                      |
| <b>Consistency of control 4 WAA</b> |                            |            |                           |
| None                                | —                          | 67.9       | 38.9                      |
| Pyraflufen-ethyl/2,4-D              | 64.9                       | 42.9       | 36.3                      |
| Halaxifen-methyl <sup>b</sup>       | 57.2                       | 39.2       | 36.5                      |
| 2,4-D ester                         | 45.4                       | 43.9       | 37.9                      |
| Dicamba                             | 41.8                       | 40.4       | 36.6                      |
| <b>Consistency of control 8 WAA</b> |                            |            |                           |
| None                                | —                          | 63.3       | 37.7                      |
| Pyraflufen-ethyl/2,4-D              | 60.9                       | 41.1       | 34.6                      |
| Halaxifen-methyl <sup>b</sup>       | 50.5                       | 35.7       | 33.8                      |
| 2,4-D ester                         | 44.8                       | 39.9       | 35.6                      |
| Dicamba                             | 37.7                       | 36.2       | 34.5                      |

<sup>a</sup>Treatments with saflufenacil included the surfactant Merge (1 L ha<sup>-1</sup>).<sup>b</sup>Treatments with halaxifen-methyl included the surfactant MSO (1% v/v).

biomass were not reduced (Table 6). Similar results were reported by Eubank et al. (2008) but are in contrast to Byker et al. (2013a), who observed a minimum of 97% GR horseweed control using glyphosate (900 g ae ha<sup>-1</sup>) plus metribuzin (1,120 g ai ha<sup>-1</sup>) at 8 WAA. The improved GR horseweed control in the Byker et al. (2013a) study can be ascribed to the greater rate of metribuzin used in that study. When dicamba, 2,4-D ester, halaxifen-methyl, or pyraflufen-ethyl/2,4-D were added to glyphosate plus metribuzin, GR horseweed control improved by 27%, 26%, 40%, and 41% at 2 WAA, 40%, 34%, 43%, and 38% at 4 WAA, and 50%, 42%, 50%, and 43% at 8 WAA, respectively. When dicamba, 2,4-D ester, halaxifen-methyl, or pyraflufen-ethyl/2,4-D were added to glyphosate plus metribuzin, GR horseweed biomass was reduced by 96%, 82%, 97%, and 96%, respectively; however, there was no decrease in GR horseweed density.

### Glyphosate-Resistant Horseweed Consistency in Control

At 2, 4, and 8 WAA, adding saflufenacil or metribuzin to glyphosate plus dicamba, 2,4-D ester, halaxifen-methyl, or pyraflufen-ethyl/2,4-D improved the consistency of GR horseweed control (Table 7). The addition of saflufenacil consistently reduced the CV more than metribuzin, indicating improved consistency of GR horseweed control with saflufenacil-based mixes relative to metribuzin-based mixes. Similar to the current study, Budd et al. (2016) observed that when a third herbicide such as metribuzin (400 g ai ha<sup>-1</sup>) was added into the tank with glyphosate (900 g ae ha<sup>-1</sup>) plus saflufenacil (25 g ai ha<sup>-1</sup>) applied PP in soybean, there was improved consistency of GR horseweed control. Soltani et al. (2020b) also reported improved consistency of GR horseweed control when metribuzin (400 g ai ha<sup>-1</sup>) was added to glyphosate (900 g ae ha<sup>-1</sup>) plus saflufenacil (25 g ai ha<sup>-1</sup>) or 2,4-D ester (500 g ae ha<sup>-1</sup>).

### Soybean Yield

Soybean yield was reduced up to 26% due to GR horseweed interference. The main effects are presented (Table 5), as there



was no interaction between Factor One (control, dicamba, 2,4-D ester, halauxifen-methyl, and pyraflufen-ethyl/2,4-D) and Factor Two (control, saflufenacil, and metribuzin) on soybean yield. When averaged across Factor Two, reduced GR horseweed interference with the application of dicamba, 2,4-D ester, or halauxifen-methyl resulted in a soybean yield increase of 38% to 40% (Table 5). Similar soybean yields were found across all herbicide treatments. Soltani et al. (2020a) reported a similar soybean yield loss due to GR horseweed interference. In contrast, Eubank et al. (2008) reported a high soybean yield reduction of 97% from GR horseweed interference.

In conclusion, the addition of saflufenacil or metribuzin to glyphosate plus halauxifen-methyl or pyraflufen-ethyl/2,4-D applied PP in soybean improved the level and consistency of GR horseweed control at 8 WAA. GR horseweed control was not improved when saflufenacil or metribuzin was added to glyphosate plus dicamba or 2,4-D ester, though the consistency of control was improved. When saflufenacil was added to glyphosate plus halauxifen-methyl or pyraflufen-ethyl/2,4-D, there was reduced GR horseweed density and biomass. When metribuzin was added to glyphosate plus halauxifen-methyl or pyraflufen-ethyl/2,4-D, there was reduced GR horseweed biomass. However, GR horseweed control was improved with the addition of saflufenacil or metribuzin to glyphosate plus halauxifen-methyl or pyraflufen-ethyl/2,4-D; all treatments including saflufenacil resulted in the highest level and most consistent control.

**Acknowledgments.** The authors would like to acknowledge Chris Kramer, Dr. Michelle Edwards, and Christy Shropshire for their technical assistance to this project. Funding for this project was in part by the OMAFRA Alliance program, Grain Farmers of Ontario (GFO), and the agricultural products companies: BASF Canada Inc., Bayer CropScience Inc., Dow AgroSciences Canada Inc., and Nufarm Canada. No other conflicts of interest are declared.

## References

- Bhowmik PC, Bekech MM (1993) Horseweed (*Conyza canadensis*) seed production, emergence, and distribution in no-tillage and conventional tillage corn (*Zea mays*). *Agron Trends Agric Sci* 1:67–71
- Bruce JA, Kells JJ (1990) Horseweed (*Conyza canadensis*) control in no-tillage soybeans (*Glycine max*) with preplant and preemergence herbicides. *Weed Technol* 4:642–647
- Budd CM, Soltani N, Robinson DE, Hooker DC, Miller RT, Sikkema PH (2016) Control of glyphosate-resistant Canada fleabane with saflufenacil plus mix partners in soybean. *Can J Plant Sci* 96:989–994
- Budd CM, Soltani N, Robinson DE, Hooker DC, Miller RT, Sikkema PH (2017) Distribution of glyphosate and cloransulam-methyl resistant Canada fleabane [*Conyza canadensis* (L.) Cronq.] in Ontario. *Can J Plant Sci* 98:492–497
- Byker HP, Soltani N, Robinson DE, Tardif FJ, Lawton MB, Sikkema PH (2013a) Occurrence of glyphosate- and cloransulam-resistant Canada fleabane [*Conyza canadensis* (L.) Cronq.] in Ontario. *Can J Plant Sci* 93:851–855
- Byker HP, Soltani N, Robinson DE, Tardif FJ, Lawton MB, Sikkema PH (2013b) Control of glyphosate-resistant horseweed (*Conyza canadensis*) with dicamba applied preplant and postemergence in dicamba-resistant soybean. *Weed Technol* 27:492–496
- Byker HP, Soltani N, Robinson DE, Tardif FJ, Lawton MB, Sikkema PH (2013c) Control of glyphosate-resistant Canada fleabane [*Conyza canadensis* (L.) Cronq.] with preplant herbicide mixes in soybean [*Glycine max* (L.) Merr.]. *Can J Plant Sci* 93:659–667
- Byker HP, Soltani N, Robinson DE, Tardif FJ, Lawton MB, Sikkema PH (2013d) Glyphosate-resistant Canada fleabane [*Conyza canadensis* (L.) Cronq.]: dose response to glyphosate and control with postemergence herbicides in soybean in Ontario. *Can J Plant Sci* 93:1187–1193
- Canadian Weed Science Society (2018) Description of 0–100 rating scale for herbicide efficacy and crop phytotoxicity. Department of Plant Agriculture, University of Guelph, Ridgetown, ON, Canada. [https://weedsociety.ca/cwss\\_scm-rating-scale/](https://weedsociety.ca/cwss_scm-rating-scale/). Accessed: April 13, 2022
- Colby SR (1967) Calculating synergistic and antagonistic responses of herbicide combinations. *Weeds* 15:20–22
- Davis VM, Kruger GR, Stachler JM, Loux MM, Johnson WG (2009) Growth and seed production of horseweed (*Conyza canadensis*) populations resistant to glyphosate, ALS-inhibiting, and multiple (glyphosate + ALS-inhibiting) herbicides. *Weed Sci* 57:494–504
- Eubank TW, Nandula VK, Reddy KN, Poston DH, Shaw DR (2013) Saflufenacil efficacy on horseweed and its interaction with glyphosate. *Weed Biol Manag* 13:135–143
- Eubank TW, Poston DH, Nandula VK, Koger CH, Shaw DR, Reynolds DB (2008) Glyphosate-resistant horseweed (*Conyza canadensis*) control using glyphosate-, paraquat-, and glufosinate-based herbicide programs. *Weed Technol* 22:16–21
- Ford L, Soltani N, Robinson DE, Nurse RE, McFadden A, Sikkema PH (2014) Canada fleabane (*Conyza canadensis*) control with preplant applied residual herbicides followed by 2, 4-D choline/glyphosate DMA applied postemergence in corn. *Can J Plant Sci* 94:1231–1237
- Frankton C, Mulligan GA (1987) Weeds of Canada (revised). Publication 948. Ministry of Supply and Services Canada. Toronto: NC Press Ltd. Pages 156–157
- Green JM, Owen MDK (2011) Herbicide-resistant crops: Utilities and limitations for herbicide-resistant weed management. *J Agric Food Chem* 59:5819–5829
- Hedges BK, Soltani N, Robinson DE, Hooker DC, Sikkema PH (2018) Control of glyphosate-resistant Canada fleabane in Ontario with multiple effective modes-of-action in glyphosate/dicamba-resistant soybean. *Can J Plant Sci* 99:78–83
- Holm L, Doll J, Holm E, Pancho J, Herberger J (1997) World weeds: natural histories and distribution. Toronto: John Wiley & Sons, Inc. Pp 226–235
- Ikley JT (2012) The utility of saflufenacil on glyphosate-resistant horseweed and its effect on select soybean varieties. Master's Thesis, University of Maryland. 78 p. <http://hdl.handle.net/1903/12803>. Accessed: April 13, 2022
- Kapusta G (1979) Seedbed tillage and herbicide influence on soybean (*Glycine max*) weed control and yield. *Weed Sci* 27:520–526
- Loux M, Stachler J, Johnson B, Nice G, Davis V, Nordby D (2006) Biology and management of horseweed. The Glyphosate, Weeds, and Crops Series. Purdue extension. <https://www.extension.purdue.edu/extmedia/gwc/gwc-9-w.pdf>. Accessed: October 13, 2021
- Mahoney KJ, McNaughton KE, Sikkema PH (2016) Control of glyphosate-resistant horseweed in winter wheat with pyrasulfotole premixed with bromoxynil. *Weed Technol* 30:291–296
- Mellendorf TG, Young JM, Matthews JL, Young BG (2013) Influence of plant height and glyphosate on saflufenacil efficacy on glyphosate-resistant horseweed (*Conyza canadensis*). *Weed Technol* 27:463–467
- Meyer CJ, Norsworthy JK (2019) Influence of weed size on herbicide interactions for Enlist™ and Roundup Ready® Xtend® technologies. *Weed Technol* 33:569–577
- Monks CD, Wilcut JW, Richburg III JS (1993) Broadleaf weed control in soybean (*Glycine max*) with chlorimuron plus acifluorfen or thifensulfuron mixtures. *Weed Technol* 7:317–321
- Quinn J, Ashigh J, Soltani N, Hooker DC, Robinson DE, Sikkema PH (2021) Control of glyphosate-resistant horseweed and giant ragweed in soybean with halauxifen-methyl applied preplant. *Weed Technol* 35: 324–329
- Scott RC, Shaw DR, Ratliff RL, Newsom LJ (1998) Synergism of grass weed control with postemergence combinations of dimethenamid and fluzifop-p, imazethapyr, or sethoxydim. *Weed Technol* 12:268–274
- Shechtman O (2013) The coefficient of variation as an index of measurement reliability. Pages 39–49 in Doi S, Williams G, eds. *Methods of Clinical Epidemiology*. Springer Series on Epidemiology and Public Health. Berlin: Springer. [https://doi.org/10.1007/978-3-642-37131-8\\_4](https://doi.org/10.1007/978-3-642-37131-8_4). Accessed: April 18, 2022



- Shields EJ, Dauer JT, VanGessel MJ, Neumann G (2006) Horseweed (*Conyza canadensis*) seed collected in the planetary boundary layer. *Weed Sci* 54:1063–1067
- Smisek A (1995) The evolution of resistance to paraquat in populations of *Erigeron canadensis* L. MSc. Thesis. London, ON: University of Western Ontario. 102 pp
- Soltani N, Shropshire C, Sikkema PH (2020a) Control of glyphosate-resistant marehail in identity-preserved or glyphosate-resistant and glyphosate/dicamba-resistant soybean with preplant herbicides. *Am J Plant Sci* 11:851–860
- Soltani N, Shropshire C, Sikkema PH (2020b) Glyphosate-resistant Canada fleabane control with three-way herbicide mixes in soybean. *Am J Plant Sci* 11:1478–1486
- Statistical Analysis Systems (2020) The SAS System, Version 9.4. Cary, NC: Statistical Analysis Systems Institute
- Steckel LE, Gwathmey CO (2009) Glyphosate-resistant horseweed (*Conyza canadensis*) growth, seed production, and interference in cotton. *Weed Sci* 57:346–350
- VanGessel MJ (2001) Glyphosate-resistant horseweed from Delaware. *Weed Sci* 49:703–705
- Weaver SE (2001) The biology of Canadian weeds. 115. *Conyza canadensis*. *Can J Plant Sci* 81:867–875
- Ye R, Huang H, Alexander J, Liu W, Millwood RJ, Wang J, Stewart Jr CN (2016) Field studies on dynamic pollen production, deposition and dispersion of glyphosate-resistant horseweed (*Conyza canadensis*). *Weed Sci* 64:101–111
- Zimmer M, Young BG, Johnson WG (2018) Weed control with halauxifen-methyl applied alone and in mixtures with 2, 4-D, dicamba, and glyphosate. *Weed Technol*. 32:597–602