



Integrating Cover Crops for Weed Management in the Semiarid U.S. Great Plains: Opportunities and Challenges

Authors: Kumar, Vipin, Obour, Augustine, Jha, Prashant, Liu, Rui, Manuchehri, Misha R., et al.

Source: Weed Science, 68(4) : 311-323

Published By: Weed Science Society of America

URL: <https://doi.org/10.1017/wsc.2020.29>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Review

Cite this article: Kumar V, Obour A, Jha P, Liu R, Manuchehri MR, Dille JA, Holman J, Stahlman PW (2020) Integrating cover crops for weed management in the semiarid U.S. Great Plains: opportunities and challenges. *Weed Sci.* 68: 311–323. doi: [10.1017/wsc.2020.29](https://doi.org/10.1017/wsc.2020.29)

Received: 18 November 2019
Revised: 22 March 2020
Accepted: 9 April 2020
First published online: 16 April 2020

Associate Editor:

Vijay Nandula, USDA-ARS


Keywords:

Dryland cropping systems; herbicide-resistant weeds; integrated weed management

Author for correspondence:

Vipin Kumar, Kansas State University,
Agricultural Research Center, 1232 240th
Avenue, Hays, KS 67601.
(Email: vkumar@ksu.edu)

Integrating cover crops for weed management in the semiarid U.S. Great Plains: opportunities and challenges

Vipin Kumar¹ , Augustine Obour², Prashant Jha³, Rui Liu⁴, Misha R. Manuchehri⁵, J. Anita Dille⁶, John Holman⁷ and Phillip W. Stahlman⁸

¹Assistant Professor, Kansas State University, Agricultural Research Center, Hays, KS, USA; ²Associate Professor, Kansas State University, Agricultural Research Center, Hays, KS, USA; ³Associate Professor, Department of Agronomy, Iowa State University, Ames, IA, USA; ⁴Assistant Scientist, Kansas State University, Agricultural Research Center, Hays, KS, USA; ⁵Assistant Professor, Department of Plant and Soil Sciences, Oklahoma State University, Stillwater, OK, USA; ⁶Professor, Department of Agronomy, Kansas State University, Manhattan, KS, USA; ⁷Professor, Kansas State University, Southwest Research and Extension Center, Garden City, KS, USA and ⁸Emeritus Professor, Kansas State University, Agricultural Research Center, Hays, KS, USA

Abstract

The widespread evolution of herbicide resistance in weed populations has become an increasing concern for no-tillage (NT) growers in semiarid regions of the U.S. Great Plains. Lack of cost-effective and alternative new herbicide sites of action further exacerbates the problem of herbicide-resistant (HR) weeds and threatens the long-term sustainability of prevailing cropping systems in the region. A recent decline in commodity prices and increasing herbicide costs to manage HR weeds has spurred research efforts to build a strong rationale for developing ecologically based integrated weed management (IWM) strategies in the U.S. Great Plains. Integration of cover crops (CCs) in NT dryland production systems potentially offers several ecosystem services, including weed control, soil health improvement, decline in selective pest pressure, and overall reduction in pest management inputs. This review article aims to document the role of CCs for IWM, with emphasis on exploring emerging weed issues; ecological, economic, and agronomic benefits of growing CCs; and constraints preventing adoption of CCs in NT cropping systems in the semiarid Great Plains. We attempt to focus on changes in weed management practices, their long-term impacts on weed seedbanks, weed shifts, and herbicide-resistance evolution in the most common weed species in the region. We also highlight current knowledge gaps and propose new research priorities based on an improved understanding of CC management strategies that will ultimately aid in achieving sustainable weed management goals and preserving natural resources in water-limited environments.

Introduction

The U.S. Great Plains extends from the Canadian border in the north to Texas in the south and is bordered by Mississippi River in the east and the Rocky Mountains in the west (Unger and Baumhardt 2001). This region covers the central portions of the continental United States and accounts for more than 60% of total wheat (*Triticum aestivum* L.) production in the United States (Paulsen and Shroyer 2008). The majority of the cropland in this region encompasses a semiarid agroecosystem characterized by hot summer days with cold and dry winters (Lenssen et al. 2007). Soil moisture is the most limiting factor for crop production, generally decreasing from east to west with an average annual rainfall of <762 mm in the eastern portions of the Great Plains down to <381 mm in Montana, Wyoming, and far west Texas (Nielsen 2018; Peterson and Westfall 2004; Shafer et al. 2014). In addition to limited rainfall, soils are shallow, low in soil organic matter content, and prone to wind erosion. Temporal and spatial climatic variation with extended drought periods further challenge crop production in this region (Hansen et al. 2012). The Dust Bowl period during the 1930s led to the adoption of soil conservation practices, including fallow and minimum-tillage and no-tillage (NT) crop production across the region (Hansen et al. 2012; Smika and Wicks 1968). The transition to NT production was mainly designed to prevent wind and water erosion of the topsoil layer, improve soil organic matter, and conserve soil moisture.

Winter wheat–fallow (W–F) is a major NT dryland crop rotation in the Northern Great Plains (NGP) (Figure 1). Winter wheat–summer crop [corn (*Zea mays* L.), soybean [*Glycine max* (L.) Merr.], grain sorghum [*Sorghum bicolor* (L.) Moench ssp. *bicolor*], or sunflower (*Helianthus annuus* L.)]–fallow (W–S–F) is the most dominant crop rotation in the Southern and Central Great Plains (Figure 1) (Hansen et al. 2012; Lenssen et al. 2007; Nielsen and Vigil 2018; Peterson and Westfall 2004). The fallow phase in the rotation was

© Weed Science Society of America, 2020. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution, and reproduction in any medium, provided the original work is properly cited.





Figure 1. Map of the Great Plains showing three main regions: (1) Northern Great Plains (marked by purple line), (2) Central Great Plains (marked by red line), and (3) Southern Great Plains (marked by light blue line). Adapted from Center for Great Plains Studies, University of Nebraska–Lincoln.

originally included to store and conserve moisture in the soil profile from rainfall or snow events for successful establishment of a subsequent cash crop (Nielsen and Vigil 2010). The fallow period limits the occurrence of crop failure in drier years, stabilizes crop yields across years, and more importantly, prevents soil erosion and soil nutrient depletion, particularly when NT is adopted (Lenssen et al. 2007; Nielsen and Vigil 2018). Depending on location and crop rotation scheme, the fallow period extends from 10 mo (W–S–F) to almost 14 to 15 mo (W–F) after wheat harvest. The adoption of NT practices has increased soil water storage and cropping system intensification across the Great Plains region. More recently, many growers have started intensifying crop rotations to eliminate fallow, and some are eliminating wheat in their rotations for economic reasons. For instance, pulse crops such as field pea (*Pisum sativum* L.), lentil (*Lens culinaris* Medik.), and chick pea (*Cicer arietinum* L.) have replaced a portion of this fallow period in the northern parts (Miller et al. 2003). Similarly, several new crops have been evaluated in 3- or 4-yr rotation schemes by including cereal crops (Nielsen and Vigil 2018; Schlegel et al. 2018), legumes as forage and grain crops (Holman et al. 2018; Lyon et al. 2004), and recently oilseed crops (Obour et al. 2018) for potential replacement of fallow in the Central Great Plains (Figure 1). Despite these efforts to increase cropping intensity, crop rotations in the region still include a fallow phase, and most growers practice

fallow as a strategy to minimize risk of drought and crop failure. For example, two fallow periods occur in a typical 3-yr (W–S–F) crop rotation. The first fallow period extends from wheat harvest until summer crop planting the next year, and the second fallow period extends from summer crop harvest until winter wheat planting in the following year.

Weed control during fallow is typically accomplished through a combination of various tillage practices and herbicide applications in a minimum-tillage system or with herbicide only in NT. The successful adoption of NT fallow-based crop production in the semiarid Great Plains was possible due to effective weed control with the use of herbicides (Wicks and Smika 1973; Wicks et al. 1993). Weed control in the fallow phase of the crop rotation is crucial, as weed infestations can reduce available soil moisture and nutrients, negatively affecting subsequent cash crop yield (Wicks and Smika 1973). In general, most fallow fields could have three to five applications of herbicides (predominantly glyphosate-based tank mixtures) per year for weed control. The continuous and repetitive use of herbicides with same site(s) of action (SOA) has led to the evolution of herbicide resistance in weed populations across the region (Heap 2019). A lack of diversity in weed control programs has further led to the evolution of multiple herbicide resistance in several weed populations in the Great Plains in recent years. Dwindling herbicide options coupled with ever-increasing reports of multiple herbicide-resistant (MHR) weed populations necessitates the development of ecologically based weed management options for the region. With low grain prices and the high cost of controlling MHR weeds, some producers are returning to tillage as a strategic weed management tool. This poses a major threat to gains and investments made in soil conservation practices over the last several decades. Integration of cover crops (CCs) to replace portions of the fallow period in dryland NT crop production may provide effective and sustainable management of HR weed populations while enhancing the productivity and profitability of dominant cropping systems in the region. Growing a CC in place of fallow as part of the crop rotation can suppress weeds and reduce frequency of herbicide use. Several NT dryland producers in the region have also shown interest in growing CCs for controlling HR weeds and gaining soil health benefits. This recent trend has renewed interest among land managers, academic weed scientists, ecologists, agronomists, and soil scientists to improve their understanding of CC management strategies for weed suppression and soil health benefits in the region. In the literature, several research and review articles have documented the effects of CCs on weed suppression, soil health benefits, and subsequent cash crop yields across various regions in the United States. However, there is a paucity of published information on this topic in NT dryland production systems of the Great Plains. Previous review articles on CCs in the semiarid Great Plains have focused on soil health (Ghimire et al. 2018) or agronomy and water use of CCs (Unger and Vigil 1998), with little discussion on using CCs for weed suppression in the region. This review article aims to document weed species shifts that have occurred in response to herbicide use patterns in NT dryland production systems and to summarize the literature supporting utilization of CCs for control of these problematic species. This review will discuss opportunities and scenarios for use of CCs to suppress weeds, including agronomic considerations and challenges of adopting CC in this semiarid region. Finally, we will suggest future research directions and extension needs to be able to effectively use CCs as an integrated weed management (IWM) tool in the Great Plains.

Emerging Weed Management Issues in the Great Plains

Weed Control Practices and Weed Shifts

Weed control practices, including tillage, crop rotation, herbicide use patterns, and other production practices are known to influence weed community richness and population dynamics (Johnson et al. 2009). Two key changes in Great Plains agriculture have occurred over the past few decades: (1) adoption of soil conservation practices (changing from preplant conventional tillage to NT production) (Shaner 2000; Swanton et al. 1993; Wilson et al. 2007) and (2) widespread adoption of glyphosate-resistant (GR) crops.

Effective weed control with use of herbicides in the early to mid-1970s facilitated the adoption of NT production practices in the Great Plains (Hansen et al. 2012; Wicks and Smika 1973; Wicks et al. 1993). Weed control strategies shifted away from tillage and became more reliant on herbicides. However, small-seeded broadleaf weeds such as horseweed [*Conyza canadensis* (L.) Cronquist], kochia [*Bassia scoparia* (L.) A.J. Scott], Palmer amaranth (*Amaranthus palmeri* S. Watson), common lambsquarters (*Chenopodium album* L.), and Russian thistle (*Salsola tragus* L.) and grass weeds such as downy brome (*Bromus tectorum* L.), wild oat (*Avena fatua* L.), foxtail species (*Setaria* spp.), and tumble windmill grass (*Chloris verticillata* Nutt.) have become problematic weed species over time in NT production systems of the Great Plains (Buhler and Oplinger 1990; Jha et al. 2016; Nichols et al. 2015). This is mainly because these small-seeded weeds can easily germinate on or near the soil surface under NT fallow conditions. In addition, a majority of these weed species are also prolific seed producers and can quickly replenish the soil seedbank during the fallow period in the absence of crop competition. In contrast, populations of large-seeded broadleaf weeds such as velvetleaf (*Abutilon theophrasti* Medik.), common cocklebur (*Xanthium strumarium* L.), and *H. annuus* have declined over time in this region (Anderson et al. 1998; Buhler et al. 1996). The NT soil surface was probably not a biologically beneficial site for successful germination and emergence of large-seeded compared with small-seeded weed species (Buhler et al. 1996). Large seeds on the soil surface under NT conditions are also exposed to higher mortality risks through predation (insects, diseases, birds, animals, etc.) and weather variability (Baraiar et al. 2009; Nichols et al. 2015).

Nevertheless, the extensive and repeated use of glyphosate for weed control in fallow, pre-crop seeding, and postharvest (NT burndown) scenarios has led to the evolution of glyphosate resistance in several weed species in the region (Heap 2019; Jha et al. 2016; Stahlman 2016). This was partly due to glyphosate use at lower than recommended rates by growers to minimize cost of weed control in NT fallow. Currently, the reported cases of GR weed species across the Great Plains include *C. canadensis*, *B. scoparia*, *A. palmeri*, and *S. tragus* (Heap 2019). Among all these reported GR weed species, glyphosate resistance is widespread among *B. scoparia* populations across all the Great Plains states (Kumar et al. 2018a).

GR crops became available in the United States in 1996, allowing producers to use glyphosate for in-season control of broadleaf and grass weeds. By 2005, >90% of total U.S. acreage planted to soybean and cotton (*Gossypium hirsutum* L.) and nearly 50% of corn had the GR trait (Kniss 2018; Sankula 2006). The GR crops in the U.S. Great Plains occupy a significant portion of the total agricultural land [alfalfa (*Medicago sativa* L.), canola (*Brassica napus* L.), corn, cotton, soybean, and sugar beet (*Beta vulgaris* L.) being the most common GR crops]. The rapid adoption of GR crops, along with the dramatic increase in glyphosate use (almost 8-fold increase between 1996

and 2005), replaced preplant tillage, POST cultivation, and other selective herbicides as a means of weed control (Givens et al. 2009; Kniss 2018). This change in glyphosate use pattern contributed to weed species shifts in major agronomic crops in the United States (Johnson et al. 2009). For instance, Wilson et al. (2007) documented that two continuous in-crop treatments of glyphosate each year at 0.4 or 0.8 kg ae ha⁻¹ shifted *B. scoparia* and wild proso millet (*Panicum miliaceum* L.) populations to predominantly *C. album* populations in a long-term field study conducted in Scottsbluff, NE. Shaner (2000) concluded that reduced sensitivity in weed seedlings to glyphosate could occur more frequently than evolution of resistant biotypes, further shifting weed populations from susceptible to ones that are more tolerant.

Herbicide-Resistant Weeds

In NT fallow fields, growers primarily rely on multiple applications of herbicides (predominantly glyphosate-based tank mixtures) to achieve season-long weed control (Fenster and Wicks 1982; Moyer et al. 1994). Repeated use of herbicides with the same SOAs has resulted in the evolution of herbicide resistance in several weed species across the region. This section briefly reviews documented cases of herbicide resistance in common weed species across the U.S. Great Plains.

Bassia scoparia

A well-detailed review on chronology of herbicide resistance, distribution, mechanisms of resistance, seed germination and emergence characteristics, population dynamics, and IWM strategies to manage HR *B. scoparia* across North America has recently been published (Kumar et al. 2018a). The first case of HR *B. scoparia* was discovered in 1976, when atrazine-resistant biotypes were reported in cornfields in Kansas and along railroads in Idaho and Iowa (Heap 2019; Kumar 2018a; Stahlman 2016). *Bassia scoparia* resistant to acetolactate synthase (ALS)-inhibiting herbicides was first identified from a wheat field in Kansas in 1987 (Primiani et al. 1990). Currently, a majority of *B. scoparia* populations in the region are believed to be resistant to ALS-inhibiting herbicides (Kumar et al. 2018a). Cross-resistance to dicamba and/or fluroxypyr was identified in *B. scoparia* populations collected from wheat/fallow fields in Montana and North Dakota in 1995 (Cranston et al. 2001; Nandula and Manthey 2002). Currently, *B. scoparia* resistant to dicamba and fluroxypyr have been reported from six states: Montana, Idaho, North Dakota, Nebraska, Colorado, and Kansas (Heap 2019; Kumar et al. 2018a). Resistance to glyphosate was first confirmed in *B. scoparia* populations from W-S-F fields in western Kansas in 2007 and is currently reported from two-thirds of the U.S. Great Plains states and three Canadian provinces (Beckie et al. 2013; Godar et al. 2015a; Hall et al. 2014; Kumar et al. 2014; Waite et al. 2013; Wiersma et al. 2015). In addition, several *B. scoparia* populations across the region have been documented to have multiple resistance to two (glyphosate and ALS inhibitors), three (glyphosate, ALS inhibitors, and dicamba), or four (glyphosate, ALS inhibitors, dicamba, and atrazine) herbicide SOAs (Heap 2019; Kumar et al. 2015, 2019a; Stahlman 2016; Varanasi et al. 2015; Westra et al. 2019).

Amaranthus palmeri

Historically, *A. palmeri* is a common weed species in the mid-southern and southeastern United States. *A. palmeri* has recently become more problematic in the Southern and Central Great Plains (Figure 1) and is moving farther north (Ward et al. 2013). Ward et al. (2013) provided a detailed review on various

aspects of *A. palmeri*, including herbicide-resistance evolution in the United States. *Amaranthus palmeri* populations in Texas were confirmed resistant to atrazine in 1993, and in Kansas in 1995 (Heap 2019). Populations of *A. palmeri* resistant to ALS inhibitors and glyphosate are reported across the Southern and Central Great Plains (Figure 1) (Chahal et al. 2017; Garetson et al. 2019; Horak and Peterson 1995; Kumar et al. 2019b, 2020; Nakka et al. 2017). *Amaranthus palmeri* resistant to 4-hydroxyphenylpyruvate dioxygenase-inhibiting herbicides was first confirmed in Kansas in 2009, and subsequently in Nebraska and Texas (Garetson et al. 2019; Jhala et al. 2014; Singh et al. 2018). Additionally, *A. palmeri* populations with multiple resistance to two, three, four, or five herbicide SOAs have become an increasing concern in the Southern and Central Great Plains (Figure 1) (Garetson et al. 2019; Heap 2019; Jhala et al. 2014; Kumar et al. 2019b, 2020; Singh et al. 2018). More recently, an *A. palmeri* population with multiple resistance to 2,4-D, glyphosate, chlorsulfuron, mesotrione, and atrazine and reduced sensitivity to fomesafen has been reported in Kansas (Kumar et al. 2019b).

Salsola tragus

This C_4 summer annual is another troublesome broadleaf weed commonly found in semiarid regions of the Great Plains (Leeson et al. 2005). *Salsola tragus* is highly problematic in summer fallow periods and in spring/winter wheat across the Northern and Central Great Plains (Figure 1) (Schillinger 2007; Young 1986). Effective management of this weed species has been complicated by the evolution of herbicide resistance in several field populations (Heap 2019). Resistance to ALS inhibitors in *S. tragus* populations was first confirmed in Washington and Montana in 1987 (Guttieri et al. 1992; Heap 2019) and was subsequently reported in California, Idaho, Kansas, North Dakota, and Oregon (Heap 2019; Peterson 1999). Currently, >75% of the wheat fields in the NGP (Figure 1) are infested with ALS-resistant *S. tragus* populations (Jha et al. 2016; Saari et al. 1992; Stallings et al. 1994). Resistance to glyphosate in *S. tragus* populations has recently been confirmed from fallow fields of Montana, Washington, and Oregon and is further suspected in populations in the Northern and Central Great Plains (Barroso et al. 2018; Heap 2019; Kumar et al. 2017c).

Conyza Canadensis

A native species of North America, this winter annual dicot weed species commonly infests agronomic crops, orchards, pastures, roadsides, and industrial/waste areas (Gleason and Cronquist 1963; Miller and Miller 1999). *Conyza canadensis* favors NT production systems of the U.S. Great Plains, as the seedlings primarily emerge on the soil surface (Nandula et al. 2006). Glyphosate is the most common nonselective herbicide used for its control in NT production systems across the region. Resistance to glyphosate in *C. canadensis* has been reported in Montana, Nebraska, Kansas, and Oklahoma (Croese et al. 2019; Heap 2019; Kumar et al. 2017b). In addition to glyphosate resistance, *C. canadensis* populations with resistance to other herbicide SOAs, including photosystem I (PSI) inhibitors, ALS inhibitors, and PSII inhibitors, have been reported (Gadamski et al. 2000; Heap 2019; Mueller et al. 2003; Smisek et al. 1998).

Weed Management under Changing Climate

Climate change poses a serious concern to the dryland production systems of the U.S. Great Plains. Changing weather trends and extreme conditions in this region can have direct (changes in crop

growth, development, and yield) and indirect (strong selective pressures exerted by abiotic stresses and biotic stresses such as pests) effects on agriculture (Hartfield et al. 2014). The predicted changes in regional weather conditions (increasing CO_2 levels, warmer temperatures, and varying frequency and distribution of annual precipitation) would have a profound effect on invasion, distribution, establishment, composition, demography, fitness, competitiveness, and management of weed species (Bradley et al. 2010; Gritti et al. 2006; Varanasi et al. 2016; Waryszak et al. 2018; Ziska and Dukes 2011). The Great Plains region is projected to experience air temperatures increased by 2 to 4 °C, doubling of atmospheric CO_2 levels, warm winters, and pronounced droughts in some zones within the next half century (Hartfield et al. 2014; Rosenberg 1982). These climatic changes might influence the growth cycles of predominant weed species in the region, further suggesting the need of innovative and improved weed management practices in future. For instance, rising levels of atmospheric CO_2 will likely increase photosynthesis and growth of several weed species, further enhancing the ability of weeds to compete with crop plants (Ziska 2001, 2003). The process of converting CO_2 to sugar in C_3 plants is less efficient compared with C_4 plant species, suggesting that C_3 weed species will respond rapidly to increasing atmospheric CO_2 concentrations (Ogren and Chollet 1982; Ziska 2003). This increased efficiency of C_3 weed species under increasing CO_2 concentrations may result in aggressive growth, more competitive ability than other weeds or crops, and/or high seed production, indicating potential future weed shifts. Furthermore, changing weather patterns across the region, such as increasing CO_2 levels, warm winters, and changing duration and frequency of annual rainfall events, are likely to increase both the risks posed by, and the sources of, invasive weed species (Bradley et al. 2012). For instance, the northward spread of invasive weeds such as *A. palmeri*, which is highly problematic in the Southern and Central Great Plains, could potentially be a concern in the NGP (Figure 1) in near future (Ward et al. 2013).

Climate change can also influence the effectiveness of herbicides and make weed management more challenging. Environmental factors such as CO_2 , light, temperature, and relative humidity can influence the performance of different herbicide SOAs (Varanasi et al. 2016). For instance, glyphosate efficacy was reduced on weeds grown at high CO_2 levels (Koleva and Schneider 2009). Similarly, reduced efficacy of mesotrione on *A. palmeri*, glyphosate on *B. scoparia* and *C. album*, and dicamba on *B. scoparia* has been reported at high temperatures (40/30 °C) (DeGreeff et al. 2018; Godar et al. 2015b; Ou et al. 2018). These reports demonstrate that more frequent applications and higher rates of herbicides may be needed to obtain adequate weed control in the future, resulting in increased economic and environmental costs associated with herbicide use. Recent literature also suggests that global warming can induce genetic and phenotypic changes within weed populations and will pose a greater risk of evolution of metabolism-based, non-target site herbicide resistance in weed species (Matzrafi et al. 2016; Ziska et al. 2019).

Cover Crops in No-Tillage Regions of the Semiarid Great Plains

Effective weed management is one of the most important agronomic practices for producing a profitable crop (Baucom and Holt 2009). As fewer and fewer new herbicide modes of action come to market and HR weeds continue to increase in NT dryland production systems of the Great Plains, alternative weed control

strategies are needed. The implementation of infrequent tillage, including reduced tillage rather than NT and competitive crops grown in rotations, have contributed to the management of current weed problems; however, these practices alone may not be enough to achieve effective management of HR weeds in dryland production systems. Integration of CCs in dryland crop rotations is gaining popularity not only for weed control but also for other conservation practices. CCs contribute to building soil health and provide supplemental forage for grazing, nitrogen fixation, weed suppression, and habitat for beneficial insects and weed seed predators (Al-Khatib et al. 1997; Holman et al. 2018; Hoorman 2009; Kasper et al. 2007; Teasdale and Daughtry 1993).

Dryland producers are adopting CCs to suppress weeds and to reduce costs of managing HR weed populations. In the U.S. Great Plains, CCs can be planted in fallow phases of the cropping cycle between cash crops. Theoretically, CCs in place of those fallow phases can help manage weeds in two different ways: (1) through the competitive effect (competition for light, water, nutrients, and space) of CCs for reducing weed growth that would otherwise establish in fallow fields (Osipitan et al. 2018) and (2) through the suppressive effect of CC residues on weeds after termination of CCs (Pullaro et al. 2006). The suppressive effect of CCs can occur in two ways: (1) through physical suppression, with CC residue (mulching) blocking sunlight and altering the soil microclimate (Lemessa and Wakjira 2015); and (2) through chemical suppression, with CCs releasing allelochemical compounds into the environment (Al-Khatib et al. 1997; Barnes and Putnam 1986; Burgos et al. 1999; Creamer et al. 1996; Dhima et al. 2006; Ercoli et al. 2007; Teasdale and Daughtry 1993; White et al. 1989). However, the findings of many allelopathic studies have been confined to research laboratories, greenhouses, and growth chambers (Creamer et al. 1996; Sosnoskie et al. 2012) and need to be further investigated under field conditions.

In addition to weed suppression, CCs can play a key role in integrated pest management in this semiarid region. CCs are known to influence soil microbial communities, which in turn can affect the viability and persistence of weed seedbanks (Liebman et al. 2001). Kumar et al. (2008) reported that soil fungi contributed to the reduction of corn chamomile (*Anthemis arvensis* L.) and shepherd's purse [*Capsella bursa-pastoris* (L.) Medik.] seedling emergence from soil where a buckwheat CC was grown and incorporated. Similarly, specific soil fungi are known to cause mortality of germinating *A. theophrasti* seeds (Davis and Renner 2006). However, the long-term impacts of CCs on the soil microbiome are not fully understood and should be investigated under field conditions. There is insufficient research-based information on using CCs for weed suppression in the NT Great Plains, where soil moisture is limited compared with other U.S. regions. The following section reviews some opportunities for inclusion of CCs for weed management in semiarid cropping systems of the U.S. Great Plains.

Ecofriendly Approach and Key Elements of Integrated Pest Management

Growing CCs during fallow periods of NT dryland crop rotations across the Great Plains would serve as an ecofriendly approach for weed management. Effective weed management in NT fallow fields tends to rely on multiple herbicide applications (glyphosate alone or in mixtures) to provide season-long weed control each year. From 1974 to 2014, >1.6 billion kg (about 19% of estimated global use) of glyphosate was applied in U.S. agriculture (Benbrook 2016).

Due to the rapid evolution of GR weed populations, some growers have been replacing glyphosate with paraquat (a restricted-use herbicide due to its high level of acute human toxicity) in their burndown programs for fallow fields (VK and PWS, personal observation). The weed suppression achieved by CCs during fallow periods is likely to reduce the frequent and substantial use of these nonselective herbicides. The resulting decline in use of herbicides such as paraquat, dicamba, and atrazine could help reduce the chemical load on the environment (soil, water, air), further creating a more natural ecosystem.

CCs can also aid in increased weed seed predation and substantial reduction in weed seedbanks by harboring or providing suitable habitat for seed-eating insects, invertebrates, small rodents, and birds (Brust and House, 1988; Cardina et al. 1996; Maulsby 2006; Menalled et al. 2006, 2007; Westerman et al. 2005; White et al. 2007). Seed predation has been well documented for several weed species such as common ragweed (*Ambrosia artemisiifolia* L.), *A. palmeri*, *C. album*, sicklepod [*Senna obtusifolia* (L.) Irwin & Barneby], jimsonweed (*Datura stramonium* L.), *A. theophrasti*, fall panicum (*Panicum dichotomiflorum* Michx.), and giant foxtail (*Setaria faberi* Herrm.) (Menalled et al. 2006, 2007; Westerman et al. 2005; White et al. 2007). Ground beetles (*Anisodactylus* spp.), field crickets (*Gryllus pennsylvanicus* Burmeister), ants (*Formicidae* spp.), earthworms (*Lumbricina* spp.), slugs (*Gastropoda* spp.), field mice (*Peromyscus maniculatus* Wagner), and other small rodents hide below a residue layer on the soil surface for protection (Menalled et al. 2006; Westerman et al. 2005; White et al. 2007). Therefore, we hypothesize that replacing NT fallow fields with CCs could increase seed predation of problematic weed species in the semiarid regions of the Great Plains and achieve more effective weed control systems.

Cover Crops for Managing Herbicide-Resistant Weeds in the Great Plains

Raising CCs for weed suppression is not a new concept; however, this method of weed control is only recently being tested in NT dryland production systems of the Great Plains. Exploring CCs for weed management in the region will open several new opportunities for weed research. Future CC studies in this dryland environment could investigate how CC management strategies (species selection, seeding rate, time of planting, termination date, livestock grazing, etc.) will affect life-history traits (seed dormancy, germination, emergence, growth, reproduction, seed viability, seed persistence, etc.) of problem weed species. Previous studies have demonstrated that *B. scoparia* populations in fallow fields from Central Great Plains (Figure 1) had earlier and quicker emergence patterns compared with populations from the NGP (Dille et al. 2017; Kumar et al. 2017a). Rapid seed germination has also been observed in *B. scoparia* resistant to ALS inhibitors, whereas reduced and delayed seed germination has been reported in dicamba-resistant *B. scoparia* (Dyer et al. 1993; Kumar and Jha 2016; Kumar et al. 2018b). Differential seed germination characteristics have also been found for GR *B. scoparia* in the region (Kumar and Jha 2017; Osipitan and Dille 2017). Similarly, differential growth and reproductive characteristics of *B. scoparia* populations with various herbicide-resistance traits have previously been reported (Kumar and Jha 2015, 2016; LeClere et al. 2018; Martin et al. 2017; Osipitan and Dille 2017). It would be important to understand the fitness attributes of evolved herbicide-resistance traits in problematic weed species (*B. scoparia*, *A. palmeri*, *C. canadensis*, *S. tragus*, and *B. tectorum*) under diverse CC

management strategies in NT dryland cropping systems. This information will help in exploring the ecological fate of HR alleles in weed populations. For instance, if fall-planted CCs such as winter triticale (\times *Triticosecale* Wittm. ex A. Camus [*Secale* \times *Triticum*]), cereal rye (*Secale cereale* L.), and hairy vetch (*Vicia villosa* Roth) provide competitive effects that might reduce seed germination, emergence, growth, and reproduction of HR *B. scoparia* and *C. canadensis* populations, then these CCs could delay or mitigate further spread of HR alleles in these populations. Similarly, spring-planted CCs such as oat (*Avena sativa* L.), triticale, and pea (*Pisum sativum* L.) alone or in mixtures can produce enough biomass and sufficient canopy cover to provide competitive effects on growth and reproduction of HR *B. scoparia* populations (Obour et al. 2019a). However, only those CCs that produced more than 1,000 kg ha⁻¹ were effective at weed suppression (Petrosino et al. 2015), and greater CC biomass results in less available soil water for the subsequent cash crop (Holman and Obour 2019). The leftover residue upon the termination of these fall- or spring-planted CCs can also provide suppressive effects (physical mulching) on the emergence of *A. palmeri* populations during early summer in the fallow period of a W-S-F rotation. The suppression of HR *B. scoparia*, *C. canadensis*, and *A. palmeri* populations achieved by these fall- or spring-planted CCs can also reduce the number of POST herbicide applications needed during the fallow phase, which will ultimately reduce selection pressure for further evolution of herbicide-resistance traits in these species.

The long-term impacts of replacing fallow fields with CCs on soil seedbank dynamics of problem weed species (especially HR populations), as well as shifts in weed communities, will be other important research questions to explore. It would be crucial to investigate and identify key competitive traits of growing CCs in the region, including planting dates, mixtures versus single species, seeding rates, and growth attributes (above- and belowground), that can reduce the growth of predominant weed species, thus reducing weed seed production to ultimately deplete the soil seedbanks. Future research could also emphasize understanding the suppressive traits of growing CCs, including residue persistence (especially in this moisture-limited environment) that can help conserve the soil moisture, alter the soil microenvironment (temperature, moisture, microbes), and ultimately, impact weed seedbank life. In a similar context, field and laboratory studies could be designed to identify allelopathic traits of various CCs (single or multiple species) grown in a semiarid environment, including the nature of allelochemicals, timing of release and impact on weed seed germination, weed seed life, and microbes in the soil (Kelton et al. 2012; Weston 1996). Depending upon the goal of growing CCs (competitiveness vs. suppressiveness), multilocation studies should investigate different methods of termination (herbicides, mowing, haying, flailing, grazing, etc.) for different CC species, optimum timing, and their short- and long-term impacts on weed demographics and population dynamics under NT dryland production systems.

Economic and Sustainability Aspects of Using Cover Crops for Weed Management

On average, weed control costs >\$11 billion a year for U.S. agriculture, and herbicides are one of the major expenses (Hartfield et al. 2014). Weed suppression provided by CCs will potentially reduce herbicide use, which in turn will help reduce herbicide costs. Subsequently, any reductions in herbicide use during fallow periods will reduce the risk of herbicide-resistance evolution in

weed populations across the region. In addition, weed seedlings remain smaller and more susceptible to herbicides for longer periods of time under a CC canopy, which could indirectly enhance the effectiveness of chemical control of those small-sized seedlings at the time of CC termination (Wallace et al. 2019). The replacement of NT fallow periods with CCs that produce high-quality forage can provide additional revenue to growers if properly managed for livestock grazing/haying. A recent study in western Kansas and eastern Colorado found that a spring-planted CC mixture of oat/triticale/pea produced high-quality forage for early summer grazing (Brummer et al. 2018). In a separate study, the same CC mixture (oat/triticale/pea) provided 90% to 99% suppression of *B. scoparia* biomass compared with nontreated fallow plots across two different sites in western Kansas (Obour et al. 2019a). However, it is important to note that the CC was not grazed or harvested in that same study, so the overall impact of CC grazing on *B. scoparia* population is unknown (Obour et al. 2019a). Previous research found that CCs in NT fallow fields could help reduce soil compaction, soil erosion, and nutrient leaching (Blanco-Canqui et al. 2013). Legume CCs can also increase nitrogen fixation, improve soil aggregates, protect crop seedlings, and conserve soil moisture by reducing evaporation rates during drought periods through a thick layer of residue after termination (Basche et al. 2016; Holman et al. 2018; Magdof and Van Es 2009; Unger and Vigil 1998). All of these factors can potentially contribute to enhancing the economic viability and sustainability of semi-arid Great Plains cropping systems.

Agronomic Considerations for Integrating Cover Crops in No-Tillage Regions of the Semiarid Great Plains

The intended goal of including CCs in NT cropping systems will influence species selection, planting time, seeding rate, and time of termination. The extent of CC weed-suppressive ability will be greater with more biomass. In NT dryland systems, CC biomass production varies greatly because of variable precipitation amount, crop rotation, CC species, planting time, seeding rate, and soil type. In a recent meta-analysis, Osipitan et al. (2019) showed that CC management decisions, including selection of CC species, planting season, planting date, seeding rate, termination date, delay in main crop planting after termination of CCs, and tillage system, all can influence the effective use of CCs for weed suppression. Here, we briefly review major agronomic practices and challenges in growing CCs for weed suppression in NT semiarid regions of the U.S. Great Plains.

Cover Crop Species Selection

Choosing individual CC species or mixtures with vigorous growth and biomass production is critical for effective weed suppression. Previous research documents that several diverse grass and broad-leaf species could be planted as single or multiple species CCs in the semiarid Great Plains (Table 1). Cereals are generally considered more weed suppressive than broadleaf plant species (Norsworthy et al. 2011; Ruffo and Bollero 2003; Ruis et al. 2018). Cereal rye is a popular species chosen by those interested in incorporating a winter-hardy CC, as it can develop a fibrous root system, tolerates low-fertility soils, scavenges for available nitrogen, prevents soil erosion that commonly occurs when no residue or plant material is left on the soil surface, and can suppress weeds (Clark 2007). However, cereal rye also can quickly become “weedy,” known as feral rye, and is a weed of dryland agriculture in the western and central United States that causes more than \$26 million in annual wheat harvest losses (Pester et al. 2000; Western Coordinating Committee-077

Table 1. List of grass and broadleaf cover crop species commonly adapted in the Great Plains.

Grass species	Scientific names ^a
Barley	<i>Hordeum vulgare</i> L.
Cereal rye	<i>Secale cereale</i> L.
Oats	<i>Avena sativa</i> L.
Proso millet	<i>Panicum miliaceum</i> L.
Sudangrass	<i>Sorghum bicolor</i> L.
Triticale	× <i>Triticosecale</i> Wittm. ex A. Camus [<i>Secale</i> × <i>Triticum</i>]
Winter wheat	<i>Triticum aestivum</i> L.
Broadleaf species	
Austrian winter pea	<i>Pisum sativum</i> L.
Berseem clover	<i>Trifolium alexandrinum</i> Moench
Buckwheat	<i>Fagopyrum esculentum</i> L.
Common vetch	<i>Vicia sativa</i> L.
Flax	<i>Linum usitatissimum</i> L.
Hairy vetch	<i>Vicia villosa</i> Roth
Lentil	<i>Lens culinaris</i> Medik.
Phacelia	<i>Phacelia tanacetifolia</i> Benth.
Spring pea	<i>Pisum sativum</i> L.
Radish	<i>Raphanus sativus</i> L.
Canola	<i>Brassica napus</i> L.
Safflower	<i>Carthamus tinctorius</i> L.
Sunflower	<i>Helianthus annuus</i> L.
Sunn hemp	<i>Crotalaria juncea</i> L.

^aReviewed/researched by Calderon et al. 2016; Holman et al. 2018; Nielsen et al. 2015; Sanderson et al. 2018.

2019; Whitson et al. 2000). Barley (*Hordeum vulgare* L.), wheat, oat, and triticale are also competitive cereals and could possibly fit into some systems better than cereal rye. Although legumes are not considered as competitive as cereals and have a low C:N ratio (which contributes to a faster residue decomposition rate), their use may still offer benefits in the Great Plains (Creamer et al. 1997).

In field studies conducted in western Kansas, the most productive grass CCs in dryland environments were sorghum or sudangrass (*Sorghum bicolor* L.) > cereal rye > triticale > wheat > oat > barley (Holman et al. 2018). The broadleaf legume CCs were not competitive and produced little biomass when grown alone or in CC mixes. In southwest Kansas, fall-planted triticale and a triticale–hairy vetch mixture reduced *B. scoparia* density by 78% and 94%, respectively, and biomass up to 98% when compared with chemical fallow (Petrosino et al. 2015). In that study, legume CCs such as Austrian winter pea, hairy vetch, spring lentil, and spring pea grown as single-species CCs produced little biomass (600 to 1,000 kg ha⁻¹) and could not provide adequate suppression of *B. scoparia* growth. This was probably because the emergence timing of these spring-planted CCs coincided with *B. scoparia* emergence and the CCs could not compete well. This further suggests that weed emergence timing relative to CC establishment is a critical factor in selecting CC species, as previously discussed. Similarly, grass and brassica monocultures or mixtures were more effective at weed suppression than legumes (Baraibar et al. 2018). Therefore, to be effective, low biomass–producing legumes and other broadleaf CCs may need to be planted in mixtures with productive grass species to improve weed suppression. When weed suppression is the ultimate goal, then selecting a single CC species that produces more biomass can provide a cheaper weed-suppression alternative compared with multispecies mixes that may have a higher seed cost (Holman et al. 2018).

Planting multispecies CCs does not necessarily produce greater biomass or residue compared with a single species or simple CC mixes. Nielsen et al. (2015) reported CC biomass or residue after CC termination with a 10-species mixture was not different

compared with biomass produced by single-species CCs across two locations in eastern Colorado and western Nebraska. In North Dakota, weed-suppression benefits with monoculture CCs were not different from mixtures comprising millet [*Pennisetum glaucum* (L.) R. Br.]/triticale/clover (*Trifolium* spp.)/radish (*Raphanus sativus* L.) (Sanderson et al. 2018). Similarly, Florence et al. (2019) reported that multispecies CCs mixes did not increase aboveground biomass production or weed suppression when compared with productive single-species CCs across 11 sites in southeastern Nebraska. However, in a 2-yr study conducted in southcentral Montana, Khan and McVay (2019) concluded that multispecies CCs can provide more stable biomass yield than single-species CCs. In a meta-analysis, Osipitan et al. (2019) reported that the decision to use a combination of multispecies mixtures, single species, and grasses or broadleaves was driven by ability of the selected CCs to produce greater biomass and persistent residue to suppress weeds. In recent CC studies in western Kansas, it was found that actively growing CC plants were more effective at suppressing weeds compared with CC residue amounts (Obour et al. 2019a). Hence, extending CC growth over greater portions of the fallow period could provide more weed-suppression benefits, but caution should be taken to avoid depleting plant-available water, which could be detrimental to the subsequent cash crop.

Cover Crop Seeding Rates

This is an important agronomic decision that affects plant stand, growth, water use, competitiveness, and productivity of CCs. Seeding rates of CCs are highly variable, and depend on species and whether planted as single or multiple species (Calderon et al. 2016; Holman et al. 2018). Single-species CCs should be planted at seeding rates similar to those used for forage production. For example, the seeding rate for oat and triticale CCs grown in the Great Plains ranged from 45 to 100 kg ha⁻¹ (Calderon et al. 2016; Holman et al. 2018). Sudangrass is planted at 13.5 to 16.8 kg ha⁻¹, which is the recommended rate range for forage production. Typical seeding rates for multispecies CCs ranged from 40 to 60 kg ha⁻¹ (Calderon et al. 2016; Farney et al. 2018); significantly less compared with seeding rates for pea (115 to 134 kg ha⁻¹) or triticale alone (72 to 100 kg ha⁻¹) (Holman et al. 2018). Because of their smaller seed size, brassicas are generally planted at seeding rates of 5.6 to 7.5 kg ha⁻¹ as a single-species CC or reduced to 1.1 to 2.2 kg ha⁻¹ when included in CC mixes (Calderon et al. 2016; Farney et al. 2018).

There is limited research on CC seeding rate effect on weed suppressiveness in the semiarid Great Plains. However, studies in the mid-Atlantic region (Maryland, Pennsylvania, New York) showed increasing cereal rye seeding rate reduced the amount of weed biomass produced (Ryan et al. 2011). Weed biomass in that study ranged from 328 g m⁻² when rye was seeded at 90 kg seed ha⁻¹ compared with 225 g m⁻² when seeding rate was increased to 210 kg seed ha⁻¹. Haramoto (2019) reported a greater plant density of cereal rye and winter wheat CCs when seeded at 112 kg ha⁻¹ (184 plants m⁻²) compared with a lower seeding rate of 34 kg ha⁻¹ (70 to 81 plants m⁻²); however, CC biomass and weed suppression did not differ between the two seeding rates. Previous research in the Central Great Plains indicated that oat grown for forage could be planted at 25% less than the recommended seeding rate of 72 kg ha⁻¹ with no decrease in total biomass produced (Obour et al. 2019b). This is due to a high tillering ability of cereal grains that compensates for reduced seeding rates. Therefore, a moderate reduction in CC seeding rate may not negatively

influence total biomass production. Because of high seed costs associated with increasing seeding rates, CC growers should be moderate in deciding what seeding rates to use. The goal is to provide enough plant density and biomass to provide early-season weed suppression.

Cover Crop Seeding Time

In the semiarid Great Plains, CC growth and productivity varies from year to year because of the variability in precipitation amounts and seeding time. Average CC biomass in southwest Kansas ranged from 780 kg ha⁻¹ when precipitation during the CC growing season was 46% of the 30-yr average to 2,690 kg ha⁻¹ when precipitation was above average (Holman et al. 2018). The CC growing season can affect growth and subsequent impact on weed suppression. Spring- or fall-seeded CCs tend to perform better than a late summer-planted CC in this semiarid environment because of the available moisture at time of seeding. This period also coincides with the emergence of the most problematic weed species of this region: *C. canadensis* primarily emerges in the fall or early spring; *B. scoparia* emerges in early spring; whereas *A. palmeri* emerges in late spring to early summer. Therefore, depending upon soil moisture, fall- or spring-seeded CCs can best compete against and suppress problematic weed species in NT fields that otherwise would remain fallow and allow weed populations to flourish. For instance, fall-seeded cereal rye or winter triticale after wheat harvest can provide enough CC biomass to compete against *C. canadensis* and *B. scoparia*, which germinate in early spring. In contrast, spring-seeded CCs (such as oat/triticale/pea) in the fallow phase can help manage *A. palmeri* and *B. scoparia* in early summer through competition and late summer through physical suppression from CC residue following termination. Sanderson et al. (2018) found that spring-seeded CCs produced greater biomass and provided greater weed suppression compared with when CCs were seeded later in the growing season in North Dakota. This would be expected, because late-summer seedlings of CCs in semiarid regions of the Great Plains face considerable risks due to variable rainfall and a shortened growing season, which can result in reduction of CC biomass and weed-suppression benefits. Irrespective of the CC growing season, optimum planting dates are crucial for adequate plant establishment, increased biomass, and improved weed suppression.

Cover Crop Establishment Challenges

The ability of CCs to suppress weeds is contingent upon achieving good establishment. This can be challenging in semiarid environments where soil moisture at the time of seeding can be limited. Seeding into subsoil moisture will increase seedling emergence, plant vigor, and early establishment. However, CCs are generally seeded in mixtures, and the large number of plant species promoted as CCs exacerbate establishment issues because of differences in seed size within CC mixtures, preventing deep seed placement when smaller-seeded species are included in mixtures. To overcome this challenge, CC seeding should coincide with periods of adequate soil moisture to ensure conditions are favorable for germination and seedling emergence, particularly for small-seeded CC species that need to be seeded at shallow depths. Furthermore, limiting the number of species in CC mixtures to plants with similar seed sizes could ensure the entire CC mixture can be seeded at depths that provide good seed-to-soil contact to increase the chances of emergence and plant establishment.

In general, the CC seeding methods include direct planting into previous crop residue, aerial seeding, interseeding, or broadcasting into a standing crop (Curran et al. 2018; Holman et al. 2018; Wilson et al. 2014). Among these methods, direct seeding with a drill provided better establishment with uniform stands compared with aerial seeding (Holman et al. 2018; Wilson et al. 2014). Establishment was reduced with broadcast or aerial seeding because of nonuniform seed distribution, low seed-to-soil contact, lack of moisture, and seed predation (Wilson et al. 2014). Establishment of small-seeded CC species could be improved with some tillage to prepare a seedbed; however, such tillage could increase potential for wind erosion and encourage a flush of weeds. Research is needed to determine whether occasional tillage every 3 to 5 yr could be used to establish CCs as part of the crop rotation cycle. This approach can improve CC establishment and control troublesome weeds with minimal impacts on soil properties (Blanco-Canqui and Wortmann 2020). One-time strategic tillage in long-term NT fields in western Kansas controlled glyphosate-tolerant red three-awn grass (*Aristida purpurea* Nutt.) and *C. verticillata* with no effect on soil water availability, winter wheat yields, and soil properties (Obour et al. 2019c). However, more research is needed to determine the combined effects of strategic tillage and CCs in managing HR weeds and subsequent impacts on soil health and crop yields in this region.

Another major challenge that affects CC establishment is residual herbicide from the previous crop. In dryland W-S-F rotations, producers are currently relying on PRE herbicides to manage HR weeds. Many of these herbicide chemistries are persistent, with residual phytotoxic concentrations in the soil that may affect establishment when CCs are planted in the fallow phase of the rotation. For example, S-metolachlor applied at 1.68 kg ha⁻¹ severely injured and reduced annual ryegrass (*Lolium multiflorum* Lam.) CC stands regardless of the planting time following herbicide application (Tharp and Kells 2000). Similarly, biomass of Italian ryegrass [*Lolium perenne* L. ssp. *multiflorum* (Lam.) Husnot] and winter oat were significantly reduced (~67%) by pyroxasulfone herbicide carryover from the previous corn crop (Cornelius and Bradley 2017). In general, herbicides with greater persistence will have a long-term residual activity that will affect CC inclusion in the rotation. Herbicide labels typically provide information on restrictions regarding safety of rotational crops, but these are based on cash crops with no or limited information for CC usage. More research information is needed regarding herbicide carryover injury on CCs to prevent risk of establishment failure. In addition, data on the safety of the herbicide with respect to CCs, particularly CCs used as livestock feed, are essential.

Soil Moisture Trade-offs and Impacts on Cash Crop Yields

The fallow phase of the production system is critical for conservation of soil water that stabilizes wheat yields and prevents crop failure in drier years in NT regions of the semiarid Great Plains (Haas et al. 1974; Nielsen and Vigil 2010). In this water-limited environment, crop yields are directly related to available soil moisture at time of cash crop seeding (Schlegel et al. 2018). Despite the weed-suppression benefits of CCs, previous research demonstrated that replacing fallow with CCs often resulted in decreased soil water content at time of cash crop seeding (Holman et al. 2018; Nielsen and Vigil 2018; Schlegel and Havlin 1997; Unger and Vigil 1998; Zentner et al. 1996). Schlegel and Havlin (1997) reported that a hairy vetch CC reduced soil water content by 178 mm compared with fallow, and the subsequent wheat crop

yield was reduced by 42% to 83% across years. In that same study, every millimeter of soil water depleted by hairy vetch resulted in reduction of wheat yields by 15 kg ha⁻¹. Holman et al. (2018) reported wheat yield reduction of 13 kg ha⁻¹ for every millimeter of soil water depleted by CCs, whether grown as a mixture or as a single species. The impact of CCs on soil water availability will depend on amounts of precipitation and infiltration after termination to replenish water use by CCs. Therefore, growing window and time of CC termination are critical for CC water use and soil water replenishment relative to seeding the next cash crop. Wheat yields following lentil as a CC terminated at full bloom in northeastern Montana was 2,040 kg ha⁻¹ compared with 2,820 kg ha⁻¹ in fallow in a W-F system (Pikul et al. 1997). This corresponds to a more than 25% decrease in wheat yield when lentil replaced fallow in the cropping system. In eastern Colorado, Nielsen and Vigil (2005) reported that the soil water content in fallow was 320 mm compared with 265 mm when CCs were terminated in early June, 245 mm when terminated in mid-June, or 216 mm when terminated in mid-July. Barker et al. (2018) reported little to no effect of CCs on soil water availability in a 3-yr study across Nebraska.

In general, water use by CCs will vary and may depend on species, biomass production, and precipitation after termination in the growing season. In a 2-yr study conducted at a dryland site in Akron, CO, Nielsen et al. (2016) reported CC water use ranging from 127 mm for pea to 221 mm for canola and 136 mm in the first year (seasonal precipitation of 85 mm) to 202 mm in the second year (seasonal precipitation of 178 mm). The aforementioned studies demonstrate that growing CCs in the semiarid Great Plains uses soil water resources and could affect subsequent cash crop yields regardless of CC species selected, growing window (fall or spring planted), or selection of multispecies CCs. It is therefore imperative that CCs grown in semiarid regions provide some revenue stream to offset seed costs and cash crop yield depressions when CCs are seeded ahead of the main crop (Holman et al. 2018).

Cover Crops Grazing/Haying

An additional revenue stream for farmers is to use CCs for grazing or haying. This approach could provide an opportunity for dryland producers to suppress weeds, improve soil health, and produce harvestable forage for livestock. Recent research in western Kansas demonstrated that most of the species planted as CCs have excellent forage attributes in terms of dry matter production and forage nutritive value (Brummer et al. 2018; Obour et al. 2019b). Regrowth from hayed or grazed CCs can provide more residue cover compared with fallow. In southwestern Kansas, winter triticale or a winter triticale-legume CC harvested for forage was more profitable than chemical fallow (Holman et al. 2018). Notwithstanding the significant depression in wheat yields following a spring CC, overall system profitability was greater when an oat-pea forage was added to dryland cropping systems compared with fallow (Lyon et al. 2004). Therefore, using CCs for forage will provide opportunity for dual-purpose CCs in dryland systems to also provide residue cover to manage HR weeds, reduce erosion, and improve soil health. However, research data on best management practices for grazing CCs in NT dryland systems are limited. More importantly, farmers are asking questions about:

1. CC mixtures to achieve best outcomes;
2. seeding windows for grazing CCs (spring, summer, or fall);

3. amount of CCs biomass that should be grazed relative to residue cover;
4. grazing impact on soil compaction;
5. grazing impacts on weed suppression (weeds like *B. scoparia* may be palatable); and
6. soil health implications when CC biomass is removed as forage through grazing.

Currently, there is a paucity of research to address these concerns. Ongoing research demonstrates that grazing CCs could increase soil compaction near the soil surface under wet conditions. Wheat grain yields following CCs that were grazed or hayed did not differ from yields when CCs were left standing (Obour et al. 2019b). The grazing potential of CCs in semiarid dryland systems of the Great Plains needs further investigation.

Summary and Future Directions

The conundrum of incorporating CCs in crop rotations in the semiarid U.S. Great Plains is how to take advantage of weed suppression and gain soil health benefits while reducing the negative impacts on available soil water and subsequent crop yields. The potential detrimental effects of CCs on subsequent crop yields have slowed their adoption, because farmers' decisions are based on the overall economic situations of their farms. For successful adoption of CCs in NT dryland systems, CCs must be managed to increase the amount of biomass at termination for weed suppression and provide an economic value. Replacing NT fallow periods with CCs will potentially reduce the number of herbicide applications. This will hold true when there is adequate moisture for CC establishment to produce adequate biomass.

This review highlights that integration of CCs in NT regions of the semiarid Great Plains can potentially contribute to the development of cost-effective and ecologically based IWM strategies, especially for managing HR weed populations. Multistate studies are needed to investigate the systematic manipulation of specific components of CCs for successful integration into current cropping systems in the region. Research gaps exist in understanding the impacts of various CC management strategies (CC species selection, time of seeding and termination, seeding rates, method of termination, etc.) on population dynamics (seed mortality and longevity; seed dormancy; germination ecology; emergence pattern; and other life-history traits, e.g., fitness) of HR weed populations and subsequent crop yields in the region. Researchers should also investigate other ecosystem services provided by CCs in NT semiarid Great Plains, including allelopathy, attractants for beneficial insects/pollinators, and habitat for weed seed predators, with the ultimate goal of achieving more resilient weed management systems that are more likely to delay evolution of HR weed populations. Long-term field research should also validate the important components of environmental sustainability and economic viability of integrating CCs for weed management in the region.

Acknowledgments. This work was jointly supported by the USDA National Institute of Food and Agriculture (Hatch Projects 1019671 and 1019594) and the USDA North Central Sustainable Agricultural Research and Education Program (grant no. LNC 18-411). No conflicts of interest have been declared. This publication is contribution no. 20-104-J from the Kansas Agricultural Experiment Station, Manhattan, KS.

References

- Al-Khatib K, Libbey C, Boydston R (1997) Weed suppression with *Brassica* green manure crops in green pea. *Weed Sci* 45:439–445
- Anderson RL, Tanaka DL, Black AL, Schweizer EE (1998) Weed community and species response to crop rotation, tillage, and nitrogen fertility. *Weed Technol* 12:531–536
- Baraibar B, Mortensen DA, Hunter MC, Barbercheck ME, Kaye JP, Finney DM, Curran WS, Bunchek J, White CM (2018) Growing degree days and cover crop type explain weed biomass in winter cover crops. *Agron Sustain Dev* 38:65
- Baraibar B, Westerman PR, Carrión E, Recasens J (2009) Effects of tillage and irrigation in cereal fields on weed seed removal by seed predators. *J Appl Ecol* 46:380–387
- Barker JB, Heeren DM, Koehler-Cole K, Shapiro CA, Blanco-Canqui H, Elmore RW, Proctor CA, Irmak S, Francis CA, Shaver TM, Mohammed AT (2018) Cover crops have negligible impact on soil water in Nebraska maize–soybean rotation. *Agron J* 110:1718–1730
- Barnes JP, Putnam AR (1986) Evidence for allelopathy by residues and aqueous extracts of rye (*Secale cereale*). *Weed Sci* 86:384–390
- Barroso J, Goullie JA, Lucher LK, Mingyang L, Mallory-Smith CA (2018) Identification of glyphosate resistance in *Salsola tragus* in Northeastern Oregon. *Pest Manag Sci* 74:1089–1093
- Basche AD, Kasparb TC, Archontoulisa SV, Jaynesb DB, Sauerb TJ, Parkinb TB, Miguez FE (2016) Soil water improvements with the long-term use of a winter rye cover crop. *Agric Water Manag* 172:40–50
- Baucom RS, Holt JS (2009) Weeds of agricultural importance: bridging the gap between evolutionary ecology and crop and weed science. *New Phytol* 184:741–743
- Beckie HJ, Blackshaw RE, Low R, Hall LM, Sauder CA, Martin S, Brandt EN, Shirriff SW (2013) Glyphosate- and acetolactate synthase inhibitor-resistant kochia (*Kochia scoparia*) in western Canada. *Weed Sci* 61:310–318
- Benbrook CM (2016) Trends in glyphosate herbicide use in the United States and globally. *Environ Sci Eur* 28:3
- Blanco-Canqui H, Holman JD, Schlegel AJ, Tatarko J, Shaver TM (2013) Replacing fallow with cover crops in a semiarid soil: effects on soil properties. *Soil Sci Soc Am J* 77:1026–1034
- Blanco-Canqui H, Wortmann CS (2020) Does occasional tillage undo the ecosystem services gained with no-till? A review. *Soil Till Res* 198, 10.1016/j.still.2019.104534
- Bradley BA, Blumenthal DM, Early R, Grosholz ED, Lawler JJ, Miller LP, Sorte CJB, D'Antonio CM, Diez JM, Dukes JS, Ibanez I, Olden JD (2012) Global change, global trade, and the next wave of plant invasions. *Front Ecol Environ* 10:20–28
- Bradley BA, Blumenthal DM, Wilcove DS, Ziska LH (2010) Predicting plant invasions in an era of global change. *Trends Ecol Evol* 25:310–318
- Brummer J, Johnson S, Obour A, Caswell K, Moore A, Holman J, Schipanski M, Harman K (2018) Managing Spring Planted Cover Crops for Livestock Grazing under Dryland Conditions in the High Plains Region. Fort Collins: Colorado State University Extension Fact Sheet No. 0.309
- Brust GE, House GJ (1988) Weed seed destruction by arthropods and rodents in low-input soybean agroecosystems. *Am J Alternative Agr* 3:19–25
- Buhler D, Mester T, Kohler K (1996) The effect of maize residues and tillage on emergence of *Setaria faberi*, *Abutilon theophrasti*, *Amaranthus retroflexus* and *Chenopodium album*. *Weed Res* 36:153–165
- Buhler DD, Oplinger ES (1990) Influence of tillage systems on annual weed densities and control in solid-seeded soybean (*Glycine max*). *Weed Sci* 38:158–165
- Burgos NR, Talbert RE, Mattice JD (1999) Cultivar and age differences in the production of allelochemicals by *Secale cereale*. *Weed Sci* 47:481–485
- Calderon FJ, Nielsen D, Acosta-Martínez V, Vigil MF, Lyon D (2016) Cover crop and irrigation effects on soil microbial communities and enzymes in semiarid agroecosystems of the Central Great Plains of North America. *Pedosphere* 26:192–205
- Cardina, J, Norquay HM, Stinner BR, McCartney DA (1996) Postdispersal predation of velvetleaf (*Abutilon theophrasti*) seeds. *Weed Sci* 44:534–539
- Chahal PS, Varanasi VK, Jugulam M, Jhala AJ (2017) Glyphosate-resistant Palmer amaranth (*Amaranthus palmeri*) in Nebraska: confirmation, EPSPS gene amplification, and response to POST corn and soybean herbicides. *Weed Technol* 31:80–93
- Clark A (2007) Managing Cover Crops Profitably. 3rd ed. Beltsville, MD: Sustainable Agriculture Network. 244 p
- Cornelius CD, Bradley KW (2017) Carryover of common corn and soybean herbicides to various cover crop species. *Weed Technol* 31:21–31
- Cranston HJ, Kern AJ, Hackett JL, Miller EK, Maxwell BD, Dyer WE (2001) Dicamba resistance in kochia. *Weed Sci* 49:164–170
- Creamer NG, Bennett MA, Stinner BR (1997) Evaluation of cover crop mixtures for use in vegetable production systems. *HortScience* 32:866–870
- Creamer NG, Bennett MA, Stinner BR, Cardina J, Regnier EE (1996) Mechanisms of weed suppression in cover crop-based production systems. *HortScience* 31:410–413
- Croze JA, Manuchehri MR, Baughman TA (2019) Horseweed (*Conyza canadensis*) management in Oklahoma winter wheat. *Weed Technol* 34:229–234
- Curran WS, Hoover RJ, Mirsky SB, Roth GW, Ryan MR, Ackroyd VJ, Wallace JM, Dempsey MA, Pelzer CJ (2018) Evaluation of cover crops drill interseeded into corn across the mid-Atlantic region. *Agron J* 110:435–443
- Davis AS, Renner KA (2006) Influence of seed depth and pathogens on fatal germination of velvetleaf (*Abutilon theophrasti*) and giant foxtail (*Setaria faberi*). *Weed Sci* 55:30–35
- DeGreeff RD, Varansai AV, Dille JA, Peterson DE, Jugulam M (2018) Influence of plant growth stage and temperature on glyphosate efficacy in common lambsquarters (*Chenopodium album*). *Weed Technol* 32:448–453
- Dhima KV, Vasilakoglou IB, Eleftherohorinos IG, Lithourgidis AS (2006) Allelopathic potential of winter cereals and their cover crop mulch effect on grass weed suppression and corn development. *Crop Sci* 46:345–352
- Dille JA, Stahlman PW, Du J, Geier PW, Riffel JD, Currie RS, Wilson RG, Sbatella GM, Westra P, Kniss AR, Moechnig MJ, Cole RM (2017) Kochia emergence profiles across the Central Great Plains. *Weed Sci* 65:614–625
- Dyer WE, Chee PW, Fay PK (1993) Rapid germination of sulfonylurea-resistant *Kochia scoparia* (L.) Schrad. accession is associated with elevated seed levels of branched chain amino acids. *Weed Sci* 41:18–22
- Ercoli L, Masoni A, Pampana S, Arduini I (2007) Allelopathic effects of rye, brown mustard and hairy vetch on redroot pigweed, common lambsquarter and knotweed. *Allelopathy J* 19:249–256
- Farney JK, Sassenrath GF, Davis CJ, Presley D (2018) Composition, forage production, and costs are variable in three-way cover crop mixes as fall forage. *Crop Forage Turfgrass Manag* 4:1–7
- Fenster CR, Wicks GA (1982) Fallow systems for winter wheat in western Nebraska. *Agron J* 74:9–13
- Florence AM, Higley LG, Drijber RA, Francis CA, Lindquist JL (2019) Cover crop mixture diversity, biomass productivity, weed suppression, and stability. *PLoS ONE* 14:e0206195
- Gadamski G, Ciarka D, Gressel J, Gawronski SW (2000) Negative cross-resistance in triazine-resistant biotypes of *Echinochloa crusgalli* and *Conyza canadensis*. *Weed Sci* 48:176–180
- Garetson R, Singh V, Singh S, Dotray P, Bagavathiannan M (2019) Distribution of herbicide-resistant Palmer amaranth (*Amaranthus palmeri*) in row crop production systems in Texas. *Weed Technol* 33:355–365
- Ghimire R, Ghimire B, Mesbah AO, Idowu OJ, O'Neill MK, Angadi SV, Shukla MK (2018) Current status, opportunities, and challenges of cover cropping for sustainable dryland farming in the Southern Great Plains. *J Crop Improv* 34:1–20
- Givens WA, Shaw DR, Kruger GR, Johnson WG, Weller SC, Young BG, Wilson RG, Owen MD, Jordan D (2009) Survey of tillage trends following the adoption of glyphosate-resistant crops. *Weed Technol* 23:150–155
- Gleason HA, Cronquist A (1963) Manual of Vascular Plants of Northeastern United States and Adjacent Canada. Boston, MA: PWS Publishers. 734 p
- Godar AS, Stahlman PW, Jugulam M, Dille JA (2015a) Glyphosate-resistant kochia (*Kochia scoparia*) in Kansas: EPSPS gene copy number in relation to resistance levels. *Weed Sci* 63:587–595
- Godar AS, Varanasi VK, Nakka S, Prasad PV, Thompson CR, Mithila J (2015b) Physiological and molecular mechanisms of differential sensitivity of palmer amaranth (*Amaranthus palmeri*) to mesotrione at varying growth temperatures. *PLoS ONE* 10:e0126731

- Gritti E, Smith B, Sykes MT (2006) Vulnerability of Mediterranean basin ecosystems to climate change and invasion by exotic plant species. *J Biogeogr* 33:145–157
- Guttieri MJ, Eberlein CV, Mallory-Smith CA, Thill DC, Hoffman DL (1992) DNA sequence variation in Domain A of the acetolactate synthase genes of herbicide-resistant and -susceptible weed biotypes. *Weed Sci* 40:670–676
- Haas HJ, Willis WO, Bond JJ (1974) Summer Fallow in the Western United States. USDA-ARS Conservation Research Report No. 17. Washington, DC: U.S. Government Printing Office. 172 p
- Hall LM, Beckie HJ, Low R, Shirriff SW, Blackshaw RE, Kimmel N, Neeser C (2014) Survey of glyphosate-resistant kochia (*Kochia scoparia* L. Schrad.) in Alberta. *Can J Plant Sci* 94:127–130
- Hansen N, Allen B, Baumhardt RL, Lyon D (2012) Research achievements and adoption of no-till, dryland cropping in the semi-arid U.S. Great Plains. *Field Crops Res* 132:196–203
- Haramoto ER (2019) Species, seeding rate, and planting method influence cover crop services prior to soybean. *Agron J* 111:1068–1078
- Hartfield JL, Antle J, Garrett KA, Izaurrealde RC, Mader T, Marshall E, Nearing M, Robertson GP, Ziska L (2014) Indicators of climate change in agricultural systems. *Clim Change*. <https://doi.org/10.1007/s10584-018-2222-2>
- Heap I (2019) The International Survey of Herbicide Resistant Weeds. <http://www.weedscience.org>. Accessed: September 26, 2019
- Holman JD, Arnet K, Dille J, Maxwell S, Obour A, Roberts T, Roozeboom K, Schlegel A (2018) Can cover or forage crops replace fallow in the semiarid Central Great Plains? *Crop Sci* 58:932–944
- Holman JD, Obour A (2019) Cover crop use in a semi-arid wheat-sorghum-fallow cropping system. Proceedings of the 2019 American Society of Agronomy–Crop Science Society of America–Soil Science Society of America (ASA–CSSA–SSSA) International Annual Meeting, November 10–14, 2019, San Antonio, TX
- Hoorman JJ (2009) Using Cover Crops to Improve Soil and Water Quality. Lima: Ohio State University Extension. <https://ohioline.osu.edu/factsheet/anr-57>. Accessed: September 26, 2019
- Horak MJ, Peterson DE (1995) Biotypes of Palmer amaranth (*Amaranthus palmeri*) and common waterhemp (*Amaranthus rudis*) are resistant to imazethapyr and thifensulfuron. *Weed Technol* 9:192–195
- Jha P, Kumar V, Lim CA (2016) Herbicide resistance in cereal production systems of the US Great Plains. *Indian J Weed Sci* 48:112–116
- Jhala AJ, Sandell LD, Rana N, Kruger GR, Knezevic SZ (2014) Confirmation and control of triazine and 4-hydroxyphenylpyruvate dioxygenase-inhibiting herbicide-resistant Palmer amaranth (*Amaranthus palmeri*) in Nebraska. *Weed Technol* 28:28–38
- Johnson WG, Davis VM, Kruger GR, Weller SC (2009) Influence of glyphosate-resistant cropping systems on weed species shifts and glyphosate-resistant weed populations. *Eur J Agron* 31:162–172
- Kasper TC, Jaynes DB, Parkin TB, Moorman TB (2007) Rye cover crop and gamagrass strip effects on NO₃ concentration and load in tile drainage. *J Environ Qual* 36:1503–1511
- Kelton J, Price AJ, Mosjidis J (2012) Allelopathic weed suppression through the use of cover crops. Pages 115–130 in Price A, ed. *Weed Control*. Shanghai: InTech China
- Khan QA, McVay KA (2019) Productivity and stability of multi-species cover crop mixtures in the Northern Great Plains. *Agron J* 111:1817–1827
- Koleva NG, Schneider UA (2009) The impact of climate change on the external cost of pesticide applications in US agriculture. *Int J Agric Sustain* 7:203–216
- Kniiss AR (2018) Genetically engineered herbicide-resistant crops and herbicide-resistant weed evolution in the United States. *Weed Sci* 66:260–273
- Kumar V, Brainard DC, Bellinder RR (2008) Suppression of Powell amaranth (*Amaranthus powellii*), shepherd's-purse (*Capsella Bursa-pastoris*), and corn chamomile (*Anthemis arvensis*) by buckwheat residues: role of nitrogen and fungal pathogens. *Weed Sci* 56:271–280
- Kumar V, Currie RS, Jha P, Stahlman PW (2019a) First report of kochia (*Bassia scoparia*) with cross-resistance to dicamba and fluroxypyr in western Kansas. *Weed Technol* 33:335–341
- Kumar V, Jha P (2015) Growth and reproduction of glyphosate resistant and susceptible populations of *Kochia scoparia*. *PLoS ONE* 10:e0142675
- Kumar V, Jha P (2016) Differences in germination, growth, and fecundity characteristics of dicamba-fluroxypyr-resistant and susceptible *Kochia scoparia*. *PLoS ONE* 11:e0161533
- Kumar V, Jha P (2017) Effect of temperature on germination characteristics of glyphosate-resistant and glyphosate-susceptible kochia (*Kochia scoparia*). *Weed Sci* 65:361–370
- Kumar V, Jha P, Dille JA, Stahlman PW (2017a) Emergence dynamics of kochia (*Kochia scoparia*) populations from the U.S. Great Plains: a multi-site-year study. *Weed Sci* 66:25–35
- Kumar V, Jha P, Giacomini D, Westra E, Westra P (2015) Molecular basis of evolved resistance to glyphosate and acetolactate synthase-inhibitor herbicides in kochia (*Kochia scoparia*) accessions from Montana. *Weed Sci* 63:758–769
- Kumar V, Jha P, Jhala AJ (2017b) Confirmation of glyphosate-resistant horseweed (*Conyza canadensis*) in Montana cereal production and response to POST herbicides. *Weed Technol* 31:799–810
- Kumar V, Jha P, Jugulam M, Yadav R, Stahlman PW (2018a) Herbicide-resistant kochia (*Bassia scoparia*) in North America: a review. *Weed Sci* 67:4–15
- Kumar V, Jha P, Lim CA, Stahlman PW (2018b) Differential germination characteristics of dicamba-resistant kochia (*Bassia scoparia*) populations in response to temperature. *Weed Sci* 66:721–728
- Kumar V, Jha P, Reichard N (2014) Occurrence and characterization of kochia (*Kochia scoparia*) accessions with resistance to glyphosate in Montana. *Weed Technol* 28:122–130
- Kumar V, Liu R, Boyer G, Stahlman PW (2019b) Confirmation of 2,4-D resistance and identification of multiple resistance in a Kansas Palmer amaranth (*Amaranthus palmeri*) population. *Pest Manag Sci* 75:2925–2933
- Kumar V, Liu R, Stahlman PW (2020) Differential sensitivity of Kansas Palmer amaranth populations to multiple herbicides. *Agron J*. <https://doi.org/10.1002/agj2.20178>
- Kumar V, Spring JF, Jha P, Lyon DJ, Burk IC (2017c) Glyphosate-resistant Russian thistle (*Salsola tragus*) identified in Montana and Washington. *Weed Technol* 31:238–251
- LeClere S, Wu C, Westra P, Sammons RD (2018) Cross-resistance to dicamba, 2,4-D, and fluroxypyr in *Kochia scoparia* is endowed by a mutation in an AUX/IAA gene. *Proc Natl Acad Sci USA* 115:E2911–E2920
- Leeson JY, Thomas AG, Hall LM, Brenzil C, Andrews T, Brown KR, Van Acker RC (2005) Prairie Weed Surveys of Cereal, Oilseed and Pulse Crops from the 1970s to the 2000s. Saskatoon, SK, Canada: Agriculture and Agri-Food Canada Weed Survey Series Publ 05–1. 395 p
- Lemessa F, Wakjira M (2015) Cover crops as a means of ecological weed management in agroecosystems. *J Crop Sci Biotechnol* 18:123–135
- Lenzen AW, Johnson GD, Carlson GR (2007) Cropping sequence and tillage system influence annual crop production and water use in semiarid Montana. *Field Crops Res* 100:32–43
- Liebman M, Mohler CL, Staver CP (2001) *Ecological Management of Agricultural Weeds*. New York: Cambridge University Press
- Lyon DJ, Blumenthal JM, Burgener PA, Harveson RM (2004) Eliminating summer fallow reduces winter wheat yields, but not necessarily system profitability. *Crop Sci* 44:855–860
- Magdof F, Van Es H (2009) *Building Soils for Better Crops: Sustainable Soil Management*. 3rd ed. Waldorf, MD: Sustainable Agriculture Publications. 294 p
- Martin SL, Benedict L, Sauder CA, Wei W, Da Costa LO, Hall LM, Beckie HJ (2017) Glyphosate resistance reduces kochia fitness: comparison of segregating resistant and susceptible F2 populations. *Plant Sci* 261:69–79
- Matzrafi M, Seiwert B, Reemtsma T, Rubin B, Peleg Z (2016) Climate change increases the risk of herbicide-resistant weeds due to enhanced detoxification. *Planta* 244:1217–1227
- Maulsby D (2006) *Free Weed Control Service: Mice*. Kutztown, PA: Rodale Institute. <http://www.newfarm.org/features/2006/0306/weedcontrol/maulsby.shtml>. Accessed: September 26, 2019
- Menalled FD, Liebman M, Renner K (2006) The ecology of weed seed predation in herbaceous crop systems. Pages 297–327 in Singh HP, Batish DR, Kohli RK, eds. *Handbook of Sustainable Weed Management*. New York: Food Products Press

- Menalled FD, Smith RG, Dauer JT, Fox TB (2007) Impact of agricultural management on carabid communities and weed seed predation. *Agr Ecosyst Environ* 118:49–54
- Miller JH, Miller KV (1999) Horseweed. Auburn, AL: Craftmaster Printers. 42 p
- Miller PR, Gan Y, Mcconkey BG, McDonald CL (2003) Pulse crops for the Northern Great Plains: I. Grain productivity and residual effects on soil water and nitrogen. *Agron J* 95:972–979
- Moyer JR, Roman ES, Lindwall CW, Blackshaw RE (1994) Weed management in conservation tillage systems for wheat production in North and South America. *Crop Prot* 4:243–259
- Mueller TC, Massey JH, Hayes RM, Main CL, Stewart CN Jr (2003) Shikimate accumulation in both glyphosate-sensitive and glyphosate-resistant horseweed (*Conyza canadensis* L. Cronq.). *J Agric Food Chem* 51:680–684
- Nakka S, Thompson CR, Peterson DE, Jugulam M (2017) Target-site-based and non-target site based resistance to ALS inhibitors in Palmer amaranth (*Amaranthus palmeri*). *Weed Sci* 65:681–689
- Nandula VK, Eubank TW, Poston DH, Koger CH, Reddy KN (2006) Factors affecting germination of horseweed (*Conyza canadensis*). *Weed Sci* 54:898–902
- Nandula VK, Manthey FA (2002) Response of kochia (*Kochia scoparia*) inbreds to 2,4-D and dicamba. *Weed Technol* 16:50–54
- Nichols V, Verhulst N, Cox R, Govaerts B (2015) Weed dynamics and conservation agriculture principles: a review. *Field Crop Res* 183:56–68
- Nielsen DC (2018) Influence of latitude on the US Great Plains east–west precipitation gradient. *Agric Environ Lett* 3:1–5
- Nielsen DC, Lyon DJ, Hergert GW, Higgins RK, Calderon FJ, Vigil MF (2015) Cover crop mixtures do not use water differently than single-species plantings. *Agron J* 107:1025–1038
- Nielsen DC, Lyon DJ, Hergert GW, Higgins RK, Holman JD (2016) Cover crop effect on subsequent wheat yield in the Central Great Plains. *Agron J* 108:243–256
- Nielsen DC, Vigil MF (2005) Legume green fallow effect on soil water content at wheat planting and wheat yield. *Agron J* 97:684–689
- Nielsen DC, Vigil MF (2010) Precipitation storage efficiency during fallow in wheat-fallow systems. *Agron J* 102:537–543
- Nielsen DC, Vigil MF (2018) Wheat yield and yield stability of eight dryland crop rotations. *Agron J* 110:594–601
- Norsworthy JK, McClelland M, Griffith G, Bangarwa SK, Still J (2011) Evaluation of cereal and brassicaceae cover crops in conservation-tillage, enhanced, glyphosate-resistant cotton. *Weed Technol* 25:6–13
- Obour AK, Chen C, Sintim HY, McVay K, Lamb P, Obeng E, Mohammed YA, Khan Q, Afshar RK, Zheljazkov VD (2018) *Camelina sativa* as a fallow replacement crop in wheat based crop production systems in the US Great Plains. *Ind Crops Prod* 111:22–29
- Obour AK, Holman JD, Dille JA, Kumar V (2019a) Effects of spring-planted cover crops on weed suppression and winter wheat grain yield in western Kansas. *Kansas Agricultural Experiment Station Research Reports* 5(6), [10.4148/2378-5977.7784](https://doi.org/10.4148/2378-5977.7784)
- Obour AK, Holman JD, Schlegel AJ (2019b) Seeding rate and nitrogen application effects on oat forage yield and nutritive value. *J Plant Nutr* 42:1452–1460
- Obour AK, Holman JD, Schlegel AJ (2019c) Strategic tillage in dryland no-tillage crop production systems. *Kansas Agricultural Experiment Station Research Reports* 5(6), [10.4148/2378-5977.7756](https://doi.org/10.4148/2378-5977.7756)
- Ogren WL, Chollet R (1982) Photorespiration. Pages 191–230 in Govindjee, ed. *Photosynthesis*. Cambridge, MA: Academic Press
- Osipitan OA, Dille JA (2017) Fitness outcomes related to glyphosate resistance in kochia (*Kochia scoparia*): what life history stage to examine? *Front Plant Sci* 8:1090
- Osipitan OA, Dille JA, Assefa Y, Knezevic SZ (2018) Cover crop for early season weed suppression in crops: systematic review and meta-analysis. *Agron J* 110:2211–2221
- Osipitan OA, Dille JA, Assefa Y, Radicetti E, Ayeni A, Knezevic SZ (2019) Impact of cover crop management on level of weed suppression: a meta-analysis. *Crop Sci* 59:833–842
- Ou J, Thompson CR, Stahlman PW, Blowdown N, Jugulam M (2018) Reduced translocation of glyphosate and dicamba in combination contributes to poor control of *Kochia scoparia*: evidence of herbicide antagonism. *Sci Rep* 8, [10.1038/s41598-018-23742-3](https://doi.org/10.1038/s41598-018-23742-3)
- Paulsen GM, Shroyer JP (2008) The early history of wheat improvement in the Great Plains. *Agron J* 100:S70–S78
- Pester TA, Westra P, Anderson RL, Lyon DJ, Miller SD, Stahlman PW, Northan FE, Wicks GA (2000) *Secale cereale* interference and economic thresholds in winter *Triticum aestivum*. *Weed Sci* 48:720–727
- Peterson DE (1999) The impact of herbicide-resistant weeds on Kansas agriculture. *Weed Technol* 13:632–635
- Peterson GA, Westfall DG (2004) Managing precipitation use in sustainable dryland agroecosystems. *Ann Appl Biol* 144:127–138
- Petrosino JS, Dille JA, Holman JD, Roozeboom KL (2015) Kochia suppression with cover crops in southwestern Kansas. *Crop Forage Turfgrass Manage* 1:1–8
- Pikul JL Jr, Aase JK, Cochran VL (1997) Lentil green manure as fallow replacement in the semiarid Northern Great Plains. *Agron J* 89:867–874
- Primiani MM, Cotterman JC, Saari LL (1990) Resistance of kochia (*Kochia scoparia*) to sulfonylurea and imidazolinone herbicides. *Weed Technol* 4:169–172
- Pullaro TC, Marino PC, Jackson DM, Harrison HF, Keinath AP (2006) Effects of killed cover crop mulch on weeds, weed seeds, and herbivores. *Agric Ecosyst Environ* 115:97–104
- Rosenberg NJ (1982) The increasing CO₂ concentration in the atmosphere and its implication on agricultural productivity II. Effects through CO₂-induced climatic change. *Clim Change* 4:239–254
- Ruffo ML, Bollero GA (2003) Modeling rye and hairy vetch residue decomposition as a function of degree-days and decomposition-days. *Agron J* 95:900–907
- Ruis S, Blanco H, Burr C, Olson B, Reiman M, Rudnick D (2018) Corn residue baling and grazing impacts on soil carbon stocks and other properties on a Haplustoll. *Soil Sci Soc Am J* 82:202–213
- Ryan MR, Curran WS, Grantham AM, Hunsberger LK, Mirsky SB, Mortensen DA, Nord EA, Wilson DO (2011) Effects of seeding rate and poultry litter on weed suppression from a rolled cereal rye cover crop. *Weed Sci* 59:438–444
- Saari LL, Cotterman JC, Smith WF, Primiani MM (1992) Sulfonylurea herbicide resistance in common chickweed, perennial ryegrass and Russian thistle. *Pestic Biochem Physiol* 42:110–118
- Sanderson M, Johnson H, Hendrickson J (2018) Cover crop mixtures grown for annual forage in a semi-arid environment. *Agron J* 110:525–534
- Sankula S (2006) Quantification of the Impacts on U.S. Agriculture of Biotechnology Derived Crops Planted in 2005. Washington, DC: National Center for Food and Agricultural Policy. 110 p. <http://www.ncfap.org/documents/2005biotechimpacts-finalversion.pdf>. Accessed: September 19, 2019
- Schillinger WF (2007) Ecology and control of Russian thistle (*Salsola iberica*) after spring wheat harvest. *Weed Sci* 55:381–385
- Schlegel AJ, Havlin JL (1997) Green fallow for the Central Great Plains. *Agron J* 89:762–767
- Schlegel AJ, Lamm FR, Assefa Y, Stone LR (2018) Dryland corn and grain sorghum yield response to available soil water at planting. *Agron J* 110:236–245
- Shafer M, Ojima D, Antle JM, Kluck D, McPherson R, Peterson S, Scanlon B, Sherman K (2014) Great Plains climate change impacts in the United States. The Third National Climate Assessment. Washington, DC: U.S. Global Change Research Program. Pp 441–461. <http://nca2014.globalchange.gov/report/regions/great-plains>. Accessed: September 21, 2019
- Shaner DL (2000) The impact of glyphosate-tolerant crops on the use of other herbicides and on resistance management. *Pest Manag Sci* 56:320–326
- Singh S, Burgos NR, Singh V, Alcober EAL, Salas-Perez R, Shivrain V (2018) Differential response of Arkansas Palmer amaranth (*Amaranthus palmeri*) to glyphosate and mesotrione. *Weed Technol* 32:579–585
- Smika DE, Wicks GA (1968) Soil water storage during fallow in the Central Great Plains as influenced by tillage and herbicide treatments. *Soil Sci Soc Am J* 32:591–595

- Smisek A, Doucet C, Jones M, Weaver S (1998) Paraquat resistance in horseweed (*Conyza canadensis*) and Virginia pepperweed (*Lepidium virginicum*) from Essex County, Ontario. *Weed Sci* 46:200–204
- Sosnoskie LM, Grey TL, Culpepper AS, Webster TM (2012) Allelopathy: hope or hype? Pages 76–78 in Collins G, Li C, Shurley D, eds. Cotton Research and Extension Report 2011. Athens: University of Georgia College of Agricultural and Environmental Sciences
- Stahlman PW (2016) Herbicide resistance in kochia: from single to multiple resistance. *Indian J Weed Sci* 48:117–121
- Stallings GP, Thill DC, Mallory-Smith CA (1994) Sulfonylurea resistant Russian thistle (*Salsola iberica*) survey in Washington State. *Weed Technol* 8:258–264
- Swanton CJ, Clements DR, Derksen DA (1993) Weed succession under conservation tillage: a hierarchical framework for research and management. *Weed Technol* 7:286–297
- Teasdale JR, Daughtry CST (1993) Weed suppression by live and desiccated hairy vetch (*Vicia villosa*). *Weed Sci* 41:207–212
- Tharp BE, Kells JJ (2000) Effect of soil-applied herbicides on establishment of cover crop species. *Weed Technol* 14:596–601
- Unger PW, Baumhardt RL (2001) Historical development of conservation tillage in Southern Great Plains. In Stiegler JH, ed. Proceedings of the 24th Annual Southern Conservation Tillage Conference for Sustainable Agriculture, Oklahoma City, 9–11 July 2001. Oklahoma City: Oklahoma State University
- Unger PW, Vigil MF (1998) Cover crop effects on soil water relationships. *J Soil Water Conserv* 53:200–207
- Varanasi A, Prasad PVV, Jugulam M (2016) Impact of climate change factors on weeds and herbicide efficacy. *Adv Agron* 135:107–146
- Varanasi VK, Godar AS, Currie RS, Dille AJ, Thompson CR, Stahlman PW, Jugulam M (2015) Field-evolved resistance to four modes of action of herbicides in a single kochia (*Kochia scoparia* L. Schrad.) population. *Pest Manag Sci* 71:1207–1212
- Waite J, Thompson CR, Peterson DE, Currie RS, Olson BLS, Stahlman PW, Khatib KA (2013) Differential kochia (*Kochia scoparia*) populations response to glyphosate. *Weed Sci* 61:193–200
- Wallace JM, Curran WS, Mortensen DA (2019) Cover crop effects on horseweed (*Erigeron canadensis*) density and size inequality at the time of herbicide exposure. *Weed Sci* 67:327–338
- Ward SM, Webster TM, Steckel LE (2013) Palmer amaranth (*Amaranthus palmeri*): a review. *Weed Technol* 27:12–27
- Waryszak P, Lenz TI, Leishman MR, Downey PO (2018) Herbicide effectiveness in controlling invasive plants under elevated CO₂: sufficient evidence to rethink weeds management. *J Environ Manag* 226:400–407
- Westerman PR, Liebman M, Menalled FD, Heggenstaller AH, Hartzler RG, Dixon PM (2005) Are many little hammers effective? Velvetleaf (*Abutilon theophrasti*) population dynamics in two- and four-year crop rotation systems. *Weed Sci* 53:382–392
- Western Coordinating Committee-077 (2019) Managing Invasive Weeds in Wheat. <https://www.nimss.org/projects/view/mrp/outline/18614>. Accessed: September 26, 2019
- Weston LA (1996) Utilization of allelopathy for weed management in agroecosystems. *Agron J* 88:860–866
- Westra EP, Nissen SJ, Getts TJ, Westra P, Gaines TA (2019) Survey reveals frequency of multiple resistance to glyphosate and dicamba in kochia (*Bassia scoparia*). *Weed Technol* 33:664–672
- White RH, Worsham D, Blum U (1989) Allelopathic potential of legume debris and aqueous extracts. *Weed Sci* 37:674–679
- White SS, Renner KA, Menalled FD, Landis DA (2007) Feeding preferences of weed seed predators and effect on weed emergence. *Weed Sci* 55:606–612
- Whitson TD, Burrill LD, Dewey SA, Cudney DW, Nelson BE, Lee RD, Parker R, eds (2000) Weeds of the West. 9th ed. Jackson, WY: Western Society of Weed Science
- Wicks GA, Martin AR, Mahnken GW (1993) Control of triazine-resistant kochia (*Kochia scoparia*) in conservation tillage corn (*Zea mays*). *Weed Sci* 41:225–231
- Wicks GA, Smika DE (1973) Chemical fallow in a winter wheat-fallow rotation. *Weed Sci* 21:97–102
- Wiersma AT, Gaines TA, Preston C, Hamilton JP, Giacomini D, Buell CR, Leach JE, Westra P (2015) Gene amplification of 5-enol-pyruvylshikimate 3-phosphate synthase in glyphosate resistant *Kochia scoparia*. *Planta* 241:463–474
- Wilson ML, Allen DL, Baker JM (2014). Aerially seeding of cover crops in the northern US Corn Belt: limitations, future research needs, and alternative practices. *J Soil Water Conserv* 69:67A–72A
- Wilson RG, Miller SD, Westra P, Kniss AR, Stahlman PW, Wicks GW, Kachman SD (2007) Glyphosate-induced weed shifts in glyphosate-resistant corn or a rotation of glyphosate-resistant corn, sugarbeet, and spring wheat. *Weed Technol* 21:900–909
- Young FL (1986) Russian thistle (*Salsola iberica*) growth and development in wheat (*Triticum aestivum*). *Weed Sci* 34:901–905
- Zentner RP, Campbell CA, Biederbeck VO, Selles F (1996) Indianhead black lentil as green manure for wheat rotations in the Brown soil zone. *Can J Plant Sci* 76:417–422
- Ziska LH (2001) Changes in competitive ability between a C4 crop and a C3 weed with elevated carbon dioxide. *Weed Sci* 49:622–627
- Ziska LH (2003) Evaluation of yield loss in field sorghum from a C3 and C4 weed with increasing CO₂. *Weed Sci* 51:914–918
- Ziska LH, Blumenthal DM, Franks SJ (2019) Understanding the nexus of rising CO₂, climate change, and evolution in weed biology. *Invasive Plant Sci Manag* 12:79–88
- Ziska LH, Dukes JS (2011) *Weed Biology and Climate Change*. Ames, IA: Wiley. 248 p