

Temporal variation in native bee diversity in the tropical Sub-deciduous forest of the Yucatan Peninsula, Mexico

Authors: Ramírez, Virginia Meléndez , Ayala, Ricardo, and González, Hugo Delfín

Source: Tropical Conservation Science, 9(2) : 718-734

Published By: SAGE Publishing

URL: <https://doi.org/10.1177/194008291600900210>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Research Article

Temporal variation in native bee diversity in the tropical sub-deciduous forest of the Yucatan Peninsula, Mexico

Virginia Meléndez Ramírez^{1*}, Ricardo Ayala² and Hugo Delfín González¹

¹Departamento de Zoología, Campus de Ciencias Biológicas y Agropecuarias, Universidad Autónoma de Yucatán, AP 4-116, Col. Itzimná, 97100 Mérida, Yucatan, México. ²Estación de Biología Chamela, Instituto de Biología, Universidad Nacional Autónoma de México (UNAM), AP 21, 48980 San Patricio, Jalisco, México. E-mail: *virginia.melendez@correo.uady.mx

*Corresponding author

Abstract

Native bee community diversity is known to vary temporally in the Neotropics, but little research has been done to quantify the extent of this variation. We studied monthly variation in native bee diversity in tropical semi-deciduous forest on the Yucatan Peninsula, Mexico, over a twelve-month period in 2007. Hurricane Dean traversed the study area in August of that year, causing some changes in the environment and in the native bee community. A total of 2,335 individual bee specimens were collected, belonging to 102 species from four families. The family Apidae had the highest number of species (46%) and individuals (71%). Species richness, abundance, and diversity varied throughout the study period. Species proportions of lifestyles and nesting behaviors varied from month to month, and absolute numbers changed. Low to moderate similarity values indicated high monthly species turnover, with almost half of species present for only one to two months. Species richness declined after the hurricane, with a loss of 40% of the species present beforehand, but nine new species appeared in September. The hurricane also reduced the number of solitary, parasocial, and social species, mostly those nesting in preexisting cavities and wood. Species composition in August had similarity values nearer to those of months in the dry season than to those in the rainy season. Two months after the hurricane, the native bee community had returned to previous species diversity levels, probably due to the rapid recovery of the vegetation on which the bee community depends. Effective biodiversity conservation programs will need more data on changes in vegetation community structure and how such changes influence Neotropical native bee communities after natural disturbances.

Key words: bee diversity, Hurricane Dean, Yucatan Peninsula, Neotropics, Mexico.

Resumen

Es conocido que la diversidad de la comunidad de abejas nativas en el neotropico varia temporalmente, y poco ha sido hecho para cuantificar la extensión de esta variación. La variación mensual de la diversidad de abejas nativas fue estudiada en la selva mediana subcaducifolia en la Península de Yucatán,

en México, en un periodo de 12 meses en 2007. El Huracán Dean atravesó el área de estudio en agosto de ese año, causando algunos cambios sobre el ambiente y la comunidad de abejas nativas. En total se recolectaron 2,335 individuos, de 102 especies pertenecientes a cuatro familias. La familia Apidae presentó el mayor número de especies (46%) e individuos (71%). La riqueza, abundancia y diversidad de especies vario a través de los meses. La proporción de especies en términos de estilos de vida y conductas de anidación vario mes a mes y el número de especies cambio. La baja a moderada similitud indico alto recambio de especies mensual y cerca del 50% de las especies estuvieron presentes solo uno o dos meses. La riqueza de especies declino después del paso del huracán, se perdieron el 40% de las especies presentes en agosto aunque nueve especies aparecieron como nuevas en septiembre. Después del huracán decrecieron especies solitarias, parasociales y sociales y principalmente las especies que anidan en cavidades preexistentes y en la madera. Agosto fue más similar en la composición de especies con los meses de secas que con los de lluvias. Dos meses después, presumiblemente la recuperación rápida de la vegetación posiblemente jugo un papel importante para la persistencia de la comunidad de abejas en el área de estudio. Se requieren más investigaciones para determinar los cambios en la estructura de la comunidad vegetal y los efectos en las comunidades de abejas neotropicales que permitan predecir el efecto de estos fenómenos naturales para la planeación de la conservación de la biodiversidad.

Palabras clave: diversidad de abejas, huracán Dean, Neotrópico, Península de Yucatán, México.

Received: 18 November 2015; **Accepted** 4 April 2016; **Published:** 27 June 2016

Copyright: Virginia Meléndez Ramírez, Ricardo Ayala and Hugo Delfín González. This is an open access paper. We use the Creative Commons Attribution 4.0 license <http://creativecommons.org/licenses/by/3.0/us/>. The license permits any user to download, print out, extract, archive, and distribute the article, so long as appropriate credit is given to the authors and source of the work. The license ensures that the published article will be as widely available as possible and that your article can be included in any scientific archive. Open Access authors retain the copyrights of their papers. Open access is a property of individual works, not necessarily journals or publishers.

Cite this paper as: Meléndez, R. V., Ayala, R. and Delfín, G. H. 2016. Temporal variation in native bee diversity in the tropical sub-deciduous forest of the Yucatan Peninsula, Mexico. *Tropical Conservation Science* Vol. 9 (2): 718-734. Available online: www.tropicalconservationscience.org

Disclosure: Neither Tropical Conservation Science (TCS) or the reviewers participating in the peer review process have an editorial influence or control over the content that is produced by the authors that publish in TCS.

Introduction

Native bee community diversity in the Neotropics is still an open field, since many zones have yet to be studied [1], particularly in Mexico [2]. Little is known about temporal variation in bee community diversity, which has received only occasional attention [*e.g.*, 3, 4]. In addition, no reports exist on the changes that natural phenomena such as hurricanes can cause in these bee communities. Native bee diversity had received minimal attention until the last twenty years, especially on the Yucatan Peninsula [2, 5, 6, 7]. Other studies have focused on bee diversity in some regional agricultural crops [8, 9], or on how habitat fragmentation has affected native bee diversity [10]. Most studies have been done in the northern portion of the Peninsula, and little is known about diversity in its south central portion, where intact remnants of sub-deciduous tropical forest remain, mainly in the area where the states of Yucatan, Campeche and Quintana Roo meet [11, 12].

The Yucatan Peninsula (YP) is a biotic province with Neotropical biogeographic affinity. Its climate is largely sub-humid. Because it is highly and uniformly exposed to the sun year round, there are only two clear seasons: a rainy season from late spring to early fall, and a dry season from late fall to early spring (the driest months are usually February to April). The Atlantic hurricane season officially spans from May to November, but the YP is most frequently affected during August and September. A common pattern for hurricanes that affect the YP is for a tropical storm to form east of the West Indies, to become a hurricane as it enters the Caribbean Sea, and then to sweep across the YP and into the Gulf of Mexico [13]. Relatively common in Mexico and Central America, hurricanes can cause dramatic changes in tropical forest structure when they make landfall, including generalized defoliation, loss of branches, trunk breakage, and the toppling of entire trees [14]. The almost complete defoliation caused by hurricanes can disrupt the canopy's role in providing shade and controlling soil microclimates [15], as well as rapidly increase the amount of leaf cover and wood detritus on the forest floor [16, 17]. Studies of this post-hurricane dynamic in the YP have only recently begun [*e.g.*, 18]. Although tropical forests frequently experience hurricanes, only limited attention has been given to the response of insects to such extreme natural disturbances [19]. Exacerbating this lack of data is that only a few tropical forest insect communities have been thoroughly evaluated [*e.g.* 20]. In contrast, a number of studies have documented the effects of anthropic disturbances on insects, particularly native bee communities [*e.g.*, 21].

Despite their being vital pollinators, no previous reports have been done on native bee communities' responses to high-energy natural phenomena such as hurricanes. To survive, native bee communities require floral resources and nesting sites, normally vegetation and other substrates. The food needs of each bee species in a community vary according to their feeding habits (generalist or specialist), social habits (*e.g.*, solitary, social, cleptoparasitic), nesting behavior, and life cycles [22, 23, 1]. Thus, their highly variable individual biology probably determines how each native bee species responds to large natural disturbances such as hurricanes. We describe monthly variation in the diversity of native bees in semi-deciduous forest on the Yucatan Peninsula before and after a hurricane affected the forest. These data will help to explain the response of the bee fauna to this type of natural phenomenon and its effect on bee ecological services.

Methods

Study area. The study area is located in the southern extreme of the state of Yucatan, in the municipality of Tekax ($19^{\circ}47'04''$ and $19^{\circ}43'03''$ N, $89^{\circ}18'13''$ and $89^{\circ}09'31''$ W), very near the point at which the Mexican states of Yucatan, Campeche and Quintana Roo meet, known as the Punto PUT (*i.e.* Punto de Unión Territorial) (Fig. 1). The area's climate is type Awo(i)gw, hot sub-humid with summer rains [24], with annual rainfall ranging from 1,000 to 1,229 mm [11]. Mean annual temperature is 24.9°C ; the hottest month is May, and the rainiest is September [25]. Soils in the area are vertisol, gleisol, nitisol, rendzinas, and lithosols [26]. Vegetation type is tropical sub-deciduous forest (TSF), which occurs as a wide band through the center of the state of Yucatan, narrowing to the south. It continues into northern Campeche and occupies only a small area in Quintana Roo, on its border with Yucatan. Trees in the TSF reach an average height of 10 to 20 m, although maximum height can be 25 to 30 m in some areas. During the dry season, typically 50% to 75% of trees in the TSF drop their leaves.

The main tree species in the TSF are *Acacia pennatula* (Schlech and Benth), *Caesalpinia gaumeri* Grenm., *Caesalpinia platyloba* S.Watson., *Lysiloma latisiliquum* (L.) Benth., *Enterolobium cyclocarpum* (Jacq.) Griseb., *Mimosa bahamensis* Benth., *Spondias mombin* L., *Metopium brownie* (Jacq.) Urban., *Cochlospermum vitifolium* (Willd) Spreng., *Guazuma ulmifolia* Lam., *Trema micrantha* (L.) Blume., *Annona reticulata* L., *Gyrocarpus americanus* Jacq., *Piscidia piscipula* (L.) Sarg., *Pithecellobium dulce* (Roxb) Benth. and *Pithecellobium albicans* (Kunth) Benth. Epiphytes in the area include some Areaceae (*Anthurium tetragonum* Schott), Bromeliaceae (*Tillandsia brachycaulos* Schltdl) and Orchidaceae (*Catasetum integerrimum* Hook) [11].

Hurricane Dean reached category 5 on the Saffir-Simpson scale off the southeast coast of the YP on 21 August 2007 (Fig. 1). As it crossed the YP from east to west and moved over the study area, it varied from category 4 to 3 [27, 28, 25].




Fig. 1. Sampling area location in Tekax Municipality (black dot ●), Yucatan Peninsula, Mexico. Trajectory of Hurricane Dean is marked with ☄ (symbol size indicates relative intensity from 5 to 1)[modified from 28].

Bee sampling. The native bee fauna was sampled monthly from January to December 2007. In August, sampling was done before Hurricane Dean traversed the area. Every month, sampling was done in 100 x 200 m quadrants, each quadrant using 15 McPhail traps [29], baited with 50% honey and placed at 10 m intervals; three Malaise traps placed at 50 m intervals; and three lots (12 traps per lot) of yellow pan traps placed at 50 m intervals. The three types of traps were left to collect specimens for 48 hours a month. During this same time, a research team member surveyed each quadrant for two hours to collect bees with an entomological aerial net. Bees were also collected by one team member along four 1-km-long transects. Collected bees were preserved in 70% ethanol, and later prepared and dried using a conventional treatment for Hymenoptera. Specimen identification was done following Ayala [30] and Michener [1], and corroborated by comparison with bee specimens in the Regional Entomological Collection of the Autonomous University of Yucatan (CER-UADY). Some individual specimens were established as “morphospecies” [*sensu* 31], because a number of bee genera have not been taxonomically reviewed. All collected specimens were deposited in the CER-UADY (Collecting permit-SEMARNAT- authorization code: SGPA/DGVS/ 10220).

Data analysis. We estimated species richness and total abundance by month. Representation of species richness values throughout the year was calculated with the quantitative Chao 2 richness estimator, one of the least biased estimators for small samples [32]. We calculated diversity using the proportional abundance indices classified as evenness indices (Shannon-Wiener). A random test was applied to the Shannon-Wiener index to detect differences among data, and calculations were done with the programs Biodiversity Pro [33] and Species Diversity and Richness [34]. The number of singletons and doubletons (species represented by only one or two bees, respectively, in the sample) was determined using the EstimateS program [35]. Environmental variable data were mean monthly temperature and rainfall during 2007, collected at the federal meteorological station nearest the study area (Nohalal Station, Tekax Municipality) [25]. Any correlations between environmental variables (temperature and rainfall) and bee species richness and abundance were identified with the Statistica program [36]. Native bee richness, abundance and diversity, as well as lifestyles and nesting behavior, were compared among the sampling data sets.

The Jackard index was used to evaluate shared species month to month; we used a cluster analysis to identify gathered species. The Jackard measures of beta diversity (month-to-month species turnover) were done using the Unweighted Pair Group Method with Arithmetic Mean (UPGMA) with the MVSP program [37]. Species lifestyle was defined based on several studies [23, 22, 38, 1]: solitary; parasocial (*sensu lato*: quasisocial, semisocial, subsocial, primitively eusocial), eusocial and cleptoparasitic. Nesting behavior was similarly defined [23, 1, 22]: preexisting cavities (nests in holes in rocks, banks of soil, dead tree limbs, etc.); nests excavated in soil; and nests excavated in hard or soft wood. For many native bee species, no data are available on their lifestyles and nesting behavior; when necessary, these parameters were deduced by considering taxonomic relationships at the genus and subgenus levels. The honeybee, *Apis mellifera* L. was not sampled because its presence, abundance and distribution depend more on the location of beehives than on environmental factors [39].

Results

Collected specimens totaled 2,335 bees, from 102 species belonging to four families. The family Apidae had the highest number of species and individuals ($\approx 45\%$ were stingless bees), followed by the Megachilidae, Halictidae and Colletidae families (Table 1). Estimated richness for the study area was 122 species, meaning our sample accounted for 85% of estimated richness, and was therefore highly representative.

Table 1. Number of species and total abundance by family of the native bee community in tropical semi-deciduous forest, Yucatan Peninsula, Mexico.

Family	Number of species	%	Number of individuals	%
Apidae	46	45.1	1660	71.0
Colletidae	6	6.0	37	1.6
Halictidae	23	22.5	466	20.0
Megachilidae	27	26.4	172	7.4
Total	102	100	2355	100

Rainfall was highest in the months following Hurricane Dean (September and October) (Fig. 2A). Species richness and total abundance were variable across months. The highest richness was in April and August, while the strongest declines were from June to July, and from November to December. The strongest decline in abundance was from October to November, during the rainy season. The number of individuals increased from August to September, and then again in October, two months after the hurricane (Fig. 2B). Forty-three species were documented in August, of which 17 (40%) were not found in September. However, nine species appeared in September that had not been recorded in August (26% of September's species). Twelve species present in August did not return in the months after the hurricane (Appendix 1). Diversity (H') was highest in April and August, declined in September and October, and then rose again in November (Fig. 2C). Values differed between August and September, were moderately variable in other months, and were lowest in January. Overall, 22 singleton species and 12 doubleton species were identified.

Lifestyles in the sampled native bee community were mostly solitary (50%), followed by parasocial (30%), eusocial (14%) and cleptoparasitic (6%) (Fig. 3A). Nesting in preexisting cavities was the most common strategy, followed by nesting in the ground, in wood, and the cleptoparasites (Fig. 3B). The number of species in each functional group changed from month to month, but the proportions remained constant (Fig. 3C and D). The number of solitary species was highest in April, and decreased steadily in later months, particularly in September and October. The lowest number of solitary species occurred in the months of January and February. Parasocial species were most frequent in August, declined notably in number from August to September, and remained constant from November to December. Present every month, eusocial species were most frequent in March and July, and declined from March to April.

Cleptoparasite species were most numerous in April, very much like the solitary species, but were not recorded in June and August. They reappeared after the hurricane (Fig. 3C).

Species nesting in cavities were most numerous in April and August, and then declined notably from June to July and again from August to September. Ground nesting species exhibited the highest numbers in November, and then declined to December. Those nesting in wood reached their maximum numbers in August, with declines in September and from June to July; their lowest numbers were in January (Fig. 3D).




Fig. 2. Total monthly rainfall and average monthly temperature (A), species richness and total monthly abundance (B), and diversity (H') of bees (C) in the study area. Figure C: letters above bar indicate differences between months ($p < 0.05$).

Overall, the month-to-month bee community species profile had generally low to moderate similarity (20 to 54%) (Fig. 4). Two groups had 20% similarity (or 80% dissimilarity): the first was present in January-December, and the second in the remaining months, including post-hurricane. Similarity was moderate from August to September and highest from September to October. Indeed, species composition in August (rainy season) had more similarity to compositions in the dry months of June and March than to other months in the rainy season (September and October) (Fig. 4).




Fig. 3. Lifestyles (A) and nesting behavior (B) in total native bee community; Lifestyles (C) and nesting behavior (D) over time. Figure C: So=Solitary, Ps=Parasocial, Eu= Eusocial and Cl= Cleptoparasitic and figure D: Cl= Cleptoparasitic, C= Preexisting cavity, G= Ground and W=Wood.




Fig. 4. Similarity dendrogram of native bee species activity in the months of 2007.

Positive correlations were identified between abundance and monthly rainfall ($r = 0.71$, $p < 0.5$), and between richness and monthly temperature ($r = 0.68$, $p < 0.5$). Bee species exhibited different activity periods throughout the year. Forty-nine species (48%) were caught in only one or two months, 29 species (28.5%) were present during three to six months, and the remaining 24 species (23.5%) were recorded during seven to twelve months (Fig. 5).




Fig. 5. Number of native bee species and number of months they were present during 2007.

Discussion

Apidae family species were the main components of the native bee community in the studied TSF, and had the highest number of recorded individuals (mainly eusocial stingless bees). This corresponds with reports for bee communities in the Neotropics [22, 40] and in other zones of the YP [5, 6, 7]. The number of recorded species represented 85% of the estimated number of species in the studied community, meaning approximately twenty more species have yet to be recorded. Complementary sampling would be needed to record the less common species that fill out bee community diversity in the Neotropics [40, 41]. Nearly 30% of the recorded species were rare (singletons and doubletons), a trend observed in other areas of the YP [5].

Weather in the YP area in 2007 was highly unusual. Rainfall in February was atypically high (this month is normally quite dry), and Hurricane Dean crossed the study area in August. The latter was an extremely high-energy event, a category 5 hurricane (wind speed: ≥ 252 km/h) when it made landfall [27], and the zone it affected was declared a disaster area [28]. When it crossed the study area, Hurricane Dean varied from category 3 to 4 (wind speed: ≥ 178 -249 km/h), and clearly impacted the TSF.

Impacts of this degree could have had negative effects on the area's native bee community. However, identifying which effects are specifically due to the hurricane is difficult, since native bee species' richness, diversity, and abundance exhibit wide monthly variation in response to several factors (*e.g.*, environmental conditions and availability of resources)[22]. Abundance increased after Hurricane Dean due to the presence of several species of stingless bees and other species (Appendix 1) that were apparently unaffected by the disturbance. Abundance peaked in October, two months after Dean, possibly indicating that vegetation (particularly

herbaceous species) quickly recovered after the event [14], and those trees and bushes that flower at this time [45] continued to do so. In addition to variations in response to resource availability [22, 39], native bee abundance patterns can also vary within and between years due to changes in environmental conditions [46, 22]. This was corroborated in the present study, as species richness was correlated to temperature, and abundance to rainfall, highlighting these factors' role in Neotropical bee activity.

Floral resources and nesting site availability are the two factors that most influence bee species' seasonal activity [42]. Bee abundance and frequency on flowers of specific plants are known to vary in response to annual flowering patterns [43]. Temporal variation in habitats occurs because important resources become available at different times [44]. Monthly variation in bee community species profiles in the studied TSF was high, with many species active for only a few months. This may be a common pattern in neotropical forests, since the same trend was reported in a study done in Chamela (Jalisco, Mexico) [4]. In our study, we observed seasonal changes in bee species composition and numbers; many were active during a few months and this activity was associated with flowering in the plants on which they feed. The presence of Cleptoparasite species is closely related to the activity of their hosts [22]. Our results confirm this, as their presence coincided with the activity of their host species (*e.g. Coelioxys* and *Megachile* species) (Appendix 1). Eusocial species are active year round [1], but they were not observed every month, except *Trigona fulviventris*, which was present in all twelve months (Appendix 1). These species are the most abundant in tropical bee communities, and are these communities' main components [22].

Since local and regional environmental conditions and biotic interactions influence native bee community composition, community structure and function exhibit marked spatial changes among locations in different zones worldwide [47]. However, only infrequent analyses have been done of temporal variation in bee communities, and this limited data can lead to erroneous conclusions about diversity in different kinds of habitat [44], and in response to natural disturbances such as hurricanes. In the TSF studied here, species composition in the native bee community varied notably throughout the year, mainly in response to the rainy and dry seasons. The dry season begins in November [13], and the lower temperature and rainfall in this season reduced bee numbers, consequently lowering bee abundance; indeed these two parameters were correlated. Richness remained relatively high in November, suggesting that species turnover exhibited marked changes between seasons. July had the lowest species richness and very low abundance, marking the transition from the dry to the rainy season. The rainy season commonly begins in May, but in 2007, both May and June were unusually dry. August saw increased rainfall, leading to the appearance of several species and the highest species numbers.

Diversity (H') varied throughout the year, but was highest in April. Similar levels can normally be expected during the rainy season, but the hurricane's impact reduced diversity in September and October. It did not fully recover until November. Hurricane Dean apparently altered the normal annual diversity pattern by displacing 17 of the species present before its impact, and introducing nine species afterwards, perhaps due to physical movement by high winds.

Community species composition also changed, losing species in the rainy season (*i.e.*, August had low similarity compared to September and October). Some of these lost species had been present in samples for months beforehand (12 species, *e.g.*, the solitary species *Megachile azteca*) (Appendix 1), but they did not reappear afterwards.

There are apparently species in the study area that occur for one generation annually in which adult bee activity occurs during two months or less. Examples of these species include *Gaesischia exul* and *Megachile texensis*, both active in March and April, or *Tetrapedia maura*, *Triepeolus cameroni* and *Heriades variolosus*, all active in April and May (Appendix 1). More research needs to be done on each of these species to confirm their phenology. Of the species active for seven or more months, most were eusocial [*e.g.* *Auglochloa aurífera*, *A. nigrocyanea* and *A. smaradigma* (Halictidae), *Frieseomelitta nigra*, *Melipona yucatanica*, *Nannotrigona perilampoides* and *Trigona fulviventris* (Apidae), some were parasocial (*e.g.*, *Ceratina eximia*, *C. nautlana*, *Euglossa viridissima* and *Eulaema polychroma*), and a few were solitary (*e.g.*, *Anthidiellum apicale* [Halictidae] and *Centris analis* [Apidae]).

When Hurricane Dean impacted the YP in August, its strong winds caused severe damage, including downed trees and branches, almost complete defoliation, almost total destruction of bush and herbaceous species, and removal of any floral resources present at the time [14]. As suggested in other studies [*e.g.*, 48], it may also have displaced some bee species, particularly those small in size, to other areas. However, the hurricane's effects were not catastrophic for the overall bee community, since two months later it had regained much of its lost activity. Although Hurricane Dean was a category 5 and had degraded to a category 3/4 over the study area, it differed from other category 5 hurricanes that have impacted the YP (*e.g.*, Gilbert and Wilma) in that it brought much less precipitation, and its force was exercised mainly as wind. It did cause almost complete defoliation, but within a month leaf cover had reached 70 to 80% of normal, indicating rapid vegetation recovery [14]. This may have allowed a commensurately rapid recovery in native bee activity, since species diversity had recovered (increased) notably within two months after the impact. The hurricane's impact on the native bee community in the study area was apparently temporary. Two possible reasons are that its winds may have had only moderate effects on the bee community, and/or Neotropical bee faunas have the capacity to recover rapidly from natural disturbances as long as the flowering plant community is resilient and adequate nesting sites and floral resources are available.

Implications for conservation

Disturbances from natural phenomena such as hurricanes could be much greater under circumstances different from those documented here; for instance, if strong winds are accompanied by copious rainfall, and/or if an area is already affected by anthropic activities. In tandem, these factors could seriously damage bee populations by altering their phenologic cycles and reducing the number of available nesting sites. This could be especially acute for eusocial species, which nest in cavities in trees, and solitary species, whose nesting sites can be destroyed by flooding, landslides, and soil churning. Further study and a more thorough understanding of pollinator species like native bees are vital because they play vital roles in maintaining vegetation communities and aiding in their recovery after disturbances.

Acknowledgements

VMR was financed by the project “Evaluación de la Biodiversidad de las Áreas Naturales Protegidas del estado de Yucatán, usando grupos indicadores, propuesta de nuevas áreas y estrategias de manejo y conservación” (SEMARNAT-2004-C01-180).

References

- [1] Michener, C. D. 2007. *The Bees of the World*. The Johns Hopkins University Press. Baltimore and London.
- [2] Ayala, R., Griswold, T. L. and Yanega, D. 1996. Apoidea (Hymenoptera). In: *Biodiversidad, taxonomía y biogeografía de artrópodos de México: Hacia una síntesis de su conocimiento*. Llorente-Bousquets, J., García-Aldrete, A. N. and González-Soriano, E. (Eds.). UNAM, CONABIO. México.
- [3] Heithaus, E. R. 1979. Community structure of Neotropical flower visiting bees and wasps: Diversity and phenology. *Ecology* 60(1): 190-202.
- [4] Ayala, R. 2004. Fauna de abejas silvestres (Hymenóptera: Apoidea) In: *Artrópodos de Chamela*. García, A. A. and Ayala, R. (Eds.). Instituto de Biología. UNAM. México.
- [5] Roubik, D. W., Villanueva, R., Cabrera, E. and Colly, U. F. 1991. Abejas nativas de la Reserva de la Biosfera de Sian Ka'an, Quintana Roo, México. In: *Biodiversidad Biológica en la Reserva de la Biosfera de Sian Ka'an, Quintana Roo, México*. Navarro, L. D. and Robinson, J. G. (Eds.). CIQRO. Chetumal, Quintana Roo.
- [6] Novelo, R. L., Delfín G. H., Ayala R. and Contreras, A. H. 2003. Estructura de las comunidades de abejas nativas de cuatro tipos de vegetación en el estado de Yucatán, México. *Folia Entomologica Mexicana* 42(2):177-190.
- [7] Reyes, N. E., Meléndez R. V., Ayala, R. and Delfín G. H. 2009. Bee faunas (Hymenoptera: Apoidea) of six natural protected areas in Yucatan, Mexico. *Entomological News*. 120 (5):530-544.
- [8] Meléndez, R. V., Magaña, R. S, Parra, T. V., Ayala, R. and Navarro, J. 2002. Diversity of native bee visitor of cucurbit crops (Cucurbitaceae) in Yucatán, México. *Journal of Insect Conservation*. 6(3): 135-147.
- [9] Meléndez R. V., Parra, T. V. Kevan, P. G., Ramírez, M. I., Zizumbo, V. D., Harries, H. and Fernandez, B. M. 2004. Mixed Mating Strategies and Pollination by Insects and Wind in Coconut Palm (*Cocos nucifera* L. (Arecaceae)). *Agriculture, Forestry and Entomology*. 6:155-163.
- [10] Meneses C. L., Meléndez R. V., Parra T. V. and Navarro A. J. 2010. Bee diversity in fragmented landscapes of the Mexican neotropic. *Journal of Insect Conservation*. 14(4): 323-334.
- [11] Flores, G. S. and Espejel, C. I. 1994. Tipos de vegetación de la Península de Yucatán. *Etnoflora Yucatanense*. UADY.
- [12] Durán, R. and Olmsted, I. 1999. Capítulo III. 3. Vegetación. *Atlas de procesos territoriales de Yucatán*. In: García de Fuentes A. and Córdoba-Ordoñez J. (Eds.). Universidad Autonoma de Yucatán. México. pp. 187-194.
- [13] Vidal, Z. R. 2005. *Las Regiones Climáticas de México*, Instituto de Geografía, Universidad Nacional Autonoma de México, México, Colección. Temas Selectos de Geografía de México (I.2.2).

- [14] Lugo, A. E. 2008. Visible and invisible effects of hurricanes on forest ecosystems: an international review. *Austral Ecology* 33:368-398.
- [15] Turton, S.M. and Siegenthaler, D. T. 2004. Immediate impacts of a severe tropical cyclone on the microclimate of a tropical rainforest canopy in northeast Australia. *Journal Tropical Ecology* 20:583-586.
- [16] Ostertag, R., Silver, W. L., and Lugo, A. E. 2005. Factors effecting mortality and resistance to damage following hurricanes in a rehabilitated subtropical moist forest. *Biotropica* 37:16-24.
- [17] Catterall, C. P., Mckenna, S., Kanowski, J. and Piper, S. D. 2008. Do cyclones and forest fragmentation have synergistic effects? A before-after study of rainforest structure at multiple sites. *Austral Ecology* 33:471-484.
- [18] Islebe, G. A., Torrescano, V. N., Valdez, H. M., Tuz, N. M. and Weissenberger, H. 2009. Efectos del impacto del huracán Dean en la vegetación del sureste de Quintana Roo, México. *Foresta Veracruzana* 11(1):1-6.
- [19] Schowater, T. D. 2012. Insect responses to major landscape-level disturbance. *Annual Review of Entomology* 57:1-20
- [20] Grimbacher, P. S. and Stork, N. E. 2009. How do beetle assemblages respond to cyclonic disturbance of a fragmented tropical rainforest landscape? *Oecologia* DOI 10.1007/s00442-009-1399-5
- [21] Cariveau, D. P. and Winfree, R. 2015. Causes of variation in wild bee responses to anthropogenic drivers. *Current Opinion in Insect Science* 10:104-109.
- [22] Roubik, D. W. 1992. *Ecology and natural history of tropical bees*. Cambridge University Press. NewYork, USA.
- [23] Michener, C. D. 1974. *The Social Behavior of the Bees. A Comparative Study*. The Bekknap Press of Harvard University Press, Cambridge, Massachusetts.
- [24] Orellana R, Balam M, Bañuelos I, García E, González, I. A, Herrera, F. and Vidal J. 1999. Evaluación climática. *Atlas de Procesos Territoriales de Yucatán*. In: García de Fuentes A. and Córdoba-Ordoñez J. (Eds.). Universidad Autónoma de Yucatán. México. p. 163-182
- [25] CONAGUA, 2008. Datos climatológicos mensuales en la estación metereológica de Nohalal, Municipio de Tekax, Yucatán. Comisión Nacional del Agua: Gerencia Regional Península de Yucatán. México.
- [26] Bautista, F., Batllori, E., Ortiz, M. A., Palacio, G. and Castillo, M. 2005. Integración del conocimiento actual sobre los paisajes geomorfológicos de la Península de Yucatán. 33-58 pp. In: *Caracterización de suelos en la Península de Yucatán, implicaciones agropecuarias, forestales y ambientales*. Universidad Autónoma de Campeche, Universidad Autónoma de Yucatán, Instituto Nacional de Ecología.
- [27] Hernández, U. A. 2007. Reseña del Huracán "Dean" del Océano Atlántico. CONAGUA-Servicio Meteorológico Nacional, México.
- [28] SAGARPA 2007. El paso del huracán "Dean" Informe al 30 de agosto de 2007. Secretaria de Agricultura, Ganadería, Desarrollo Rural, Pesca y Alimentación.
- [29] Delfín, G. H., Manrique, S. P., Meléndez, R. V. and Reyes, N. E. 2011. Insectos terrestres. In: Bautista, Z. F. Delfín, G. H. and J. L. Palacio Prieto (Eds.). *Técnicas de muestreo para manejadores de recursos naturales*. Centro de Investigaciones en Geografía Ambiental. UNAM. pp. 415-448.

- [30] Ayala, R. 1988. Abejas silvestres (Hymenoptera: Apoidea) de Chamela, Jalisco, México. *Folia Entomologica Mexicana* 77:395-493.
- [31] Mayr, E. and Ashlock, P. D. 1991. *Principles of Systematic Zoology*, McGraw-Hill, New York, U.S.A.
- [32] Colwell, R. K. and Coddington, J. A. 1994. Estimating terrestrial biodiversity through extrapolation. *Philosophical Transactions of the Royal Society of London Series B* 345:101-118.
- [33] Biodiversity Pro. 1997. <http://www.sams.ac.uk/dml/projects/benthic/bdpro>.
- [34] Species Diversity and Richness ver. SDR 3.03 (Pisces Conservation, 2004).
- [33] Solow, AR. 1993. A simple test for change in community structure. *Journal Animal Ecology* 62(1): 191-193.
- [35] Colwell, R. K. 2006. Statistical estimation of species richness and shared species from samples. Version 8.0. University of Connecticut, USA. Available at: <http://purl.oclc.org/estimates>
- [36] Statistica ver. 7 program (2006)
- [37] MVSP ver. 3.13 program (Kovach 2004)
- [38] Michener, C. D., McGinley, R. J. and Danforth, B. 1994. *The Bee Genera of North and Central America (Hymenoptera: Apoidea)*. Smithsonian Institution Press. Washington, USA.
- [39] Oertli, S., Müller, A. and Dorn, S. 2005. Ecological and seasonal patterns in the diversity of a species-rich bee assemblage (Hymenoptera: Apoidea: Apiformes). *European Journal Entomology* 102: 53-63.
- [40] Smith, P. A. and González, V. H. 2007. Diversidad de abejas (Hymenoptera: Apoidea) en estados sucesionales del bosque húmedo tropical. *Acta Biologica Colombiana* 12(1): 43-56.
- [41] Nates, P. G. and González, V. H. 2000. Las abejas silvestres de Colombia: Por qué y cómo conservarlas. *Acta Biológica Colombiana* 5(2):5-37.
- [42] Potts, S. G., B. Vulliamy, S. Roberts, C. O'Toole, A. Dafni, G. Ne'eman, and Willmer, P. 2005. Role of nesting resources in organising diverse bee communities in a Mediterranean landscape. *Ecological Entomology* 30:78–85.
- [43] Cane J. H. and Payne J. A. 1993. Regional, annual and seasonal-variation in pollinator guides intrinsic traits of bees (Hymenoptera, Apoidea) underlie their patterns of abundance at *Vaccinium ashei* (Ericaceae). *Annals of Entomological Society of America* 86:577-588.
- [44] Tylanakis, J. M., Klein, A. M., Lozada, T. and Tscharntke, T. 2006. Spatial scale of observation affects alfa, beta and gama diversity of cavity-nesting bees and wasps across a tropical land-use gradient. *Journal of Biogeography* 33: 1295-1304.
- [45] Flores, S. J. 1990. The flowering periods of leguminosae in the Yucatan Peninsula in relation to honey flows. *Journal of Apicultural Research* 29(2): 82-88.
- [46] Kevan, P. G. and Baker, H. G. 1983. Insects as flower visitors and pollinators. *Annual Review Entomology* 28:407-53.
- [47] Williams, N. M., Minckley, R. L. and Silveira, F. A. 2001. Variation in native bee faunas and its implications for detecting community changes. In: Conservation Ecology: Variation in Native Bee Faunas and Its Implications, <http://www.ecologyandsociety.org/vol5/iss1/art7/>
- [48] Genaro, J. A. 2004. Las abejas de la Isla de la Juventud, Cuba (Hymenoptera: Apoidea). *Boletín S. E .A.* 34: 177-179.

Appendix 1. Bee species in the Tropical Subdeciduous Forest of the Yucatan Peninsula, Mexico.

F	Native bee Species	Jn	Fe	Ma	Ap	May	Ju	Jl	Ag	Se	Oc	No	Dc	LS	Nb
Meg	<i>Anthidiellum (Loyolanthidium) apicale</i> (Cresson, 1878)	--	■	■	■	■	--	--	■	■	■	■	--	So	C
Hal	<i>Augochloropsis metallica</i> (Fabricius, 1793)	--	--	--	--	■	■	■	■	■	■	--	--	Ps	W
Hal	<i>Augochlora (Augochlora) nigrocyanea</i> Cockerell, 1897	--	■	■	--	■	■	--	■	■	--	■	--	Ps	W
Hal	<i>Augochlora (Augochlora) smaragdina</i> Friese, 1917	--	--	■	■	■	--	■	■	■	■	■	--	Ps	W
Hal	<i>Augochlora (Oxystoglossella) aurifera</i> Cockerell, 1897	--	■	■	■	■	■	■	■	■	■	--	--	Eu	G
Hal	<i>Augochlora</i> sp 2	--	■	■	--	■	■	■	■	■	■	■	■	Ps	W
Hal	<i>Augochlora</i> sp 4	--	--	--	--	--	--	■	--	■	■	■	■	Ps	W
Hal	<i>Augochlora</i> sp 5	--	--	--	--	--	■	--	--	■	--	--	■	Ps	W
Hal	<i>Augochlora</i> sp 6	--	--	--	--	--	--	--	--	--	--	■	--	Ps	W
Hal	<i>Augochlora</i> sp 7	--	--	--	--	--	--	■	--	--	--	--	--	Ps	W
Hal	<i>Augochlorella neglectula</i> (Cockerell, 1897)	--	--	--	--	■	--	■	--	--	--	■	--	Eu	G
Hal	<i>Augochloropsis</i> sp 1	■	■	■	--	--	--	■	■	■	■	■	■	Ps	G
Hal	<i>Augochloropsis</i> sp 2	■	--	--	--	--	■	■	--	■	■	■	■	Ps	G
Hal	<i>Augochloropsis</i> sp 3	■	--	■	--	--	--	--	■	■	■	■	■	Ps	G
Hal	<i>Caenaugochlora (Caenaugochlora) gemmella</i> (Cockerell, 1912)	--	--	--	--	■	--	--	--	--	--	■	--	Ps	G
Hal	<i>Caenaugochlora</i> sp 1	--	■	--	--	--	--	--	■	--	--	--	--	Ps	G
Hal	<i>Caenaugochlora</i> sp 2	--	--	--	--	--	--	--	--	--	--	■	--	Ps	G
Api	<i>Centris (Heterocentris) analis</i> (Fabricius, 1804)	--	--	--	■	--	■	--	■	■	■	■	■	So	C
Api	<i>Centris (Heterocentris) nitida</i> Smith, 1874	--	--	■	■	■	■	--	■	■	--	--	--	So	C
Api	<i>Centris (Heterocentris) trigonoides</i> Lepeletier, 1841	--	--	■	--	--	■	--	■	■	--	■	■	So	C
Api	<i>Centris (Trachina) eurypatana</i> Snelling, 1984	--	--	■	--	--	--	--	--	--	--	--	--	So	G
Api	<i>Centris (Xanthemis) lutea</i> Friese, 1899	--	--	--	--	--	--	--	--	■	--	--	--	So	C
Api	<i>Cephalotrigona zexmeniae</i> (Cockerell, 1912)	■	--	■	--	--	--	■	■	--	--	■	■	Eu	G
Api	<i>Ceratina (Calloceratina) eximia</i> Smith, 1862	■	■	--	■	--	--	--	■	■	■	■	■	Ps	W
Api	<i>Ceratina (Calloceratina) itzarum</i> Cockerell, 1931	--	--	--	--	--	--	■	■	■	--	■	--	Ps	W
Api	<i>Ceratina (Ceratinula) sp1</i>	--	--	--	--	--	■	--	--	--	--	--	--	Ps	W
Api	<i>Ceratina (Ceratinula) sp2</i>	--	--	--	--	--	■	--	--	--	--	--	--	Ps	W
Api	<i>Ceratina (Zadontomerus) cf. nautlana</i> Cockerell, 1897	--	■	--	■	--	■	■	■	■	■	■	■	Ps	W
Api	<i>Ceratina (Zadontomerus) sp1</i>	--	--	--	--	--	■	--	■	--	--	--	--	Ps	W
Api	<i>Ceratina</i> sp 7	--	--	--	--	--	--	--	■	--	--	--	--	Ps	W
Meg	<i>Coelioxys (Acrocoelioxys)</i>	■	--	--	--	--	--	--	--	--	--	--	--	Cl	--
Meg	<i>Coelioxys (Cyrtocoelioxys) sp 1</i>	--	--	--	--	--	--	--	--	■	--	--	--	Cl	--
Meg	<i>Coelioxys (Glyptocoelioxys) sp 2</i>	--	--	--	--	--	--	--	--	--	■	■	--	Cl	--
Meg	<i>Coelioxys vituperabilis</i> Holmberg, 1903	--	--	--	--	--	--	--	--	--	--	--	■	Cl	--
Coll	<i>Colletes arizonensis</i> Stephen, 1954	--	--	--	--	--	■	--	■	--	--	--	--	So	G

Coll	<i>Colletes rugicollis</i> Friese, 1900	--	--	--	■	--	--	■	■	■	--	--	--	So	G
Meg	<i>Epanthidium (Carloticola) cf boharti</i> Stange, 1983	--	--	--	■	--	--	--	--	--	--	--	--	So	C
Api	<i>Eufriesea mexicana</i> (Mocsáry, 1897)	--	--	--	--	--	--	--	--	■	--	--	--	So	C
Api	<i>Euglossa (Euglossa) viridissima</i> Friese, 1899	■	--	■	--	■	■	■	■	■	■	--	Ps	C	
Api	<i>Eulaema (Apeulaema) polychroma</i> (Mocsáry, 1899)	■	--	--	■	■	--	■	■	■	■	--	Ps	C	
Api	<i>Exaerete smaragdina</i> (Guérin-Ménéville, 1845)	--	--	■	■	■	--	■	--	■	--	--	Cl	--	
Api	<i>Exomalopsis (Exomalopsis) boharti</i> Timberlake, 1980	■	■	■	--	--	■	■	--	--	■	--	So	G	
Api	<i>Exomalopsis (Exomalopsis) mellipes</i> Cresson, 1878	--	■	■	--	--	■	--	--	■	■	--	So	G	
Api	<i>Frieseomelitta nigra</i> (Cresson, 1878)	■	■	■	--	--	■	■	■	■	■	■	Eu	C	
Api	<i>Gaesischia exul</i> Michener, LaBerge y Moure, 1955	--	--	■	■	--	--	--	--	--	--	--	So	G	
Hal	<i>Halictus (Odontalictus) ligatus</i> Say, 1837	--	--	--	--	■	--	■	--	--	--	--	Eu	G	
Meg	<i>Heriades (Neotrypetes) variolosus</i> (Cresson, 1872)	--	■	--	■	■	--	--	--	--	--	--	So	C	
Coll	<i>Hylaeus (Hylaeopsis) sp1</i>	--	--	--	--	--	■	--	--	--	--	--	So	C	
Coll	<i>Hylaeus sp 4</i>	--	--	--	--	--	--	--	--	--	■	--	So	C	
Meg	<i>Hypanthidium (Hypanthidium) yucatanicum</i> Cockerell, 1931	--	--	--	--	■	■	--	■	■	■	--	So	C	
Hal	<i>Lasioglossum (Dialictus) sp 1</i>	--	--	--	--	--	--	--	--	--	--	■	Ps	G	
Hal	<i>Lasioglossum (Dialictus) sp 2</i>	--	■	■	■	■	■	--	■	■	■	--	Ps	G	
Hal	<i>Lasioglossum (Dialictus) sp 3</i>	--	--	■	■	■	--	■	■	--	■	--	Ps	G	
Hal	<i>Lasioglossum (Dialictus) sp 4</i>	--	■	■	■	--	■	■	--	--	--	--	Ps	G	
Hal	<i>Lasioglossum (Dialictus) sp 5</i>	--	■	■	■	■	■	--	■	--	■	--	Ps	G	
Meg	<i>Megachile (Leptorachis) sp 1</i>	--	--	--	--	--	■	--	■	■	■	■	So	C	
Meg	<i>Megachile (Acentron) sp 1</i>	--	--	--	■	■	--	--	--	--	--	--	So	C	
Meg	<i>Megachile (Acentron) albitarsis</i> Cresson, 1872	■	--	--	■	■	--	--	--	--	--	■	So	C	
Meg	<i>Megachile (Acentron) sp 2</i>	--	--	--	■	--	--	--	--	■	--	--	So	C	
Meg	<i>Megachile (Austromegachile) sp 1</i>	--	--	--	--	--	--	--	--	--	--	■	So	C	
Meg	<i>Megachile (Austromegachile) sp 2</i>	■	■	--	--	--	--	--	--	--	--	--	So	C	
Meg	<i>Megachile (Chelostomoides) texensis</i> Mitchell, 1956	--	--	■	■	--	--	--	--	--	--	--	So	C	
Meg	<i>Megachile (Leptorachis) sp 1</i>	■	--	--	--	--	■	--	--	--	--	■	So	C	
Meg	<i>Megachile (Melanosarus) nigripennis</i> Spinola, 1841	--	--	--	--	--	■	--	■	--	--	--	So	C	
Meg	<i>Megachile (Neochelynia) chichimeca</i> Cresson, 1878	--	--	--	--	■	--	--	■	--	--	--	So	C	
Meg	<i>Megachile (Pseudocentron) azteca</i> Cresson, 1878	--	■	--	■	■	■	--	■	--	--	■	So	C	
Meg	<i>Megachile (Pseudocentron) inscita</i> Mitchell, 1930	--	--	■	■	--	--	--	■	--	--	--	So	C	
Meg	<i>Megachile (Pseudocentron) sp 4</i>	--	--	■	--	■	■	--	■	--	--	--	So	C	
Meg	<i>Megachile (Sayapis) zaptlana</i> Cresson, 1878	--	--	--	■	--	--	--	--	--	--	--	So	C	
Meg	<i>Megachile (Sayapis) frugalis</i> Cresson, 1872	--	--	■	--	--	--	--	■	--	--	--	So	C	
Meg	<i>Megachile (Sayapis) sp1</i>	--	--	--	■	--	--	--	--	--	--	--	So	C	
Meg	<i>Megachile sp 1</i>	--	--	--	■	--	--	--	--	--	--	--	So	C	
Meg	<i>Megachile sp 2</i>	--	--	--	--	--	--	--	--	--	■	--	So	C	
Meg	<i>Megachile sp 3</i>	--	--	--	--	■	--	--	--	--	--	--	So	C	

Api	<i>Melipona beecheii</i> Bennett, 1831	■	--	--	--	--	--	■	■	■	■	--	■	Eu	C
Api	<i>Melipona yucatanica</i> Camargo, Moure and Roubik, 1988	--	■	■	■	--	■	--	■	--	■	■	■	Eu	C
Api	<i>Melissodes tepaneca</i> Cresson, 1878	■	--	--	■	■	■	--	--	--	--	■	--	So	G
Api	<i>Melitoma marginella</i> (Cresson, 1872)	--	--	--	--	--	--	--	■	--	■	■	--	So	G
Api	<i>Nannotrigona perilampoides</i> (Cresson, 1878)	■	--	■	--	--	■	■	--	■	■	■	■	Eu	C
Api	<i>Osiris semiatratus</i> Shanks, 1987	--	--	--	■	--	--	--	--	--	■	--	--	Cl	--
Api	<i>Paratetrapedia (Lophopedia) apicalis</i> (Cresson, 1878)	--	■	--	--	--	--	--	--	--	--	--	--	So	G
Api	<i>Paratetrapedia (Paratetrapedia) calcarata</i> (Cresson, 1878)	■	■	--	■	■	■	--	--	■	■	■	■	So	G
Api	<i>Paratetrapedia (Paratetrapedia) moesta</i> (Cresson, 1878)	--	--	■	■	■	■	■	■	■	■	■	--	So	G
Api	<i>Paratetrapedia (Xanthopedia) swainsonae</i> (Cockerell, 1909)	--	--	--	■	■	--	--	--	--	--	--	--	So	G
Api	<i>Peponapis limitaris</i> (Cockerell, 1906)	--	--	--	--	--	--	--	--	--	--	--	■	So	G
Api	<i>Peponapis</i> sp 1	--	--	--	--	--	--	--	--	--	--	--	■	So	G
Api	<i>Peponapis utahensis</i> (Cockerell, 1905)	--	--	--	--	--	--	--	--	--	--	■	--	So	G
Api	<i>Plebeia frontalis</i> (Friese, 1911)	■	■	■	■	■	■	■	■	■	■	■	--	Eu	C
Api	<i>Plebeia pulchra</i> Ayala, 1999	--	■	■	--	■	■	■	■	--	--	--	--	Eu	C
Hal	<i>Pseudaugochloropsis graminea</i> (Fabricius, 1804)	--	--	--	■	■	--	■	■	■	■	■	--	Ps	G
Col	<i>Ptiloglossa mexicana</i> (Cresson, 1878)	--	--	--	--	■	--	--	--	--	--	--	--	So	G
Col	<i>Ptiloglossa eximia</i> (Smith, 1861)	--	--	■	■	--	■	■	■	--	--	■	--	So	G
Api	<i>Scaptotrigona pectoralis</i> (Dalla Torre, 1896)	■	--	--	--	--	--	--	--	--	■	■	■	Eu	C
Api	<i>Tetrapedia (Tetrapedia) maura</i> Cresson, 1878	--	--	--	■	■	--	--	--	--	--	--	--	So	C
Api	<i>Triepeolus cameroni</i> (Meade-Waldo, 1913)	--	--	--	■	■	--	--	--	--	--	--	--	Cl	--
Api	<i>Trigona fulviventrtris</i> Guérin-Méneville, 1845	■	■	■	■	■	■	■	■	■	■	■	■	Eu	C
Api	<i>Trigona fuscipennis</i> Friese, 1900	--	--	■	■	--	--	--	--	--	--	--	--	Eu	C
Api	<i>Trigonisca pipioli</i> Ayala, 1999	■	■	■	■	■	■	--	--	■	■	--	--	Eu	C
Api	<i>Xylocopa (Neoxylocopa) mexicanorum</i> Cockerell, 1912	--	--	■	--	■	■	--	■	--	--	--	--	Ps	W
Api	<i>Xylocopa (Neoxylocopa) nautlana</i> Cockerell, 1904	--	--	--	--	■	--	--	--	--	--	--	--	Ps	W
Api	<i>Xylocopa (Schoenherria) subvirescens</i> Cresson, 1879	--	--	--	--	--	■	--	■	--	--	--	--	Ps	W
Api	<i>Xylocopa (Schoenherria) muscaria</i> Fabricius, 1775	--	■	--	■	■	■	--	■	--	--	--	--	Ps	W

F= Family, Api= Apidae, Coll= Colletidae, Hal= Halictidae and Meg= Megachilidae. Months: Ja= January, Fe= February, Ma= March, Ap= April, My= May, Ju= June, Jl= July, Ag=August, Se= September, Oc= October, No=November and Dc= December. LS=Lifestyle: So=Solitary, Ps=Parasocial, Eu= Eusocial and Cl= Cleptoparasitic. Nb=Nesting behavior: C= Preexisting cavity, G=Ground and W=Wood. Specie: ■ = present, -- = absent