

## Letters

Authors: Anderson, Sean S., Peterson, Charles H., Cherr, Gary N.,

Hampton, Stephanie, and Blum, Michael

Source: BioScience, 62(12): 1010-1011

Published By: American Institute of Biological Sciences

URL: https://doi.org/10.1525/bio.2012.62.12.17

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at <a href="https://www.bioone.org/terms-of-use">www.bioone.org/terms-of-use</a>.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.




Figure 2. Horizontal surface location (in meters) of large oil droplets relative to the well head (the center circle). The numbers next to the plus sign (+) show the number of hours after midnight on 9 May. To calculate a surface location at a given time, the measured current profile from the Deepwater Discovery III was applied to a 10-millimeter oil droplet (the diameter of a large oil droplet that might form when dispersants are not used) that was assumed to rise at 20 centimeters per second. Although this model is quite simple, it should be adequate to give a relative measure of the importance of ocean currents on the rise of the oil.

surface. During the crude oil discharge, spotter planes observed a  $9 \times 1$  kilometer sheen at the surface. The sheen was estimated to have a thickness of between 0.3 and 5 microns, which represents 2.7–45 cubic meters of oil. Considering that 50 cubic meters of crude oil was released, a significant amount reached the surface at the low estimate. At the high estimate, 90% of the oil reached the surface!

Furthermore, the role of dispersants is not just to facilitate the formation of small, slowly rising oil droplets but also to hinder recoalescence into larger droplets (Young 1945, Ivanov et al. 1979, Vincent 1983) that rise more rapidly to the surface.

Evidence indicates that subsea dispersants reduced the volatile oil surfacing near the DWH well, which helped protect responders attempting to control the well. Clearly, some oil surfaced, but it was mostly away from the well and in smaller amounts, which helped reduce the amount of oil reaching sensitive shorelines.

> ROBIN RORICK TIM NEDWED GREG DEMARCO CORTIS COOPER

Robin Rorick (rorickr@api.org) is affiliated with the American Petroleum Institute, in Washington, DC. Tim Nedwed and Greg DeMarco are affiliated with the Exxon Mobil Corporation, in Irving, Texas. Cortis Cooper is affiliated with the Chevron Corporation, in San Ramon, California.

## References cited

Ivanov IB, Jain RK, Somasundaran P, Traykov TT. 1979. The role of surfactants on the coalescences of emulsion droplets. Pages 817–840 in Mittal KL, ed. Solution Chemistry of Surfactants, vol. 2. Plenum. Johansen Ø, Rye H, Cooper C. 2003. DeepSpill— Field study of a simulated oil and gas blowout in deep water. Spill Science Technology Bulletin 8: 433–443.

Peterson CH, et al. 2012. A tale of two spills: Novel science and policy implications of an emerging new oil spill model. BioScience 62: 461–469.

Vincent B. 1983. Emulsions and foams. Pages 175–196 in Tadros TF, ed. Surfactants. Academic Press.

Young CBF, Coons KW. 1945. Emulsions. Pages 159–173 in Young CBF, Coons KW, eds. Surface Active Agents: Theoretical Aspects and Applications. Chemical Publishing Co.

doi:10.1525/bio.2012.62.12.16

## Casual Observations on DWH Dispersant Effects Expose the Lack of Rigorous Science: Response to Rorick and Colleagues

Robin Rorick, of the American Petroleum Institute (API), and his colleagues question our doubt that subsurface dispersant application was required to prevent Deepwater Horizon (DWH) oil from reaching the sea surface. The photographs provided as support for the need for and efficacy of subsurface dispersant do not allow scientifically rigorous conclusions in the absence of quantitative measurements of subsurface processes and dynamics of materials transport. Despite our close connectivity to industry, government, and academic research, we are unaware of any compelling, peer-reviewed data documenting the efficacy, necessity, and consequences of subsurface dispersant application during the DWH spill. Rorick and his colleagues' assertions reinforce long-standing concerns that oil spill responses are not based on sufficient science. We expect a higher level of commitment to excellence from an industry capable of so much more.

We questioned the unsupported assumption that the vast majority of hydrocarbon retention at depth is attributable solely to subsurface dispersant use. Rorick and his colleagues' argument regarding Johansen and colleagues (2003) contradicts our understanding of that experiment. In a review of the study, Adams and Socolofsky (2005) noted that, in the absence of dispersants and under even less favorable conditions (shallower, colder, less turbulent water) for generating natural

dispersion than those of the DWH blowout, "most oil was not recovered at the surface, suggesting it was in the form of widely dispersed fine droplets" (p. 1). Even assuming that Rorick and his colleagues are correct, their own calculations suggest as little as 5% of the oil may have reached the surface in that study. By emphasizing the dubious high-end estimate of 90%, they reinforce our argument that unequivocal data are not available and that rigorous, process-oriented study is necessary.

Quantifying the behavior, transport, fate, and effects of hydrocarbons from deepwater blowouts, with and without dispersants at various application rates, is fundamental to the public interest. Because subsurface dispersant application is increasingly touted as a necessary new response method, rigorous scientific answers to questions not only of efficacy but also of the impact of subsurface retention of highly dispersed oil-dispersant droplets on particle feeders and an array of other organisms throughout the water

column are urgently needed. Such answers do not come from a few photographs. Rorick and his colleagues' restricted focus solely on preventing oil from reaching the sea surface reflects a failure to acknowledge the need for a new oil spill model that includes an assessment of risk below the surface following deepwater blowouts. We welcome formal, process-oriented data and urge API to pursue such studies.

> SEAN S. ANDERSON CHARLES H. PETERSON GARY N. CHERR STEPHANIE HAMPTON MICHAEL BLUM

Sean S. Anderson (sean.anderson@ csuci.edu) is an associate professor in the Environmental Science and Resource Management Program and director of the Pacific Institute for Restoration Ecology at California State University Channel Islands, in Camarillo, California. Charles H. Peterson (cpeters@email.unc.edu) is a distinguished professor at the Institute of Marine Sciences, at the University of North Carolina at Chapel Hill,

in Morehead City. Gary N. Cherr is a professor in the Departments of Environmental Toxicology and Nutrition and director of the University of California, Davis, Bodega Marine Laboratory, in Bodega Bay, California. Stephanie Hampton is the deputy director of the National Center for Ecological Analysis and Synthesis, in Santa Barbara, California. Michael Blum is an assistant professor in the Ecology and Evolutionary Biology Department at Tulane University, in New Orleans, Louisiana.

## References cited

Adams EE, Socolofsky SA. 2005. Review of Deep Oil Spill Modeling Activity Supported by the DeepSpill JIP and Offshore Operators Committee. US Bureau of Ocean Energy Management, Regulation and Enforcement. (1 October 2012; www.boemre.gov/tarprojects/ 377/Adams%20Review%204.pdf)

Johansen Ø, Rye H, Cooper C. 2003. DeepSpill-Field study of a simulated oil and gas blowout in deep water. Spill Science Technology Bulletin 8: 433-443.

doi:10.1525/bio.2012.62.12.17