

DIET, PLASMA CAROTENOIDS, AND SEXUAL COLORATION IN THE ZEBRA FINCH (TAENIOPYGIA GUTTATA)

Authors: McGraw, Kevin J., Gregory, Alexander J., Parker, Robert S.,

and Adkins-Regan, Elizabeth

Source: The Auk, 120(4): 1212

Published By: American Ornithological Society

URL: https://doi.org/10.1642/0004-

8038(2003)120[1212:DPCASC]2.0.CO;2

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Erratum

Kevin J. McGraw, 1 Alexander J. Gregory, 1 Robert S. Parker, 2 and Elizabeth Adkins-Regan 1,3

DIET, PLASMA CAROTENOIDS, AND SEXUAL COLORATION IN THE ZEBRA FINCH (TAENIOPYGIA GUTTATA)

¹Department of Neurobiology and Behavior, Cornell University, Ithaca, New York 14853, USA; ²Division of Nutritional Sciences, Cornell University, Ithaca, New York 14853, USA; and ³Department of Psychology, Cornell University, Ithaca, New York 14853, USA

Due to an omission during final proofing, Table 3 in McGraw et al. (Auk 120:400–410) was inadvertently omitted. It is reproduced here in full.

TABLE 3. Spearman-rank correlations between plasma carotenoid concentrations and beak coloration in captive male and female Zebra Finches (*Taeniopygia guttata*). In all comparisons n=12, and sequential Bonferroni adjustments were applied to correct for multiple comparisons (carotenoid types) within each sex.

Sex	Plasma carotenoid	$r_{\rm s}$	P
Male	Lutein	-0.69	0.01
	Zeaxanthin	-0.54	0.07
	Anhydrolutein	-0.64	0.02
	β-cryptoxanthin	-0.42	0.16
Female	Lutein	-0.71	0.01
	Zeaxanthin	-0.74	0.008
	Anhydrolutein	-0.74	0.008
	β-cryptoxanthin	-0.59	0.03