BioOne COMPLETE

Trichospilus diatraeae (Hymenoptera: Eulophidae): A Potential Biological Control Agent of Lepidopteran Pests of Oil Palm in the Brazilian Amazon

Authors: Ribeiro, Rafael C., Lemos, Walkymário De P., Castro, Ancidériton A. De, Poderoso, Júlio C. M., Serrão, José E., et al.

Source: Florida Entomologist, 96(2) : 676-678

Published By: Florida Entomological Society

URL: https://doi.org/10.1653/024.096.0245

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Downloaded From: https://complete.bioone.org/journals/Florida-Entomologist on 28 Mar 2024
Terms of Use: https://complete.bioone.org/terms-of-use

TRICHOSPILUS DIATRAEAE (HYMENOPTERA: EULOPHIDAE): A POTENTIAL BIOLOGICAL CONTROL AGENT OF LEPIDOPTERAN PESTS OF OIL PALM IN THE BRAZILIAN AMAZON

Rafael C. Ribeiro[1,*], Walkymário De P. Lemos[2], Ancidériton A. De Castro[3], Júlio C. M. Poderoso[3], José E. Serrão[4] and José C. Zanuncio[3]

[1]Departamento de Fitotecnia, Universidade Federal de Viçosa, Viçosa, Minas Gerais, Brazil

[2]Laboratório de Entomologia, Embrapa Amazônia Oriental, CPATU, Belém, Pará, Brazil

[3]Departamento de Entomologia, Universidade Federal de Viçosa, Viçosa, Minas Gerais, Brazil

[4]Departamento de Biologia Animal, Universidade Federal de Viçosa, Viçosa, Minas Gerais, Brazil

*Corresponding author; E-mail: rafael.c.ribeiro@ufv.br

Oil palm (*Elaeis guineensis*) monoculture may favor insect pests and disease vectors (Gitau et al. 2009). The defoliating caterpillars, *Opsiphanes invirae* Hübner and *Brassolis sophorae* L. (Lepidoptera: Nymphalidae), and the crown-leaf borer, *Eupalamides cyparissias* (Fabricius) (Lepidoptera: Castniidae), affect the development and productivity of this crop in northern Brazil (Vasquez et al. 2008; Ribeiro et al. 2010). Studies involving control methods of these pests are needed.

Trichospilus diatraeae Margabandhu & Cherian (Hymenoptera: Eulophidae) is a pupal parasitoid of many lepidopteran families, such as Arctiidae (Bennett et al. 1987; Paron & Berti-Filho 2000), Crambidae (Bennett et al. 1987), Oecophoridae (Oliveira et al. 2001), Pieridae (Torres-Bauza 1994), Geometridae (Pereira et al. 2008; Zaché et al. 2010; Pastori et al. 2012), Saturniidae (Pastori et al. 2012), Lymantriidae (Zaché et al. 2011a), Riodinidae (Zaché et al. 2011b), Noctuidae and Nymphalidae (Bennett et al. 1987; Zaché et al. 2012).

Female parasitoids must overcome the host's defensive behavior and immune response (Andrade et al. 2010). These factors can affect the biological characteristics and the host choice of parasitoids (Gross 1993; Völkl & Stadler 1996; Farias & Hopper 1999; Walker & Hoy 2003). The objective of this study was to evaluate the potential parasitism of lepidopteran pests of oil palm by *T. diatraeae*.

Opsiphanes invirae, *B. sophorae* and *E. cyparissias* pupae were obtained from oil palms in the district of Tailandia, Pará State, Brazil (S 24° 2.04' W 48° 08.02'). *Trichospilus diatraeae* was reared on *Tenebrio molitor* L. (Coleoptera: Tenebrionidae) pupae at the Federal University of Viçosa, Viçosa, Minas Gerais State, Brazil, at 25 ± 2 °C, 70 ± 10% RH and 12:12 h L:D (Favero 2009). Originally, *T. diatraeae* was obtained from *Thyrinteina arnobia* (Stoll) (Lepidoptera: Geometridae) pupae collected on *Eucalyptus cloeziana* F. Muell. (Myrtales: Myrtaceae) plants (Pereira et al. 2008).

Groups of fifty 2-day old *T. diatraeae* females were placed inside test tubes (14.5 cm H × 2.0 cm diam) together with one pupa of one of the following hosts: *O. invirae*, *B. sophorae* or *E. cyparissias,* for 48 h. The tubes were closed with cotton and parasitoids females fed with a drop of pure honey applied with a surgical needle on the wall of the test tubes (Pereira et al. 2009).

Trichospilus diatraeae parasitized *E. cyparissias* (Fig. 1A), *B. sophorae* (Fig. 1B) and *O. invirae* pupae (Fig. 1C), with parasitism rates of 90%, 100% and 90%, respectively. Releases of this parasitoid may provide an alternative biological control in oil palm crops, where several species of Lepidoptera pests can occur simultaneously (Ribeiro et al. 2010). However, further studies are needed to assess the mobility and dispersal of *T. diatraeae* in the field (Zappalà et al. 2012) and determine whether mass release of this parasitoid could interfere with other natural enemies (Charles 2012).

Trichospilus diatraeae developed in pupae of *O. invirae* and *B. sophorae.* An average of 447.83 ± 51.52 and 669.00 ± 89.62 parasitoid adults/pupa were obtained from these hosts, respectively. These results show the potential of this parasitoid to control defoliators and its possible establishment in the field through inoculative releases (Eilenberg et al. 2001). However, *T. diatraeae* adults did not emerge from *E. cyparissias* pupae probably because of host defense mechanisms (Gross 1993; Pennacchio & Strand 2006). The number of *T. diatraeae* females (50) per *E. cyparissias* pupae may have been insufficient to generate the amount of toxins and immature individuals required to overcome the host defense mechanism (Zaki et al. 1994; Andrade et al. 2010). Encapsulated parasitoids were found inside dissected pupae of *E. cyparissias*.

Downloaded From: https://complete.bioone.org/journals/Florida-Entomologist on 28 Mar 2024
Terms of Use: https://complete.bioone.org/terms-of-use




Fig. 1. *Eupalamides cyparissias* (Lepidoptera: Castniidae) (A), *Brassolis sophorae* (B) and *Opsiphanes invirae* (C) (Lepidoptera: Nymphalidae) pupae parasitized by *Trichospilus diatraeae* (Hymenoptera: Eulophidae).

Trichospilus diatraeae can be reared on pupae of the oil palm defoliator caterpillars, *O. invirae* and *B. sophorae,* indicating the potential of this parasitoid for the control of these pests.

Summary

The expansion of oil palm (*Elaeis guineensis*) monoculture may favor insect pests such as the caterpillars of *Opsiphanes invirae* Hübner, *Brassolis sophorae* L. (Lepidoptera: Nymphalidae) and *Eupalamides cyparissias* (Fabricius) (Lepidoptera: Castniidae). The objective of this study was to evaluate the potential of *Trichospilus diatraeae* Cherian & Margabandhu (Hymenoptera: Eulophidae) to parasitize pupae of lepidopteran pests of oil palm. Groups of fifty females of *T. diatraeae* were maintained inside test tubes with one pupa of one of the following hosts: *O. invirae*, *B. sophorae* or *E. cyparissias* for 48 h. *Trichospilus diatraeae* parasitized the 3 Lepidopteran species, but reproduced only on *O. invirae* and *B. sophorae*. These results suggest that this parasitoid, of polyphagous habit, may represent an alternative for the control of lepidopteran pests of oil palm in the Brazilian Amazon.

Resumo

A expansão da monocultura da palma do óleo (*Elaeis guineensis*) pode favorecer pragas como as lagartas *Opsiphanes invirae* Hübner, *Brassolis sophorae* L. (Lepidoptera: Nymphalidae) e *Eupalamides cyparissias* (Fabricius) (Lepidoptera: Castniidae). O objetivo foi avaliar o potencial de *Trichospilus diatraeae* Cherian & Margabandhu (Hymenoptera: Eulophidae) como parasitóide de pupas de lepidópteras-praga da palma do óleo. Cinquenta fêmeas de *T. diatraeae* foram mantidas um tubo de ensaio por 48 horas com uma pupa dos seguintes hospedeiros: *O. invirae*, *B. sophorae* ou *E. cyparissias*, com, até, dois dias de

Downloaded From: https://complete.bioone.org/journals/Florida-Entomologist on 28 Mar 2024
Terms of Use: https://complete.bioone.org/terms-of-use

idade. *Trichospilus diatraeae* realizou o parasitismo nas três espécies de Lepidoptera, mas obteve potencial reprodutivo em pupas de *O. invirae* e *B. sophorae*. Isto sugere que este parasitoide, de hábito polífago, pode representar uma alternativa para o controle de pragas de lepidópteros da palma de óleo na Amazônia brasileira.

Acknowledgments

We thank "Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq)", "Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES)", "Fundação de Amparo à pesquisa do Estado de Minas Gerais (FAPEMIG)" and "Grupo Agropalma" for financial support.

References Cited

Andrade, G. S., Serrão, J. E., Zanuncio, J. C., Zanuncio, T. V., Leite, G. L. D., and Polanczyk, R. A. 2010. Immunity of an alternative host can be overcome by higher densities of its parasitoids *Palmistichus elaeisis* and *Trichospilus diatraeae*. PLoS One 5(10): 13231.

Bennett, F. D., Glenn, H., Yaseen, M., and Baranowski, R. M. 1987. Records of *Trichospilus diatraeae*, an Asian parasite (Hymenoptera: Eulophidae) from the Caribbean and Florida. Florida Entomol. 70: 184-186.

Charles, J. G. 2012. Assessing the non-target impacts of classical biological control agents: is host-testing always necessary? BioControl 57: 619-626.

Eilenberg, J., Hajek, A., and Lomer, C. 2001. Suggestions for unifying the terminology in biological control. BioControl 46: 387-400.

Farias, A. M. I., and Hopper, K. R. 1999. Oviposition behavior of *Aphelinus asychis* (Hymenoptera: Aphelinidae) and *Aphidius matricariae* (Hymenoptera: Aphidiidae) and defense behavior of their host *Diuraphis noxia* (Homoptera: Aphididae). Environ. Entomol. 28: 858-862.

Fávero, K. 2009. Biologia e técnicas de criação de *Trichospilus diatraeae* (Hymenoptera: Eulophidae) em pupas de *Tenebrio molitor* (Coleoptera: Tenebrionidae) e *Diatraea saccharalis* (Lepidoptera: Crambidae). Dourados, MS, (Mestrado em Entomologia e Conservação da Biodiversidade) - Universidade Federal da Grande Dourados, 77 p.

Gitau, C. W., Gurr, G. M., Dewhurst, C., Fletcher, M. J., and Mitchell, A. 2009. Insect pests and insect-vectored diseases of palms. J. Australian Entomol. 48: 328-342.

Gross, P. 1993. Insect behavioral and morphological defenses against parasitoids. Annu. Rev. Entomol. 38: 251-27.

Oliveira, M. A. S., Junqueira, N. T. V., Icuma, I. M., Alves, R. T., Oliveira, J. N. S., and Andrade, G. A. 2001. Incidência de danos da broca do fruto da graviola no Distrito Federal. Comun. Técnol. 51: 01-06.

Pastori, P. L., Pereira, F. F., Andrade, G. S., Silva, R. O., Zanuncio, J. C., and Pereira, A. Í. A. 2012. Reproduction of *Trichospilus diatraeae* (Hymenoptera: Eulophidae) in pupae of two lepidopterans defoliators of eucalypt. Rev. Colombiana Entomol. 38: 91-93.

Paron, M. R., and Berti Filho, E. 2000. Capacidade reprodutiva de *Trichospilus diatraeae* (Hymenoptera: Eulophidae) em pupas de diferentes hospedeiros (Lepidoptera). Sci. Agríc. 57: 355-358.

Pennacchio, F., and Strand, M. R. 2006. Evolution of developmental strategies in parasitic Hymenoptera. Annu. Rev. Entomol. 51: 233-258.

Pereira, F. F., Zanuncio, J. C., Tavares, M. T., Pastori, P. L., Jacques, G. C., and Vilela, E. F. 2008. New record of *Trichospilus diatraeae* as a parasitoid of the eucalypt defoliator *Thyrinteina arnobia* in Brazil. Phytoparasitica 36: 304-306.

Pereira, F. F., Zanuncio, J. C., Serrão, J. E., Oliveira, H. N., Fávero, K., and Grance, E. L. V. 2009. Progênie de *Palmistichus elaeisis* Delvare & LaSalle (Hymenoptera: Eulophidae) parasitando pupas de *Bombyx mori* L. (Lepidoptera: Bombycidae) de diferentes idades. Neotrop. Entomol. 38: 660-664.

Ribeiro, R. C., Lemos, W. P., Bernardino, A. S., Buecke, J., and Müller, A. A. 2010. Primeira ocorrência de *Alcaeorrhynchus grandis* (Dallas) (Hemiptera: Pentatomidae) predando lagartas desfolhadoras do dendezeiro no Estado do Pará. Neotrop. Entomol. 39: 131-132.

Torres-Bauza, J. A. 1994. Hymenopterous parasitoids of *Dismorfia spio* (Pieridae: Dismorphiinae). J. Lepidopt. Soc. 48: 266.

Vasquez, J., Delgado, C., Couturier, G., Mejia, K., Freitas, L., and Castillo, D. 2008. Pest insects of the palm tree *Mauritia flexuosa* L.f., dwarf form, in Peruvian Amazonia. Fruits 63: 227-238.

Völkl, W., and Stadler, B. 1996. Colony orientation and successful defense behavior in the conifer aphid, *Schizolachnus pineti*. Entomol. Exp. Appl. 78: 197-200.

Walker, A. M., and Hoy, M. A. 2003. Responses of *Lipoplexis oregmae* (Hymenoptera: Aphidiidae) to different instars of *Toxoptera citricida* (Homoptera: Aphididae). J. Econ. Entomol. 96: 1685-1692.

Zaki, F. N., Elsaadany, G., Gomaa, A., and Saleh, M. 1994. Some biological factors affecting the production of the larval parasitoid *Bracon brevicornis* Wesm. (Hymenoptera: Braconidae). J. Appl. Entomol. 118: 413-418.

Zaché, B., Wilcken, C. F., Da Costa, R. R., and Soliman, E. P. 2010. *Trichospilus diatraeae* Cherian & Margabandhu, 1942 (Hymenoptera: Eulophidae), a new parasitoid of *Melanolophia consimilaria* (Lepidoptera: Geometridae). Phytoparasitica 38: 355-357.

Zaché, B., Wilcken, C. F., Zaché, R. R. C., and Souza, N. M. 2012. New occurrence of *Trichospilus diatraeae* (Hymenoptera: Eulophidae) as a parasitoid of *Spodoptera cosmioides* Walker, 1858 (Lepidoptera: Noctuidae) in Brazil. Biota Neotrop. 12: 1-4.

Zaché, B., Zaché, R. R. C., Soliman, E. P., and Wilcken, C. F. 2011a. Evaluation of *Trichospilus diatraeae* (Hymenoptera: Eulophidae) as parasitoid of the eucalyptus defoliator *Euselasia eucerus* (Lepidoptera: Riodinidae). Intl. J. Trop. Insect Sci. 20: 1-5.

Zaché, B., Zaché, R. R. C., Souza, N. M., Dias, T. K. R., and Wilcken, C. F. 2011b. New record of *Trichospilus diatraeae* margabandhu & cherian, 1942 (Hymenoptera: Eulophidae) parasitizing *Sarsina violascens* (Herrich-Schaeffer, 1856) (Lepidoptera: Lymantriidae) in Brazil. J. Plant. Protect. Res. 51: 420-422.

Zappalà, L., Campolo, O., Grande, S. B., Saraceno, F., Biondi, A., Siscaro, G., and Palmeri, V. 2012. Dispersal of *Aphytis melinus* (Hymenoptera: Aphelinidae) after augmentative releases in citrus orchards. European J. Entomol. 109: 561-568.

Downloaded From: https://complete.bioone.org/journals/Florida-Entomologist on 28 Mar 2024
Terms of Use: https://complete.bioone.org/terms-of-use