

Microsatellite Markers for *Aleurocanthus spiniferus* (Hemiptera: Aleyrodidae) and Their Potential use in Whiteflies

Authors: Tang, Xiao-Tian, Tao, Huan-Huan, Wang, Ji-Rui, and Du, Yu-Zhou

Source: Florida Entomologist, 97(3) : 1035-1040

Published By: Florida Entomological Society

URL: <https://doi.org/10.1653/024.097.0305>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

MICROSATELLITE MARKERS FOR *ALEUROCANTHUS SPINIFERUS* (HEMIPTERA: ALEYRODIDAE) AND THEIR POTENTIAL USE IN WHITEFLIES

XIAO-TIAN TANG, HUAN-HUAN TAO, JI-RUI WANG AND YU-ZHOU DU*

School of Horticulture and Plant Protection and Institute of Applied Entomology,
Yangzhou University, Yangzhou 225009, China

*Corresponding author; E-mail: yzdu@yzu.edu.cn

ABSTRACT

The citrus (or orange) spiny whitefly, *Aleurocanthus spiniferus* (Quaintance) (Hemiptera: Aleyrodidae), is an important pest of various economic crops such as citrus and tea, causing severe economic losses. However, the population genetics of *A. spiniferus* is poorly understood, both in China and in other countries. To improve our knowledge of the population structure and genetic variation of this species, 10 polymorphic microsatellite loci for *A. spiniferus* were developed and characterized using FIASCO (Fast Isolation by AFLP of Sequences Containing Repeats). Our results showed that the polymorphic information contents (PIC) of all 10 loci were greater than 0.5, showing a high degree of polymorphism. The number of alleles ranged from 12 to 27 across 60 individuals from 6 populations. In addition, the expected heterozygosity (H_e) and the observed heterozygosity (H_o) ranged from 0.851 to 0.958 and from 0.621 to 0.850, respectively. Interestingly, all loci deviated from the Hardy-Weinberg equilibrium, although - according to our study - they were not affected by the species' specific reproductive strategy, and thus must be related to other unknown factors. Furthermore, linkage disequilibrium analysis revealed that C19-2 and D14, and also I20 and F12 showed linkage disequilibrium. Cross-species amplification was also tested in 5 closely related whitefly species (*Aleurodicus disperses* Russell, *Pealius mori* (Takahashi), *Aleuroclava aucubae* (Kuwana), *Bemisia tabaci* (Gennadius) MEAM1 and *B. tabaci* MED) in this study. Nine pairs of primers were successfully amplified at different levels except for F12. In particular, A21-2, D14, F13-2, and F42-2, coupled with I20 were successfully amplified across all the above species. Consequently, the 10 loci identified here can be used to study the population genetic structure of *A. spiniferus* and other closely related whitefly species.

Key Words: *Aleurocanthus spiniferus*, disequilibrium population genetic structure, genetic variation, microsatellite, cross-species amplification

RESUMEN

La mosca blanca espinosa de los cítricos (o naranja), *Aleurocanthus spiniferus* (Quaintance) (Hemiptera: Aleyrodidae), es una plaga importante de diversos cultivos económicos como los cítricos y el té, que causa graves pérdidas económicas. Sin embargo, la genética de las poblaciones de *A. spiniferus* es poco conocida, tanto en China como en otros países. Para mejorar nuestro conocimiento de la estructura de la población y la variación genética de esta especie, se desarrollaron y caracterizaron 10 loci microsatélites polimórficos para *A. spiniferus* utilizando FIASCO (Aislamiento Rápido por AFLP de Secuencias que Tienen Repeticiones, ARASTR). Nuestros resultados mostraron que el contenido de información polimórfica (CIP) de todos los 10 loci fueron mayor de 0.5, mostrando un alto grado de polimorfismo. El número de alelos varió de 12 a 27 a través de 60 individuos de 6 poblaciones. Además, la heterocigosidad esperada (H_e) y la heterocigosidad observada (H_o) oscilaron 0.851 a 0.958 y 0.621 a 0.850, respectivamente. Curiosamente, todos los loci se desviaron del equilibrio de Hardy-Weinberg, aunque de acuerdo a nuestro estudio ellos no fueron afectados por la estrategia reproductiva específica de la especie, y por lo tanto deben estar relacionados con otros factores desconocidos. Además, el análisis de disequilibrio de ligamiento reveló que C19-2 y D14, y también I20 y F12 mostraron disequilibrio de ligamiento. La amplificación de cruz-especie también fue probada en 5 especies de mosca blanca estrechamente relacionadas (*Aleurodicus dispersus* Russell, *Pealius mori* (Takahashi), *Aleuroclava aucubae* (Kuwana), *Bemisia tabaci* (Gennadius) MEAM1 y *B. tabaci* MED) en este estudio. Se amplificaron con éxito nueve pares de cebadores a diferentes niveles con la excepción de F12. En particular, se amplificaron con éxito A21-2, D14, F13-2, y F42-2 junto con I20 a través de todas las especies anteriores. En consecuencia, los 10 loci identificados aquí pueden ser utilizados para estudiar la estructura genética de la población de *A. spiniferus* y otras especies de mosca blanca estrechamente relacionadas.

Palabras Clave: *Aleurocanthus spiniferus*, disequilibrio estructura genética de la población, variación genética, microsatélites, amplificación entre especies

The citrus spiny whitefly *Aleurocanthus spiniferus* (Quaintance) (Hemiptera: Aleyrodidae), is a pest of citrus (*Citrus* spp.; Rutaceae) and tea (*Camellia sinensis* (L.) Kuntze; Ericales; Theaceae), and several serious outbreaks of a tea-infesting whitefly in China and Japan have been attributed to this species over the last 20 yr (Kanmiya et al. 2011). The citrus spiny whitefly affects host plants by removal of plant sap, but whiteflies also cause indirect damage by producing honeydew, which promotes sooty mold. In addition to citrus and tea, this whitefly infests many other host plants, as noted in Evans (2008). Furthermore, *A. spiniferus* is an EPPO (European and Mediterranean Plant Protection Organization) quarantine species that has recently been moved from the A1 to the A2 list (Anonymous 2011). It is worth mentioning that commonly *A. spiniferus* reproduces sexually and infrequently by arrhenotoky (Huang et al. 1999), which may influence the Hardy-Weinberg equilibrium (see Discussion).

Because *A. spiniferus* is a well-known insect pest, some aspects of its biology, behavior, ecology, and management have been thoroughly investigated (Van den Berg et al. 2000; Han and Cui. 2002; Guo et al. 2006; Muniappan et al. 2006; Peng et al. 2010; Niu et al. 2014). In contrast, little is known about the genetic diversity and population structure of *A. spiniferus*, and no microsatellite markers have been developed for this species. Fu & Han (2007) preliminarily analyzed the genetic diversity of *A. spiniferus* from 7 populations in eastern China using random amplified polymorphic DNA (RAPD). Kanmiya et al. (2011) proposed new specific status for tea-infesting populations of the nominal citrus spiny whitefly *A. spiniferus* using molecular marker (mtCOI gene), morphological and acoustic analysis to distinguish it from *A. spiniferus* that constitutes the citrus-infesting population. However, genetic diversity for *A. spiniferus* has not been comprehensively investigated, and it is poorly understood. Such gaps in our knowledge highlight the need for the development of highly polymorphic and informative molecular markers in order to accurately assess and monitor genetic variation in and population structure of *A. spiniferus* populations.

Microsatellite DNA, also known as simple sequence repeats (SSRs), is a technique that has been widely used in many areas of research (Carleton et al. 2001; Peterlunger et al. 2003). In the field of entomology, it has been used for gene mapping, population genetic structure analysis, identification of genetic relationships, and genetic map constructions (Gorman 1997; Zenger et al. 2005). In this study, we developed 10 high polymorphic microsatellite markers for *A. spiniferus*, and cross-species amplifications were also conducted of the 5 closely related whitefly species (*Aleurodicus disperses* Russell, *Pealius mori* Takahashi, *Aleuroclava aucubae* Kuwana, *Bemisia tabaci* (Genna-

dus) MEAM1 and *B. tabaci* MED). These markers are expected to be used widely in studies on the genetic diversity and population genetic structure of whitefly species in their native and invasive ranges and in tracing their global invasion history.

MATERIALS AND METHODS

Aleurocanthus spiniferus samples were collected from tea plants in 6 provinces: Shandong (N 34.73° E 117.29°), Jiangsu (N 32.41° E 119.42°), Zhejiang (N 30.19° E 120.09°), Hunan (N 28.20° E 113.09°), Guizhou (N 27.66° E 107.60°), and Yunnan (N 25.13° E 102.75°) in China. At first, we did not identify the gender of the 60 *A. spiniferus* individuals, until we noticed that all 10 study loci deviated from Hardy-Weinberg equilibrium. To ensure the accuracy of our results, another 48 female individuals from Jiangsu (N 32.41° E 119.42°) and Sichuan (N 30.57° E 104.07°) provinces were tested later. Gender was determined as per Dubey & Ko (2012). All specimens (including those from the 5 closely related whitefly species) were preserved in 100% ethanol and stored at -20 °C until DNA extractions were performed.

Genomic DNA was extracted using the salting-out method as described by Teng et al. (2009). The enriched library was constructed by FIASC (Fast Isolation by AFLP of Sequences Containing Repeats) protocol according to Zane et al. (2002) with slight modifications in terms of PCR amplification conditions. The genomic DNA was first digested with the restriction enzyme *Mse*I (BioLabs, Beijing, China) and approximately 250 ng of DNA was ligated to 1 µg *Mse*I adaptor (5'-TACTCAGGACTCAT-3'/5'-GACGATGAGTCCTGAG - 3'). The digestion-ligation products was diluted (1:10) and amplified with adaptor-specific primers (5'-GATGAGTCCT-GAGTAAN-3', *Mse*I-N) in 20 µL reactions containing 1 × PCR buffer (10 mM Tris-HCl, pH 9.0, 25 °C; 50 mM KCl), MgCl₂ 1.5 mM, dNTPs 250 µM, *Mse*I-N 0.5 µM, 1 U of *Taq* DNA polymerase (TaKaRa, Dalian, China) and 5 µL diluted digestion-ligation DNA. The PCR conditions were 5 min at 94 °C followed by 20 cycles of 30 s at 94 °C, 1 min at 53 °C, 1 min at 72 °C with a final extension of 10 min at 72 °C. The PCR products were hybridized with biotinylated (AG)₁₂ and (GT)₁₂ for 1 h at 68 °C, respectively, after denaturation for 5 min at 95 °C. The DNA fragments that hybridized to biotinylated probes were selectively captured by streptavidin-coated magnetic beads (Streptavidin Magnesphere Paramagnetic Particles, Promega, Shanghai, China). Microsatellite-rich fragments were amplified by *Mse*I-N primers for 35 cycles and cloned into an *Escherichia coli* strain (DH5α) using pGEM-T Easy vectors (Promega). Insert-positive bacterial clones identified by blue-white selection were amplified using M13 primers and visualized by agarose gel electrophoresis. A total of 132 positive clones were chosen for sequencing.

Because the flanking regions of some sequences were too short, and only 41 pairs of primers were designed and synthesized in the end. In addition, we attached FAM, HEX and TAMRA fluorophores at the 5' ends of each forward primer. PCR was conducted in 25 μ L volumes containing PCR buffer, 1.5 mM MgCl₂, 200 μ M of each dNTP, 50 ng genomic DNA, 0.75 U *Taq* polymerase, and 4 pmol of primer. Amplification included an initial denaturation step at 94 °C for 4 min, followed by 42 cycles of 50 s at 94 °C, 50 s at 54–63 °C depending on the primer pair (Table 1), 1 min at 72 °C, and a final extension for 10 min at 72 °C. The PCR products of 3 fluorophores (FAM, HEX and TAMRA) were mixed in a ratio that was based on the brightness of bands visualized by agarose gel electrophoresis. Parameters relevant to multiplex analysis of 10 loci are included in Table 1. PCR products were analyzed using an ABI 3730XL DNA sequencer. Electropherograms were derived using Gene Scan 4.0 and used to deduce DNA fragment sizes using Gene Mapper 4.0 (Sangon Biotech, Shanghai).

Cross-species amplification was tested in 5 closely related whitefly species (*Aleurodicus dispersus*, *Pealius mori*, *Aleuroclava aucubae*, *Bemisia tabaci* MEAM1 and *B. tabaci* MED). DNA samples from 10 individuals of each species were tested.

The number of alleles at each polymorphic locus, size range, and heterozygosities (both observed and expected) were calculated using Cervus 2.0 software (Marshall et al. 1998). Deviation from Hardy-Weinberg equilibrium (HWE) and linkage disequilibrium at each locus were calculated using GenePop 3.4 software (Rousset 2008). The null allele frequency was calculated using Micro-Checker (Van Oosterhout et al. 2004).

RESULTS

The 41 pairs of primers described above were used for screening microsatellite polymorphism. Twenty-eight loci successfully amplified the target regions, while only 10 loci revealed high microsatellite polymorphisms. All of the polymorphic information contents (PIC) of these 10 primers were greater than 0.5. The number of alleles ranged from 12 to 27 across the 60 individuals tested. The expected heterozygosity (H_e) and the observed heterozygosity (H_o) ranged from 0.851 to 0.958 and from 0.621 to 0.850, respectively (Table 1). Null allele frequencies at those loci ranged from 0.0352 to 0.1619 and all loci exhibited no significant evidence of null alleles, but deviated from Hardy-Weinberg equilibrium based on testing of 60 unsexed individuals. Table 2 provided characteristics of the loci for *A. spiniferus* from 48 female individuals in which all loci also deviated from the Hardy-Weinberg equilibrium, and no loci exhibited significant evidence of null alleles. In addition, loci C19-2 and D14, and also I20 and

F12 manifested linkage disequilibrium after sequential Bonferroni correction.

The results of cross-species amplification showed that all loci could be used for amplification at different levels except F12 (Table 3). In particular, loci A21-2, D14, F13-2, F42-2, and I20 were readily amplified in all 5 additional whitefly species.

DISCUSSION

Traditionally, microsatellite markers are developed by extensive screening for microsatellite containing clones through repetitive hybridization of a repeat motif probe to a large number of random clones (Rassmann et al. 1991). Such an isolation strategy resulted in low rate of the number of positive clones (containing microsatellites) detection (Zane et al. 2002). Using modified protocols of Hamilton et al (1999) and Glenn et al (2000) to construct and clone genomic libraries increased proportions of inserts that contained tandem repeat arrays. A great number of microsatellite repeat regions have been detected, sequenced and used to design specific flanking primers for microsatellite amplification.

Our studied loci indicated a high degree of polymorphism (PIC > 0.5), and can be successfully used to study the population genetic structures of *A. spiniferus* and other closely related whitefly species. The 10 loci still deviated from Hardy-Weinberg equilibrium when another 48 female whiteflies were tested, although we avoided the influence factor of reproduction (Table 2), which was consistent with the findings of Kobmoo et al. (2009), who found that 4 out of 10 loci in *Ceratosolen fusciceps* Mayr (Hymenoptera: Agaonidae) deviated from the Hardy-Weinberg equilibrium (HWE), and the global test of deviation from the HWE equilibrium was significant ($P < 0.01$) even when only females were tested. This deviation occurred because the involved individuals were indeed inbreeding. It is important to note that the HWE test is a low power statistical test to identify genotyping errors (Leal 2005). Thus, the reasons for deviating from the HWE by *A. spiniferus* may be due to other factors such as mutation, natural selection, small population size (genetic drift and population bottle necks) or presence of population substructures including Wahlund's effect (<http://www.dorak.info/genetics/popgen.html>). Specifically, Martins et al. (2012) reported that 11 of the 15 microsatellite loci in *Planococcus citri* (Hemiptera: Pseudococcidae) significantly deviated from the HWE, probably due to data structuring, since individuals from 2 significantly distant geographic areas were used. Probably this is a potential factor affecting the HWE for *A. spiniferus*, since we used 48 females from Jiangsu (N 32.41° E 119.42°) and Sichuan (N 30.57° E 104.07°) provinces for testing. Moreover, small population size results in random sampling errors as well as unpredictable

TABLE 1. CHARACTERISTICS OF POLYMORPHIC MICROSATELLITE LOCI OF *ALEUROCANTHUS SPINIFERUS* FROM 60 INDIVIDUALS COLLECTED FROM TEA PLANTS IN 6 PROVINCES (SHANDONG, JIANGSU, ZHEJIANG, HUNAN, GUIZHOU, AND YUNNAN) IN CHINA.

| Locus | Genbank Accession no. | Repeat motif | Primer sequences (5'–3') | Size range (bp) | Ta (°C) | NA | HO | HE | PIC | Null freq |
|-------|--------------------------|---|---|--------------------|---------|----|-------|-------|-------|-----------|
| A21-2 | JF422773 | (GA) ₆ GC(GA) ₃₆ | F: CTGGTGATGGACAAAGATA R: GACATTGAGACTGTGATAAC | 254-284 | 61 | 14 | 0.830 | 0.919 | 0.903 | 0.0469 |
| B48-2 | JF422774 | (GT) ₇ AT(GT) ₃ | F: CAGTTTACGCCCTTTTAC R: AATAGAGTTCAAGTCCC | 164-206 | 54 | 21 | 0.800 | 0.922 | 0.908 | 0.0656 |
| C19-2 | JF422775 | (CT) ₂₅ | F: CTCGCAATTCCTTCAACC R: ACATCGCCAACTGCCTT | 179-207 | 65 | 15 | 0.850 | 0.922 | 0.907 | 0.0352 |
| D14 | JF422776 | (GT) ₂₄ | F: GCTGCTATCCCCACTCT R: AACGATTGCTCGCCTC | 203-257 | 63 | 20 | 0.846 | 0.926 | 0.911 | 0.0404 |
| F12 | JF422777 | (GT) ₂₅ GA(GT) ₁₁ | F: GGTTAGGCTGGGTAAAT R: ACACCTCGGTAGGATAGT | 225-281 | 63 | 27 | 0.717 | 0.958 | 0.948 | 0.1394 |
| F13-2 | JF422778 | (CA) ₃₁ | F: ACGATTTGCTCGCCTCA R: GCTGCTATCCCCACTCT | 188-244 | 63 | 21 | 0.828 | 0.928 | 0.915 | 0.0536 |
| F42-2 | JF422779 | (CA) ₂₃ | F: ACGATTTGCTCGCCTCA R: CTGCTATCCCCACTCTGT | 187-243 | 63 | 18 | 0.732 | 0.930 | 0.917 | 0.1173 |
| I20 | JF422780 | (TG) ₁₇ | F: CTAAAGGGACTCTGGTTC R: GGATGTTGCGTAAGGTGT | 133-161 | 56 | 12 | 0.621 | 0.851 | 0.825 | 0.1494 |
| I29 | JF422781 | (CA) ₂₂ | F: ACGATTTGCTCGCCTCAG R: TGCTGCTATCCCCACTCTG | 190-242 | 63 | 17 | 0.706 | 0.918 | 0.902 | 0.1267 |
| I47-2 | JF422782 | (GA) ₆ GC(GA) ₃₆ | F: CGTGTAACCAACATAACCC R: TCCTCGGAAGTGACAGAT | 245-277 | 59 | 14 | 0.635 | 0.877 | 0.856 | 0.1619 |

Acronyms: Ta, annealing temperature; N_a, observed number of alleles; H_o, observed heterozygosity; H_e, expected heterozygosity; PIC, Polymorphic information con tent; Null freq, null allele frequency.

TABLE 2. CHARACTERISTICS OF POLYMORPHIC MICRESATELLITE LOCI FOR *ALEUROCANTHUS SPINIFERUS* FROM 48 FEMALE INDIVIDUALS FROM TEA PLANTS IN JIANGSU AND SICHUAN PROVINCES.

| Locus | N _A | H _O | H _E | PIC | Null freq | P-HW |
|-------|----------------|----------------|----------------|-------|-----------|---------|
| A21-2 | 13 | 0.632 | 0.784 | 0.749 | 0.0655 | 0.0000* |
| B48-2 | 19 | 0.814 | 0.908 | 0.890 | 0.0434 | 0.0000* |
| C19-2 | 17 | 0.310 | 0.853 | 0.830 | 0.4788 | 0.0000* |
| D14 | 21 | 0.977 | 0.874 | 0.853 | -0.0724 | 0.0000* |
| F12 | 21 | 0.809 | 0.874 | 0.854 | 0.0058 | 0.0000* |
| F13-2 | 31 | 0.938 | 0.877 | 0.858 | -0.0491 | 0.0068* |
| F42-2 | 26 | 0.872 | 0.785 | 0.752 | -0.0715 | 0.0000* |
| I20 | 27 | 0.938 | 0.922 | 0.907 | -0.0163 | 0.0000* |
| I29 | 16 | 0.636 | 0.790 | 0.768 | 0.0849 | 0.0032* |
| I47-2 | 32 | 0.800 | 0.876 | 0.859 | 0.0174 | 0.0000* |

Acronyms: N_A, observed number of alleles; H_O, observed heterozygosity; H_E, expected heterozygosity; PIC, Polymorphic information content; Null freq, null allele frequency; P-HW, value from the exact test for Hardy–Weinberg equilibrium; *denotes a significant deviation from Hardy–Weinberg equilibrium (P < 0.01), after sequential Bonferroni’s correction of the significance threshold.

TABLE 3. RESULTS OF CROSS-SPECIES AMPLIFICATION TESTS INVOLVING DNA SAMPLES FROM 10 INDIVIDUALS OF EACH OF 5 CLOSELY RELATED WHITEFLY SPECIES.

| Locus | <i>Aleurodicus disperses</i> (n = 10) | <i>Pealius mori</i> (n = 10) | <i>Aleuroclava aucubae</i> (n = 10) | <i>Bemisia tabaci</i> MEAM1 (n = 10) | <i>Bemisia tabaci</i> MED (n = 10) |
|-------|--|---------------------------------|--|---|---------------------------------------|
| A21-2 | ✓ | ✓ | ✓ | ✓ | ✓ |
| B48-2 | × | × | × | ✓ | ✓ |
| C19-2 | × | × | × | ✓ | ✓ |
| D14 | ✓ | ✓ | ✓ | ✓ | ✓ |
| F12 | × | × | × | × | × |
| F13-2 | ✓ | ✓ | ✓ | ✓ | ✓ |
| F42-2 | ✓ | ✓ | ✓ | ✓ | ✓ |
| I20 | ✓ | ✓ | ✓ | ✓ | ✓ |
| I29 | ✓ | × | × | × | × |
| I47-2 | ✓ | ✓ | ✓ | × | ✓ |

✓, Successful amplification; ×, unsuccessful amplification.

genotype frequencies; and in small populations, departure from expected genotype frequencies occurs more easily. Probably this is another factor because we used 60 samples from 6 provinces (10 from each location) in the beginning. Furthermore, as mentioned above, a very high mutation rate in the population (typical mutation rates are < 10⁻⁴ per generation) and selection of one or a combination of genotypes (Ineichen & Batschelet 1975) also could violate the HWE. Clearly, further population genetic structure studies are urgently needed to assess these hypotheses.

Our results may also provide new insights about the reproductive mechanism of *A. spiniferus*. Since few individuals engaged in arrhenotokous reproduction, the reproductive strategies of this insect may not significantly affect Hardy–Weinberg equilibrium of the loci studied.

Microsatellites are particularly reliable for studying recent biological invasions (Zygouridis et al. 2009) and are a powerful means to detect the

origin and invasion route of exotic insect species (Fonseca et al. 2010). Our cross-species amplification results indicated that the 10 microsatellite polymorphic loci may possess high versatility, and can be used to study gene flow, genetic diversity and population genetic structure of related whitefly species. Meanwhile, this information will facilitate the selection of molecular markers for *B. tabaci*, a controversial invasive species worldwide. Overall, these microsatellite markers show suitable resolution to meet our needs in subsequent studies and to give us insights into the genetic variability, gene flow and the mating system of whitefly species.

ACKNOWLEDGMENTS

This research was funded by the Special Fund for Agro-scientific Research in the Public Interest of China (201303019, No. 200803005).

REFERENCES CITED

- ANONYMOUS. 2011. EPPO A2 List of pests recommended for regulation as quarantine pests (version 2011-09). URL: <http://www.eppo.int/QUARANTINE/listA2.htm>
- CARLETON, K. L., STREELMAN, J. T., LEE, B. Y., GARNHART, N., KIDD, M., AND KOCHER, T. D. 2001. Rapid isolation of CA microsatellite from the tilapia genome. *Animal Genetics* 33: 140-144.
- DUBEY, A. K., AND KO, C. C. 2012. Sexual dimorphism among species of *Aleurocanthus* Quaintance & Baker (Hemiptera: Aleyrodidae) in Taiwan, with one new species and an identification key. *Zootaxa* 3177: 1-23.
- EVANS, G. A. 2008. The whiteflies (Hemiptera: Aleyrodidae) of the world and their host plants and natural enemies. http://www.sel.barc.usda.gov:591/1WF/whitefly_catalog.htm. (Accessed 23-IX-2008).
- FONSECA, D. M., WIDDEL, A. K., HUTCHINSON, M., SPICHER, S. E., KRAMER, L. D. 2010. Fine-scale spatial and temporal population genetics of *Aedes japonicus*, a new US mosquito, reveal multiple introductions. *Mol Ecol* 19: 1559-1572.
- FU, J. Y. AND HAN, B. Y. 2007. A RAPD analysis on genetic diversity of populations of *Aleurocanthus spiniferus* from tea gardens in eastern China. *Acta Ecol. Sinica* 27(5): 1887-1894.
- GLENN, T. C., CARY, T., DUST, M., HAUSWALDT, S., PRINCE, K., RAMSDELL, C., AND SHUTE, I. M. 2000. Microsatellite Isolation. www.uga.edu/srel/DNA_Laboratory/protocols.htm.
- GORMAN, M. J., SEVERSON, D. W., CORNEL, A. J., COLLINS, F. H., AND PASKEWITZ, S. M. 1997. Mapping a quantitative trait locus involved in melanotic encapsulation of foreign bodies in the malaria vector, *Anopheles gambiae*. *Genetics* 146: 965-971.
- GUO, L., QIU, B. L., REN, S. X., AND WU, H. J. 2006. Summary on the classify of natural enemy germplasm resources of *Aleurocanthus spiniferus* (Homoptera: Aleyrodidae). *Guangdong Agric. Sci.* 2: 002.
- HAMILTON, M. B., PINCUS, E. L., FIORE, A. D., AND FLEISCHER, R. C. 1999. Universal linker and ligation procedures for construction of genomic DNA libraries enriched for microsatellites. *BioTechniques* 27: 500-507.
- HAN, B., AND CUI, L. 2002. Natural population life table of citrus spiny whitefly (*Aleurocanthus spiniferus*) in tea garden. *Acta Ecol. Sinica* 23(9): 1781-1790.
- HUANG, J., LUO, X. N., HUANG, B. K., AND ZHANG, X. Y. 1999. Studies on *Aleurocanthus spiniferus* (Quaintance) and its natural enemies. *Entomol. J. East China* 8(1): 35-40.
- INEICHEN, R., AND BATSCHELET, E. 1975. Genetic selection and de Finetti diagrams. *J. Math. Biol.* 2(1): 33-39.
- KANMIYA, K., USDA, S., KASAI, A., YAMASHITA, K., SATO, Y., AND YOSHIYASU, Y. 2011. Proposal of new specific status for tea-infesting populations of the nominal citrus spiny whitefly *Aleurocanthus spiniferus* (Homoptera: Aleyrodidae). *Zootaxa* 2797: 25-44.
- KOBMOO, N., VIGNES, H., HOSSARET-MCKEY, M., WEI, Z., AND KJELLBERG, F. 2009. Isolation and characterization of microsatellite primers for *Ceratosolen fusciceps*, the fig-pollinating wasp of *Ficus racemosa*, and amplification in two populations. *Mol. Ecol. Resour.* 9(4): 1147-1150.
- LEAL, S. M. 2005. Detection of genotyping errors and pseudo-SNPs via deviations from Hardy-Weinberg equilibrium. *Genet. Epidemiol.* 29(3): 204-214.
- MARSHALL, T. C., SLATE, J., KRUIK, L. E. B., AND PEMBERTON, J. M. 1998. Statistical confidence for likelihood-based paternity inference in natural populations. *Mol. Ecol.* 7: 639-655.
- MARTINS, R. F., ZINA, V. E. R. A., DA SILVA, E. B., REBELO, M. T., FIGUEIREDO, E., MENDEL, Z., PAULO, O. S. FRANCO, J. C., AND SEABRA, S. G. 2012. Isolation and characterization of fifteen polymorphic microsatellite loci for the citrus mealybug, *Planococcus citri* (Hemiptera: Pseudococcidae), and cross-amplification in two other mealybug species. *J. Genet.* 91: e75-e78.
- MUNIAPPAN, R., PUREA, M., SENGEBAU, F., AND REDDY, G. V. P. 2006. Orange spiny whitefly, *Aleurocanthus spiniferus* (Quaintance) (Homoptera: Aleyrodidae), and its parasitoids in the Republic of Palau. *Proc. Hawaiian Entomol. Soc.* 38: 21-25.
- NIU, J. Z., HULL-SANDERS, H., ZHANG, Y. X., LIN, J. Z., DOU, W., AND WANG, J. J. 2014. Biological control of arthropod pests in citrus orchards in China. *Biol. Control* 68: 15-22.
- PENG, P., TANG, M., HOU, Y. J., LIN, Q., HUANG, S. J., DENG, M., HU, X., AND ZHANG, Y. 2010. Study on the Effect and Characters of Yellow Sticky Trap Sticking *Aleurocanthus spiniferus* and *Empoasca vitis* Gothe in Tea Garden. *Southwest China J. Agric. Sci.* 23(1): 87-90.
- PETERLUNGER, E., GASPERO, G. D., CIPRIANI, G., SIVILOTTI, P., ZULINI, L., MARRAZZO, M. T., ANDRETTA, D., AND TESTOLIN, R. 2003. Breeding strategy for the introgression of disease resistance genes into European grapevine. *Acta Hort.* 603: 665-670.
- RASSMANN, K., SCHLOTTERER, C., AND TAUTZ, D. 1991. Isolation of simple sequence loci for use in polymerase chain reaction-based DNA fingerprinting. *Electrophoresis* 12: 113-118.
- ROUSSET, F. 2008. GenePop'007: a complete re-implementation of the GenePop software for Windows and Linux. *Mol. Ecol. Resour.* 8: 103-106.
- TENG, X., WU, Q., AND WAN, F. H. 2009. A salting-out method for genomic DNA extraction from *Bemisia tabaci* (Gennadius). *Biotechnol. Bull.* (8): 166-168.
- VAN DEN BERG, M. A., HOPPER, G., AND GREENLAND, J. 2000. An economic study of the biological control of the spiny blackfly, *Aleurocanthus spiniferus* (Hemiptera: Aleyrodidae), in a citrus orchard in Swaziland. *Biocontrol Sci. Technol.* 10(1): 27-32.
- VAN OOSTERHOUT, C., HUTCHINSON, W. F., WILLS, D. P., AND SHIPLEY, P. 2004. Micro-Checker: software for identifying and correcting genotyping errors in microsatellite data. *Mol. Ecol.* 4: 535-538. www.uga.edu/srel/DNA_Lab/protocols.htm.
- ZANE, L., BARGELLONI, L., AND PATANELLO, T. 2002. Strategies for microsatellite isolation: a review. *Mol. Ecol.* 11(1): 1-16.
- ZENGER, K. R., ELDRIDGE, M. D. B., AND JOHNSTON, P. G. 2005. Phylogenetics, population structure and genetic diversity of the endangered southern brown bandicoot (*Isodon obesulus*) in south-eastern Australia. *Conserv. Genetics* 6: 193-204.
- ZYGOURIDIS N. E., AUGUSTINOS A. A. ZALOM, F. G., AND MATHIOPOULOS, K. D. 2009. Analysis of olive fly invasion in California based on microsatellite markers. 102: 402-412.