

## **Molecular cloning of heat shock protein gene HSP90 and effects of abamectin and double-stranded RNA on its expression in *Panonychus citri* (Trombidiformes: Tetranychidae)**

Authors: Tian, Huixia, Yu, Shijiang, Liu, Bin, Yang, Juan, Li, Xiaojiao, et al.

Source: Florida Entomologist, 98(1) : 37-43

Published By: Florida Entomological Society

URL: <https://doi.org/10.1653/024.098.0107>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

# Molecular cloning of heat shock protein gene HSP90 and effects of abamectin and double-stranded RNA on its expression in *Panonychus citri* (Trombidiformes: Tetranychidae)

Huixia Tian<sup>1</sup>, Shijiang Yu<sup>1</sup>, Bin Liu<sup>3</sup>, Juan Yang<sup>2</sup>, Xiaojiao Li<sup>1</sup>, Jiansu Yue<sup>1</sup>, Fei Chen<sup>3</sup>, Lili Ding<sup>2</sup>, Haoqiang Liu<sup>2</sup>, Hongjun Li<sup>2</sup>, Lin Cong<sup>2</sup> and Chun Ran<sup>1,2,\*</sup>

## Abstract

*Panonychus citri* McGregor (Trombidiformes: Tetranychidae), the widely distributed citrus red mite, has developed resistance to most registered acaricides. Adaptation of arthropods to extreme environmental conditions has been related to increased expression of their heat shock proteins (HSPs). The objectives of this study were to explore the relationship between HSPs and resistance of *Panonychus citri* to the acaricide abamectin and the adaptation of *Panonychus citri* to high temperatures. The full-length cDNA of the HSP90 gene was cloned from an abamectin-sensitive strain of *Panonychus citri*. This gene consisted of 2,495 nucleotides with a complete open reading frame (ORF) of 2,169 nucleotides. This gene encoded a polypeptide of 721 amino acids with a predicted molecular weight of 83.44 kDa, a theoretical isoelectric point of 5.06, a 3' untranslated region (UTR) of 228 bp, and a 5' UTR of 98 bp. The results of real-time PCR analyses indicated that the expression of the HSP90 gene in *P. citri* was markedly affected by the concentration of abamectin, the duration of exposure to it and the temperature, suggesting that the up-regulation of the HSP90 gene may play an important role in abamectin resistance and adaptation to high temperatures in *Panonychus citri*. The results of RNA interference experiments indicated that the HSP90 gene from adult female *Panonychus citri* was sensitive to down-regulation by double-stranded RNA (0.1–0.2 µg/µL). This study provides a molecular basis for further analysis of the relationships between the HSP90 gene and the resistance of *Panonychus citri* to abamectin and to high temperatures.

Key Words: citrus red mite; abamectin; HSP90; double-stranded RNA; resistance mechanism

## Resumen

*Panonychus citri* McGregor (Trombidiformes: Tetranychidae), el ácaro rojo de los cítricos ampliamente distribuido, ha desarrollado resistencia a la mayoría de los acaricidas registrados. La adaptación de los artrópodos a condiciones ambientales extremas se ha relacionado con el aumento de expresión de sus proteínas de choque térmico (HSPs, en inglés). El objetivo de este estudio fue explorar la relación entre el HSP y la resistencia de *Panonychus citri* al acaricida, abamectina, y la adaptación de *Panonychus citri* a altas temperaturas. El ADNc de longitud completa del gen HSP90 fue clonado a partir de una cepa de *Panonychus citri* sensible de abamectina. Este gen consistió en 2,495 nucleótidos con un marco de lectura abierto completo (ORF) de 2169 nucleótidos. Este gen codifica un polipéptido de 721 aminoácidos con un peso molecular predicho de 83.44 kDa, un punto isoelectrónico teórico de 5.06, una región 3' no traducida (UTR) de 228 pb y una 5' UTR de 98 pb. Los resultados del análisis de PCR en tiempo real indicaron que la expresión del gen HSP90 en *P. citri* fue marcadamente afectada por la concentración de abamectina, la duración de la exposición a la misma y la temperatura, lo que sugiere que la sobre regulación del gen HSP90 puede jugar un papel importante en la resistencia a la abamectina y a la adaptación a las altas temperaturas en *P. citri*. Los resultados de los experimentos de interferencia de ARN indicaron que el gen HSP90 de la hembra adulta de *Panonychus citri* fue sensible a la baja regulación de ARN de doble cadena (0.1-0.2 g/l). Este estudio provee una base molecular para el análisis adicional de las relaciones entre el gen HSP90 y la resistencia de *Panonychus citri* a la abamectina y a altas temperaturas.

Palabras Clave: ácaro del rojo de los cítricos; abamectina; HSP90; ARN de doble cadena; mecanismo de resistencia

The citrus red mite, *Panonychus citri* McGregor (Trombidiformes: Tetranychidae), is a citrus pest widely distributed throughout the world, including in the citrus growing areas of China (Liu et al. 2011; Xia et al. 2014). *P. citri* punctures and feeds on the contents of cells,

<sup>1</sup>College of Horticulture and Landscape Architecture, Southwest University, Key Laboratory of the Southern Mountain Horticulture of the Ministry Education, Chongqing 400716, China

<sup>2</sup>Citrus Research Institute, Chinese Academy of Agricultural Sciences, National Citrus Engineering Research Center, Chongqing 400712, China

<sup>3</sup>Key Laboratory of Entomology and Pest Control Engineering, College of Plant Protection, Southwest University, Chongqing 400716, China

\*Corresponding author; E-mail: ranchun@cric.cn

mainly of leaves, tender shoots, flower buds and fruit of citrus plants. The severe damage these mites cause induces loss of leaves, flowers and fruits and reduces citrus yield. Chemical control remains the most effective method for controlling *P. citri*. However, *P. citri* easily develops resistance to acaricides because of the mite's minuscule size, rapid production of generations and strong reproductive performance. *Panonychus citri* is resistant to most currently registered acaricides (Furuhashi 1994; Hu et al. 2010; Ouyang et al. 2012). The new antibiotic acaricide abamectin is effective against eggs, larvae and adult mites. Since abamectin was first introduced onto the Chinese market in 1991, it has played an important role in control of mite pests in China. Currently, more than 10 companies produce the active ingredient, abamectin, and approximately 300 organizations have registered abamectin formulations, of which approximately 140 enterprises have registered abamectin formulations to control *P. citri* (Chemical Book. 2010; China Pesticide Information Network). However, due to the long-term application of abamectin, many pest mite species have evolved resistance to it, and abamectin-resistant *P. citri* have appeared in some areas of citrus production (Zhang et al. 2013), hindering the sustainable control of this pest. Revealing the molecular mechanism of this resistance is crucial for coping with the resistance problem.

Heat shock proteins (HSPs) are rapidly generated in both prokaryotes and eukaryotes under environmental stress. The increase of HSPs in arthropods is related to their adaptation to extreme temperature, drought, radiation and pesticides (Feder & Hofmann 1999; Padmini 2010; Akerfelt et al. 2011; Zhu et al. 2013). A recent study indicated that the abamectin resistance in *Tetranychus cinnabarinus* (Boisduval) (Trombidiformes: Tetranychidae) may be associated with an increase in HSP90 gene expression (Feng et al. 2010). Some abamectin resistance has been generated in *P. citri*, but the molecular mechanism for this resistance remains unclear. Therefore, it is essential to elucidate the relationship between HSPs and abamectin resistance in *P. citri*.

RNA interference (RNAi) has been widely applied in the analysis of gene function (Fire et al. 1998; Campbell et al. 2010; Li et al. 2011). It is a post-transcriptional gene silencing process induced by double-stranded RNA (dsRNA), with gene expression inhibited by interference in the translation or transcription of specific genes. Due to the small size of individual mites, few studies have been conducted by employing RNAi technology in mites, and to date, no studies have applied RNAi technology to examine gene function in *P. citri*. Thus, we sought to establish an appropriate RNAi methodology to systematically analyze the function of HSPs in *P. citri*.

To determine the relationship between HSPs and the abamectin resistance of *P. citri*, as well as to examine the mechanism for the adaptation of *P. citri* to high temperatures, the complete HSP90 gene sequence from an abamectin-sensitive strain of *P. citri* was cloned. The influence of abamectin on the expression of the HSP90 gene in *P. citri* adult females by various abamectin concentrations and temperatures was analyzed. In addition, a methodology for silencing the HSP90 gene was developed using RNAi to systematically analyze the function of HSPs in *P. citri*.

## Materials and Methods

### EXPERIMENTAL MATERIALS

A relatively abamectin-sensitive *P. citri* mite strain, bred on a lemon tree in an abamectin-free field in the Citrus Research Institute of the Chinese Academy of Agricultural Sciences, was collected in 2013. The experimental population was established with lemon leaves as the host plant, and adult female mites (15 days old) were

the experimental materials. *Panonychus citri* was reared indoors using the leaf disc method (Li et al. 2014). The culture dish was 9 cm in diam, and the dish was covered with a wet sponge of the same diam. An explanted fresh lemon leaf was placed over the sponge with the abaxial side facing upward, and the leaf margin was surrounded by a degreased cotton strand. This strain was maintained in an indoor laboratory at  $25 \pm 1$  °C with 70–80% relative humidity (RH) and a 14:10 h light: dark cycle.

The 1.8% abamectin EC was purchased from Liuzhou Huinong Chemical Co., Ltd. (Liuzhou, Guangxi).

### CLONING AND SEQUENCE ANALYSIS OF THE HSP90 GENE FROM *P. CITRI*

After the protein sequences of HSP90 from *Ixodes scapularis*, *Xenopus tropicalis*, *Chiromantes haematocheir*, *Tigriopus japonicus*, *Liriomyza sativae*, and *Equus caballus* were obtained from NCBI, 2 pairs of degenerate primers were designed with Primer Premier 5.0 to amplify the HSP90 fragment: HSP90-DF1 (5'-GGACCAAGGCCTTCATGgargcnytnca-3'), HSP90-DR1 (5'-GCCCGGAAGTCCAGTgncctcncac-3') and HSP90-DF2 (5'-GCCAAGTCCGGCACCAGGcNtntatgarg-3'), HSP90-DR2 (5'-CATGATGAACCCGCGCacrtanarytt-3'). The same sequence was obtained while the middle fragment was compared with transcriptome database, and 2 specific primers were designed for each fragment: HSP90-5GSPR1 (5'-CCACCAGCCGAGGACTCCCACACAT-3'), HSP90-5GSPR2 (5'-TACCACGACCAAGGGGCTCACCAGT-3') and HSP90-3GSPF1 (5'-GAAGACAAGGGACAAGGAAGTCTAGCGA-3'), HSP90-3GSPF2 (5'-AAACCAAGCCTATCTGGACCCGCAAT-3'). The sequences of the 5'-end and 3'-end of HSP90 gene from *P. citri* were amplified.

The total RNA of *P. citri* was extracted based on the instructions provided in the RNAPrep Pure Micro kit (TIANGEN). The PrimeScriptRT reagent kit with gDNA Eraser (TaKaRa) was used to synthesize the cDNA. The PCR amplification was conducted according to the method of Staheli et al (2009). The amplification of the cDNA ends was conducted according to the instructions provided in the SMARTerRACE cDNA Amplification kit and the Advantage 2 PCR kit (Clontech).

DNA sequences were subjected to the BLAST tool on the NCBI website (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>). The 5'- and 3'-end sequences were spliced with DNAMAN 5.0. The online software ORF Finder from NCBI (<http://www.ncbi.nlm.nih.gov/gorf/gorf.html>) was used to analyze the open reading frame of the complete HSP90 gene sequence in *P. citri*. Using Mega 4.0 and ClustalX, *P. citri* was compared with 4 insect/mite species showing high amino acid sequence homology to determine the degree of homology.

### BIOASSAY METHOD

The slide impregnation method recommended by Food and Agriculture Organization of the United Nations was employed for treatment applications (Busvine 1980). Based on preliminary experiments, the abamectin solutions were prepared with 5–7 concentrations using a geometric series of mortalities (20%–90%) in *P. citri* adult females. Double-sided adhesive tape (3 cm) was adhered to one end of a clean slide. Healthy adult female mites were picked using a small brush and the dorsal side of the mite was affixed to the tape. Each slide held 30 individuals, and 4 replicates were used for each treatment. For the treatment group, the slides with the adhered mites were immersed in an abamectin solution, mildly agitated for 5 s and then removed. Mites in the control group were similarly treated but with clean water instead of abamectin. Excess solution on the surface of the *P. citri* females was removed with absorbent paper. The mites were then air-dried at room temperature for 15 min and placed into an illumination incubator ( $25 \pm 1$  °C, RH 70–80%). After 24 h, the mites were observed using a bin-

### EFFECT OF ABAMECTIN CONCENTRATIONS AT VARIOUS TEMPERATURES ON HSP90 GENE EXPRESSION IN *P. CITRI* ADULT FEMALES

The RNA and cDNA were extracted based on the method described in section 1.2.2. The upstream and downstream primers were designed with Primer Premier 5.0 to detect the relative expression of HSP90 mRNA from *P. citri*, i.e., HSP90-F: 5'-TGAAAGTGATCCGCAAGAACC-3' and HSP90-R: 5'-CTGTCTCTGTGAATCCAAG-3'. The upstream and downstream primers for the reference gene ELF1A (Niu et al. 2012) were as follows: ELF1A-F: 5'-GGCACTTCGTCTTCCACTTC-3', and ELF1A-R: 5'-ATGATTCTGGTGCATCTCA-3'.

Reference genes are considered genes that maintain stable expression levels under different experimental conditions and in various tissues and cells. Although the relative HSP90 gene expression by the adult female mites changed with temperature and abamectin concentration, the data could be normalized through corrections using the expressed amount of reference genes in the corresponding sample.

The double-stranded RNA (dsRNA) synthesized by Wuhan Cell Marker Biotechnology Co., Ltd. (5'-GCTAGGTTTGGGTATTGAT-3') was diluted (0.1 µg/µL, 0.2 µg/µL and 0.4 µg/µL) and then used to feed adult female mites that were at the same developmental stage. The dsRNA solution was applied under a microscope to the mouth parts of the mites by a clean fine brush (number 0) that had been treated with DEPC (diethyl pyrocarbonate) and dipped in the dsRNA solution. The mites in the control group were supplied with clean water. The adults were placed into an illumination incubator (25 ± 1 °C, RH 70–80%), and the surviving individuals were collected after 6, 12, 24 and 48 h and placed into 1.5 mL centrifuge tubes. After freezing in liquid nitrogen, the specimens were stored at -80 °C.

The mortality rates were corrected by Abbott's formula, and the data were used to fit a toxicity regression equation using SPSS 17.0. A  $\chi^2$  test was used to test the  $LC_{25}$ ,  $LC_{50}$  and  $LC_{75}$  values at the 95% confidence interval to verify the accuracy of the toxicity regression equation using SPSS 17.0 software. The  $2^{-\Delta\Delta CT}$  method was applied to calculate the relative expression of HSP90 mRNA in *P. citri* in the different treatment groups. The expression level in the control group was used as the background value. The difference in the relative expressions of HSP90 mRNA between treatment and control groups examined at the same temperature were analyzed with Student's *t*-test ( $P < 0.05$ ). Differences across treatment groups were examined with multiple comparisons using One-Way ANOVA (LSD method) ( $P < 0.05$ ).

#### CLONING AND SEQUENCE ANALYSIS OF FULL-LENGTH HSP90 GENE FROM *P. CITRI*

[illegible]

Downloaded From: <https://complete.bioone.org/journals/Florida-Entomologist> on 15 Apr 2024  
Terms of Use: <https://complete.bioone.org/terms-of-use>



Multiple alignments were performed with DNAMAN 5.0 on the amino acid sequences of HSP90 genes of *P. citri* McGregor and *Frankliniella occidentalis* (Pergande) (Thysanoptera: Thripidae), *Trialeurodes vaporariorum* Westwood (Hemiptera: Aleyrodidae), *Pteromalus puparum* (L.) (Hymenoptera: Pteromalidae), *Tetranychus cinabarinus* (Boisduval) (Trombidiformes: Tetranychidae). The results indicated that they shared several conserved motifs and displayed high amino acid sequence homology (Fig. 2).

#### ABAMECTIN TOXICITY BIOASSAY

Panonychus citri	.MPFEFMQGDVETFAFQAEIAQLMLNTFYSNKEIFLR	39
Frankliniella oc	.MPFEFMQGDVETFAFQAEIAQLMLNTFYSNKEIFLR	39
Trialeurodes vap	.MPEDVAMEQETFAFQAEIAQLMLNTFYSNKEIFLR	39
Pteromalus pupar	...METAGGEVETFAFQAEIAQLMLNTFYSNKEIFLR	37
Tetranychus cina	MPFETMSVEELFETFAFQAEIAQLMLNTFYSNKEIFLR	40
Consensus	etf fgaqlaqlm li ntfysnkeiflr	
Panonychus citri	ELISNSSDALDKIRYESLTDPKRLDSCKRDEIRIKENKDN	79
Frankliniella oc	ELISNSSDALDKIRYESLTDPKRLDSCKRDEIRIKENKDN	79
Trialeurodes vap	ELISNSSDALDKIRYESLTDPKRLDSCKRDEIRIKENKDN	79
Pteromalus pupar	ELISNSSDALDKIRYESLTDPKRLDSCKRDEIRIKENKDN	77
Tetranychus cina	ELISNSSDALDKIRYESLTDPKRLDSCKRDEIRIKENKDN	80
Consensus	elisnssdaldkiryeslt dp l k l i p	
Panonychus citri	RTLTIIDIGIGMTKADLVNGLTIKSGTAFMEALCAGA	119
Frankliniella oc	RTLTIIDIGIGMTKADLVNGLTIKSGTAFMEALCAGA	119
Trialeurodes vap	RTLTIIDIGIGMTKADLVNGLTIKSGTAFMEALCAGA	119
Pteromalus pupar	RTLTIIDIGIGMTKADLVNGLTIKSGTAFMEALCAGA	117
Tetranychus cina	RTLTIIDIGIGMTKADLVNGLTIKSGTAFMEALCAGA	120
Consensus	rtlt id gigmgtkadi nngltiaksgt afmealaga	
Panonychus citri	DISMIGQFGVGFYSALVADRVVYVSKHNDDCYWESSA	159
Frankliniella oc	DISMIGQFGVGFYSALVADRVVYVSKHNDDCYWESSA	159
Trialeurodes vap	DISMIGQFGVGFYSALVADRVVYVSKHNDDCYWESSA	159
Pteromalus pupar	DISMIGQFGVGFYSALVADRVVYVSKHNDDCYWESSA	157
Tetranychus cina	DISMIGQFGVGFYSALVADRVVYVSKHNDDCYWESSA	160
Consensus	dismigqfgvgfyasa lvadrvv yv skhndd cywes a	
Panonychus citri	GGSFITIKHDTTGELRCGTIVLHMKEDQTEVLEPKEKE	199
Frankliniella oc	GGSFITIKHDTTGELRCGTIVLHMKEDQTEVLEPKEKE	198
Trialeurodes vap	GGSFITIKHDTTGELRCGTIVLHMKEDQTEVLEPKEKE	199
Pteromalus pupar	GGSFITIKHDTTGELRCGTIVLHMKEDQTEVLEPKEKE	196
Tetranychus cina	GGSFITIKHDTTGELRCGTIVLHMKEDQTEVLEPKEKE	199
Consensus	ggsft itgkgti ked l e k ke	
Panonychus citri	IVKKHSQFYGYPKILVCKRERKESIDEPEE...KKDKD	238
Frankliniella oc	IVKKHSQFYGYPKILVCKRERKESIDEPEE...KKDKD	238
Trialeurodes vap	IVKKHSQFYGYPKILVCKRERKESIDEPEE...KKDKD	239
Pteromalus pupar	IVKKHSQFYGYPKILVCKRERKESIDEPEE...KKDKD	230
Tetranychus cina	IVKKHSQFYGYPKILVCKRERKESIDEPEE...KKDKD	237
Consensus	ivkhsqfygypikl vckrerk esidepee k k k k k	
Panonychus citri	EDKKEG...DDGTPKEDVGEDEE...KKKKKKKTVKE	272
Frankliniella oc	EDKKEG...DDGTPKEDVGEDEE...KKKKKKKTVKE	272
Trialeurodes vap	EDKKEG...DDGTPKEDVGEDEE...KKKKKKKTVKE	272
Pteromalus pupar	EDKKEG...DDGTPKEDVGEDEE...KKKKKKKTVKE	266
Tetranychus cina	EDKKEG...DDGTPKEDVGEDEE...KKKKKKKTVKE	272
Consensus	k e d pk edv k k k k k e	
Panonychus citri	KYDEEELNFKPIWRNNDITQDEYGEFYKSLTNDWE	312
Frankliniella oc	KYDEEELNFKPIWRNNDITQDEYGEFYKSLTNDWE	312
Trialeurodes vap	KYDEEELNFKPIWRNNDITQDEYGEFYKSLTNDWE	312
Pteromalus pupar	KYDEEELNFKPIWRNNDITQDEYGEFYKSLTNDWE	306
Tetranychus cina	KYDEEELNFKPIWRNNDITQDEYGEFYKSLTNDWE	312
Consensus	ky edeel ntkpiw rn di eygefyk sltndwe	
Panonychus citri	HLAVKHFSVEGQLEFFALLFERRPFDFLENKKKNNIK	352
Frankliniella oc	HLAVKHFSVEGQLEFFALLFERRPFDFLENKKKNNIK	352
Trialeurodes vap	HLAVKHFSVEGQLEFFALLFERRPFDFLENKKKNNIK	352
Pteromalus pupar	HLAVKHFSVEGQLEFFALLFERRPFDFLENKKKNNIK	346
Tetranychus cina	HLAVKHFSVEGQLEFFALLFERRPFDFLENKKKNNIK	352
Consensus	hlavkhfsvegqleff allf err pfdflen k k k n n i k	
Panonychus citri	LYVRRVFIMDNCELIPEYLNFGVVDSEDLPLNISRE	392
Frankliniella oc	LYVRRVFIMDNCELIPEYLNFGVVDSEDLPLNISRE	392
Trialeurodes vap	LYVRRVFIMDNCELIPEYLNFGVVDSEDLPLNISRE	392
Pteromalus pupar	LYVRRVFIMDNCELIPEYLNFGVVDSEDLPLNISRE	386
Tetranychus cina	LYVRRVFIMDNCELIPEYLNFGVVDSEDLPLNISRE	392
Consensus	lyvrrvfimdnce l ipeylnf g v v d s e d l p l n i s r e	

In adult female *P. citri* the lethal concentrations of 1.8% abamectin EC were determined by a bioassay. Thus the LC<sub>25</sub>, LC<sub>50</sub> and LC<sub>75</sub> were determined to be 0.0206 mg/L, 0.0572 mg/L and 0.1586 mg/L, respectively (Table 1).

#### EFFECTS OF ABAMECTIN CONCENTRATION AT 18 °C ON THE RELATIVE EXPRESSION OF HSP90 IN *P. CITRI*

After treatment at 18 °C with various concentrations of abamectin, the relative expression levels of HSP90 in *P. citri* increased within

Panonychus citri	LQQNKILVIRKKNLKKKLEFEEEDKEKYYKYYEQFS	432
Frankliniella oc	LQQNKILVIRKKNLKKKLEFEEEDKEKYYKYYEQFS	432
Trialeurodes vap	LQQNKILVIRKKNLKKKLEFEEEDKEKYYKYYEQFS	432
Pteromalus pupar	LQQNKILVIRKKNLKKKLEFEEEDKEKYYKYYEQFS	426
Tetranychus cina	LQQNKILVIRKKNLKKKLEFEEEDKEKYYKYYEQFS	432
Consensus	lqqnkil virknl kkc lfee edke ykk yeqfs	
Panonychus citri	KNIRLGIHEDSTERRKLABLRYETASAGDEVCSSKDYIS	472
Frankliniella oc	KNIRLGIHEDSTERRKLABLRYETASAGDEVCSSKDYIS	472
Trialeurodes vap	KNIRLGIHEDSTERRKLABLRYETASAGDEVCSSKDYIS	472
Pteromalus pupar	KNIRLGIHEDSTERRKLABLRYETASAGDEVCSSKDYIS	466
Tetranychus cina	KNIRLGIHEDSTERRKLABLRYETASAGDEVCSSKDYIS	472
Consensus	kn klgihed sterrklabl ry etasagde cs k dy	
Panonychus citri	RMKENQKCIYMITGESREQVANSFVFRVVRGRFVVMVT	512
Frankliniella oc	RMKENQKCIYMITGESREQVANSFVFRVVRGRFVVMVT	512
Trialeurodes vap	RMKENQKCIYMITGESREQVANSFVFRVVRGRFVVMVT	512
Pteromalus pupar	RMKENQKCIYMITGESREQVANSFVFRVVRGRFVVMVT	506
Tetranychus cina	RMKENQKCIYMITGESREQVANSFVFRVVRGRFVVMVT	512
Consensus	rmkenqk iy itges qva s fvrvr grf vvmv	
Panonychus citri	EPIDEMVQCCREDFGKLVSVTRKLELPEDEEKKKRE	552
Frankliniella oc	EPIDEMVQCCREDFGKLVSVTRKLELPEDEEKKKRE	552
Trialeurodes vap	EPIDEMVQCCREDFGKLVSVTRKLELPEDEEKKKRE	552
Pteromalus pupar	EPIDEMVQCCREDFGKLVSVTRKLELPEDEEKKKRE	546
Tetranychus cina	EPIDEMVQCCREDFGKLVSVTRKLELPEDEEKKKRE	552
Consensus	epidey vqq r e dfgk lvsvtrkleglpe e e e k k k e	
Panonychus citri	EDAKKEELCKVMKILDLKRVKVVVSNRIVSPCCIVTS	592
Frankliniella oc	EDAKKEELCKVMKILDLKRVKVVVSNRIVSPCCIVTS	592
Trialeurodes vap	EDAKKEELCKVMKILDLKRVKVVVSNRIVSPCCIVTS	592
Pteromalus pupar	EDAKKEELCKVMKILDLKRVKVVVSNRIVSPCCIVTS	586
Tetranychus cina	EDAKKEELCKVMKILDLKRVKVVVSNRIVSPCCIVTS	592
Consensus	d k e lckvmk ild rkvk vvr snriv spccivts	
Panonychus citri	CYGVANERIMKAQALRSTMYAAKKHLEINPDHAI	632
Frankliniella oc	CYGVANERIMKAQALRSTMYAAKKHLEINPDHAI	632
Trialeurodes vap	CYGVANERIMKAQALRSTMYAAKKHLEINPDHAI	632
Pteromalus pupar	CYGVANERIMKAQALRSTMYAAKKHLEINPDHAI	626
Tetranychus cina	CYGVANERIMKAQALRSTMYAAKKHLEINPDHAI	632
Consensus	cygv an erimkaqalr st m yaakkhle inpdh	
Panonychus citri	VEDIRKKEAENRNVKVDLYLIFETALLSGEPHBBQ	672
Frankliniella oc	VEDIRKKEAENRNVKVDLYLIFETALLSGEPHBBQ	672
Trialeurodes vap	VEDIRKKEAENRNVKVDLYLIFETALLSGEPHBBQ	672
Pteromalus pupar	VEDIRKKEAENRNVKVDLYLIFETALLSGEPHBBQ	666
Tetranychus cina	VEDIRKKEAENRNVKVDLYLIFETALLSGEPHBBQ	672
Consensus	ir k ea enr nvkdly lifetall sgf l p	
Panonychus citri	VHARIRRMIRKLGIGIDEVPEKDDYKVDENPFCATA	712
Frankliniella oc	VHARIRRMIRKLGIGIDEVPEKDDYKVDENPFCATA	712
Trialeurodes vap	VHARIRRMIRKLGIGIDEVPEKDDYKVDENPFCATA	712
Pteromalus pupar	VHARIRRMIRKLGIGIDEVPEKDDYKVDENPFCATA	705
Tetranychus cina	VHARIRRMIRKLGIGIDEVPEKDDYKVDENPFCATA	712
Consensus	ri zmiklgigid e v p e k d d y k v d e n p f c a t a	
Panonychus citri	DA SRMEEV	721
Frankliniella oc	DA SRMEEV	721
Trialeurodes vap	DA SRMEEV	721
Pteromalus pupar	DA SRMEEV	714
Tetranychus cina	DA SRMEEV	721
Consensus	da srmeev	

Fig. 2. Alignment of HSP90 amino acid sequence of *P. citri* McGregor and its homologous amino acid sequences from other species (*Frankliniella occidentalis*, *Trialeurodes vaporariorum*, *Pteromalus puparum*, *Tetranychus cinabarinus*).

**Table 1.** Toxicity of abamectin to *Panonychus citri* adult females.

Toxicity regression equation	R	$\chi^2$	LC <sub>25</sub> (CI <sub>95</sub> )* (mg/L)	LC <sub>50</sub> (CI <sub>95</sub> ) (mg/L)	LC <sub>75</sub> (CI <sub>95</sub> ) (mg/L)
Y = 6.8926 + 1.5233X	0.9987	0.5075	0.0206 (0.0148–0.0269)	0.0572 (0.0464–0.0686)	0.1586 (0.1339–0.1906)

\*CI<sub>95</sub>, 95% confidence interval.

a short time, but then gradually decreased to levels comparable with those in controls (Fig. 3).

#### EFFECTS OF ABAMECTIN CONCENTRATION AT 25 °C ON THE RELATIVE EXPRESSION OF HSP90 IN *P. CITRI*


After treatment at 25 °C with various concentrations of abamectin, the relative expression levels of HSP90 in adult female *P. citri* decreased within a short time, then fluctuated, and subsequently were gradually restored to levels comparable with those in controls (Fig. 4).

#### EFFECTS OF ABAMECTIN CONCENTRATION AT 32 °C ON THE RELATIVE EXPRESSION OF HSP90 IN *P. CITRI*


With various concentrations of abamectin at 32 °C, the relative expression levels of HSP90 in adult female *P. citri* increased within a short time, then fluctuated, and subsequently were gradually restored to levels comparable with those in controls (Fig. 5).

#### EFFECTS OF TEMPERATURE ON THE RELATIVE EXPRESSION OF HSP90 GENE IN *P. CITRI*

The adult female mites were subjected to 1 of 2 temperatures (18 °C or 32 °C) for various lengths of time to determine the effects of temperature on the relative expression of HSP90 gene in *P. citri*. The control group was maintained at 25 °C. At 18 °C, the relative expression




**Fig. 3.** Relative expressional levels of HSP90 mRNA in *P. citri* adult females after treatment with abamectin at LC<sub>25</sub>, LC<sub>50</sub> and LC<sub>75</sub> levels for the indicated times at 18 °C. Data are shown as means of 3 replicates ± SD. The relative expression level of HSP90 mRNA in the control group is equal to 1.0 and is marked with a dashed line. The asterisk above the bar indicates a significance difference between treatment with abamectin and control group (*t*-test, *P* < 0.05).




**Fig. 4.** Relative expressional levels of HSP90 mRNA in *P. citri* adult females after treatment at 25 °C with abamectin concentrations of LC<sub>25</sub>, LC<sub>50</sub> and LC<sub>75</sub> for the indicated times. Data are shown as means of 3 replicates ± SD. The relative expression level of HSP90 mRNA in the control group is equal to 1.0 and is marked with a dashed line. The asterisk above the bar indicates a significance difference between treatment with abamectin and control group (*t*-test, *P* < 0.05).

of HSP90 in *P. citri* increased within a short time, then fluctuated, and subsequently increased above the level of controls. At 32 °C, the relative expression of HSP90 in *P. citri* decreased within a short time, but then increased above the level of controls (Fig. 6).



**Fig. 5.** Relative expressional levels of HSP90 mRNA in *P. citri* adult females after treatment at 32 °C with abamectin concentrations of LC<sub>25</sub>, LC<sub>50</sub> and LC<sub>75</sub> for the indicated times. Data are shown as means of 3 replicates ± SD. The relative expression level of HSP90 mRNA in the control group is equal to 1.0 and is marked with a dashed line. The asterisk above the bar indicates a significance difference between treatment with abamectin and control group (*t*-test, *P* < 0.05).






**Fig. 6.** Relative expressional levels of HSP90 mRNA in *P. citri* adult females subjected to 18 °C or 32 °C for the length of time indicated. Data are shown as means of 3 replicates  $\pm$  SD. The relative expression level of HSP90 mRNA in the control group is equal to 1.0 and is marked with a dashed line. The asterisk above the bar indicates a significance difference between treatment with temperature and control group (*t*-test,  $P < 0.05$ ).

#### DSRNA INTERFERENCE OF HSP90 IN *P. CITRI*

After the treatment with 0.1  $\mu\text{g}/\mu\text{L}$  of dsRNA for 48 h, the relative expression levels of HSP90 in adult female mites were significantly down-regulated compared with those in the control group. With 0.2  $\mu\text{g}/\mu\text{L}$  of dsRNA, the relative expression of HSP90 in *P. citri* first decreased but then increased 5.25 fold after 48 h ( $P < 0.05$ ). After treatment with 0.4  $\mu\text{g}/\mu\text{L}$  of dsRNA, the relative expression of HSP90 in *P. citri* fluctuated and then was gradually restored to levels comparable with those of controls (Fig. 7).

#### Discussion

Development of resistance in arthropods is mainly caused by decreased target sensitivity and increased activity of detoxifying



**Fig. 7.** Relative expressional levels of HSP90 mRNA in adult female *P. citri* McGregor after treatment with dsRNA at concentrations equivalent to 0.1  $\mu\text{g}/\mu\text{L}$ , 0.2  $\mu\text{g}/\mu\text{L}$  and 0.4  $\mu\text{g}/\mu\text{L}$ . Data are shown as means of 3 replicates  $\pm$  SD. The relative expression level of HSP90 mRNA in the control group is equal to 1.0 and is marked with a dashed line. The asterisk above the bar indicates a significance difference between treatment with dsRNA and control group (*t*-test,  $P < 0.05$ ).

enzymes (Ran et al. 2009; Liu et al. 2011). Although acaricide resistance in *P. citri* is rampant, there is less research on the elucidation of resistance mechanisms in this mite than on other arthropods, and, thus, the molecular mechanism for resistance of *P. citri* to most acaricides remains unclear (Chen et al. 2009; Niu et al. 2012). A previous study suggested that abamectin resistance of *T. cinnabarinus* was related to an increase in its HSP90 expression (Zhu et al. 2013). In the present study, the full-length sequence of the HSP90 gene from *P. citri* was obtained by cloning, and then the effects of various abamectin concentrations on the HSP90 expression in *P. citri* adult females were analyzed. The results indicated that abamectin treatment at various times induced a marked up-regulation of the HSP90 gene in female adult *P. citri*, suggesting that the resistance of *P. citri* to abamectin may be related to the up-regulation of HSP90 gene.

As a chaperone protein, HSP90 maintains protein stability when stress tends to denature proteins and cause them to aggregate into nonfunctional structures. Thus under high or low temperature stress, HSP90 is overexpressed to maintain the stability of proteins in organisms, and when appropriate environmental conditions are restored, HSP90 expression returns to normal levels. This regulation is likely the reason why HSP90 was temporarily upregulated, as shown in Figs. 3–5, in the present study. The optimal temperature for normal growth of *P. citri* is 25 °C, probably because this temperature is optimal for enzyme activity. Either higher or lower temperatures may influence the regulatory element of the HSP90 gene. Therefore, as shown in Fig. 4, the variation of HSP90 expression in the adult female mites was more robust below 25 °C. Moreover, at the three temperatures, the mortality of the adult female mites increased with the pesticide concentration over time. When the effect of temperature on HSP90 gene expression in *P. citri* was further analyzed, we found that high temperature also induced an increase in the expression of this gene, suggesting that up-regulating the HSP90 gene may also promote resistance to high temperatures in *P. citri*, but these results also indicated the need for further functional analysis.

RNA interference is an important tool for analyzing gene functions (Campbell et al. 2010). Two common methods used for RNA interference in arthropods are injection and feeding (Blandin et al. 2002; Soares et al. 2005; Turner et al. 2006; Walshe et al. 2009; Huvenne & Smagghe 2010). Because of the small size of mites, the injection method affords better results than feeding for gene silencing, but mortality is higher with the injection method. By contrast, the feeding method affords a lower mortality rate, but the gene silencing is not ideal (Chen et al. 2010; Liu et al. 2010). There are no previous reports of applying RNAi technology in *P. citri*. In the present study, the *P. citri* mites were fed dsRNA (0.1  $\mu\text{g}/\mu\text{L}$ , 0.2  $\mu\text{g}/\mu\text{L}$  and 0.4  $\mu\text{g}/\mu\text{L}$ ). The results indicated that this method effectively decreased HSP90 gene expression in *P. citri*; however, the decrease was unstable. This instability may have been caused by the feeding pattern of *P. citri* and the short duration of feeding on dsRNA. Complementary features of other molecular chaperone proteins might have limited the mortality rates observed in *P. citri* adult females caused by the various concentrations of the dsRNA. Additional systematic studies will be required to perfect the RNAi methodology for use in *P. citri*.

Among the current methodologies for functional gene analysis (Cerutti et al. 2011), RNA interference is appropriate for use in *P. citri*. However, the most effective dsRNA concentrations and lengths remain to be determined. In future studies, abamectin-resistant strains will be crossed and selected in an indoor laboratory. The RNAi methodology for functional HSP90 gene analysis will be established, and the difference in the expression of the HSP90 gene in acaricide-resistant and acaricide-sensitive *P. citri* strains at different temperatures will be

compared. The RNAi technology will be applied to analyze both the difference in survival of *P. citri* at high temperatures and the variation in sensitivity of abamectin-resistant strains after silencing of the HSP90 gene. Thus, the results in the present study that examined the effects of the HSP90 gene on the resistance of *P. citri* to high temperatures and chemicals may be confirmed. In summary, the results of the present study provide the molecular basis for future research on analyzing the relationship between the HSP90 gene and resistance of *P. citri* to high temperatures and abamectin.

## Acknowledgments

This research was supported by the National Science and Technology Support Programme (2012BAD19B06), the National Sparking Plan Project (2014GA811008), the Chongqing Scientific Research Project (cstc2014fazktjcsf80031) and the Special Foundation of Chongqing Key Laboratory of Citrus. We also thank Wen-Hua Tian and Yun-Fei Zhang for their technical assistance in this study.

## References Cited

- Akerfelt M, Morimoto RI, Sistonen L. 2011. Heat shock factors: integrators of cell stress, development and lifespan. *Nature Reviews Molecular Cell Biology* 11: 545-555.
- Blandin S, Moita LF, Köcher T, Wilm M, Kafatos FC, Levashina E. A. 2002. Reverse genetics in the mosquito *Anopheles gambiae*: Targeted disruption of the Defensin gene. *Embo Reports* 3(9): 852-856.
- Busvine JR. 1980. Recommended methods for measurement of resistance to pesticides. *FAO Plant Production and Protection Paper* 21: 49-54.
- Campbell EM, Budge GE, Bowman AS. 2010. Gene-knockdown in the honey bee mite *Varroa destructor* by a non-invasive approach: Studies on a glutathione S-transferase. *Parasites and Vectors* 3: 73-82.
- Cerutti H, Ma XR, Msanne J, Repas T. 2011. RNA-mediated silencing in algae: Biological roles and tools for analysis of gene function. *Eukaryotic Cell* 10(9): 1164-1172.
- Beijing Westwood Booker Network Technology Co., Ltd. 2010. Chemical Book. 2010. [http://www.chemicalbook.com/NewsInfo\\_255.htm](http://www.chemicalbook.com/NewsInfo_255.htm)
- Chen J, Zhang D, Yao Q, Zhang J, Dong X, Tian H, Chen J, Zhang W. 2010. Feeding-based RNA interference of a *trehalose phosphate synthase* gene in the brown planthopper, *Nilaparvata lugens*. *Insect Molecular Biology* 19(6): 777-786.
- Chen ZY, Ran C, Zhang L, Dou W, Wang JJ. 2009. Susceptibility and esterase activity in citrus red mite *Panonychus citri* (McGregor) (Acari: Tetranychidae) after selection with phoxim. *International Journal of Acarology* 35(1): 33-40.
- Feder ME, Hofmann GE. 1999. Heat shock proteins, molecular chaperons, and the stress response: Evolutionary and ecological physiology. *Annual Review of Physiology* 61: 243-282.
- Feng HZ, Wang L, Liu YH, He L, Li M, Lu WC, Xue CH. 2010. Molecular characterization and expression of a heat shock protein gene (HSP90) from the carmine spider mite, *Tetranychus cinnabarinus* (Boisduval). *Journal of Insect Science* 10(112): 1-14.
- Fire A, Xu SQ, Montgomery MK, Kostas SA, Driver SE, Mello CC. 1998. Potent and specific genetic interference by double-stranded RNA in *Caenorhabditis elegans*. *Nature* 391: 806-811.
- Furuhashi K. 1994. Development of acaricide resistance in the citrus red mite, *Panonychus citri* (McGregor), in Shizuoka Prefecture. *Proceedings of the Kanto-Tosan Plant Protection Society* 41: 267-269.
- Hu JF, Wang CF, Wang J, You Y, Chen F. 2010. Monitoring of resistance to spirotolifen and 5 other acaricides in *Panonychus citri* collected from Chinese citrus orchards. *Pest Management Science* 66(9): 1025-1030.
- Huvenne H, Smagghe G. 2010. Mechanisms of dsRNA uptake in insects and potential of RNAi for pest control: A review. *Journal of Insect Physiology* 56(3): 227-235.
- Institute for Control of Agrochemical, Ministry of Agriculture of the People's Republic of China. 2014. China Pesticide Information Network. 2014. <http://www.chinapesticide.gov.cn/index.html>
- Liu B, Jiang GF, Zhang YF, Li JL, Li XJ, Yue JS, Chen F, Liu HQ, Li HJ, Zhu SP, Wang JJ, Ran C. 2011. Analysis of transcriptome differences between resistant and susceptible strains of the citrus red mite, *Panonychus citri* (Acari: Tetranychidae). *PLoS ONE* 6: 28516.
- Liu SH, Ding ZP, Zhang CW, Yang BJ, Liu ZW. 2010. Gene knockdown by intro-thoracic injection of double-stranded RNA in the brown planthopper, *Nilaparvata lugens*. *Insect Biochemistry and Molecular Biology* 40(9): 666-671.
- Livak KJ, Schmittgen TD. 2001. Analysis of relative gene expression data using real-time quantitative PCR and the 2<sup>-ΔΔCT</sup> method. *Methods* 25: 402-408.
- Li XX, Zhang MY, Zhang HY. 2011. RNA interference of four genes in adult *Bactrocera dorsalis* by feeding their dsRNA. *PLoS ONE* 6: 17788.
- Li YJ, Wang ZY, Zhang GH, Liu H. 2014. Effects of different temperatures on the growth and development of *Eotetranychus kankitus* (Ehara). *Acta Ecologica Sinica* 34(4): 862-868.
- Niu JZ, Dou W, Ding TB, Yang LH, Shen GM, Wang JJ. 2012. Evaluation of suitable reference genes for quantitative RT-PCR during development and abiotic stress in *Panonychus citri* (McGregor) (Acari: Tetranychidae). *Molecular Biology Reports* 39(5): 5841-5849.
- Niu JZ, Dou W, Wang BJ, Zhang GN, Zhang R, Yin Y, Wang JJ. 2012. Purification and partial characterization of glutathione S-transferases from three field populations of *Panonychus citri* (McGregor) (Acari: Tetranychidae). *Experimental and Applied Acarology* 56(2): 99-111.
- Ouyang YL, Montez GH, Liu L, Grafton-Cardwell EE. 2012. Spirotolifen and spirotetramat bioassays for monitoring resistance in citrus red mite, *Panonychus citri* (Acari: Tetranychidae). *Pest Management Science* 68(5): 781-787.
- Padmini F. 2010. Physiological adaptations of stressed fish to polluted environments: roles of heat shock proteins. *Reviews of Environmental Contamination and Toxicology* 206: 1-27.
- Ran C, Chen Y, Wang JJ. 2009. Susceptibility and carboxylesterase activity of five field populations of *Panonychus citri* (McGregor) (Acari: Tetranychidae) to four acaricides. *International Journal of Acarology* 35(2): 115-121.
- Soares CAG, Lima CMR, Dolan MC, Piesman J, Beard CB, Zeidner NS. 2005. Capillary feeding of specific dsRNA induces silencing of the *isac* gene in nymphal *Ixodes scapularis* ticks. *Insect Molecular Biology* 14(4): 443-452.
- Staheli JP, Ryan JT, Bruce AG, Boyce R, Rose TM. 2009. Consensus-degenerate hybrid oligonucleotide primers (CODEHOPS) for the detection of novel viruses in non-human primates. *Methods* 49(1): 32-41.
- Turner CT, Davy MW, MacDiarmid RM, Plummer KM, Birch NP, Newcomb RD. 2006. RNA interference in the light brown apple moth, *Epiphyas postvittana* (Walker) induced by double-stranded RNA feeding. *Insect Molecular Biology* 15(3): 383-391.
- Walshe DP, Lehane SM, Lehane MJ, Haines LR. 2009. Prolonged gene knockdown in the tsetse fly *Glossina* by feeding double stranded RNA. *Insect Molecular Biology* 18(1): 11-19.
- Xia WK, Ding TB, Niu JZ, Liao CY, Zhong R, Yang WJ, Liu B, Dou W, Wang JJ. 2014. Exposure to diflubenzuron results in an up-regulation of a chitin synthase 1 gene in citrus red mite, *Panonychus citri* (Acari: Tetranychidae). *International Journal of Molecular Science* 15(3): 3711-3728.
- Zhang K, Niu JZ, Ding TB, Dou W, Wang JJ. 2013. Molecular characterization of two carboxylesterase genes of the citrus red mites (Acari: Tetranychidae). *Insect Biochemistry and Physiology* 82(4): 213-226.
- Zhu JY, Wu, GX, Ye GY, Hu C. 2013. Heat shock protein genes (HSP20, HSP75 and HSP90) from *Pieris rapae*: Molecular cloning and transcription in response to parasitization by *Pteromalus puparum*. *Insect Science* 20: 183-193.