



## **Corrigenda: Body Size, Sexual Dimorphism and Ecological Succession in Grasshoppers Journal of Orthoptera Research 17(2): 177–181**

Authors: Picaud, F., and Petit, D.P.

Source: Journal of Orthoptera Research, 18(1) : 137

Published By: Orthopterists' Society

URL: <https://doi.org/10.1665/034.018.0118>

---

BioOne Complete ([complete.BioOne.org](http://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](http://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Corrigenda:


**Body size, sexual dimorphism and ecological succession in grasshoppers**  
**Journal of Orthoptera Research 17(2): 177-181**

F. PICAUD AND D.P. PETIT

(FP) Société Entomologique du Limousin, av. Baudin, 87000 Limoges, France. Email: picaud@educagri.fr

(DP) UMR INRA 1061, Faculté des sciences et techniques, Université de Limoges, 123, av. A. Thomas, F-87060 Limoges cedex, France.  
Email: dpetit@unilim.fr

Corrected Fig. 2.



**Fig. 2.** Variations of sexual dimorphism associated with tegmina length (A) and hind wing length (B) according to succession stage (p values are given for ANOVAs, N=12).