

The effect of the insecticide dichlorvos on esterase activity extracted from the psocids, *Liposcelis bostrychophila* and *L. entomophila*

Authors: Wang, Jin-Jun, Cheng, Wei-Xia, Ding, Wei, and Zhao, Zhi-Mo

Source: Journal of Insect Science, 4(23) : 1-5

Published By: Entomological Society of America

URL: <https://doi.org/10.1673/031.004.2301>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.



The effect of the insecticide dichlorvos on esterase activity extracted from the psocids, *Liposcelis bostrychophila* and *L. entomophila*

Jin-Jun Wang, Wei-Xia Cheng, Wei Ding, Zhi-Mo Zhao

College of Plant Protection, Southwest Agricultural University, Chongqing 400716, People's Republic of China

jjwang7008@tom.com

epcl@swau.cq.cn

Received 17 January 2004, Accepted 10 May 2004, Published 14 July 2004

Abstract

The inhibition kinetics of dichlorvos on carboxylesterase and acetylcholinesterase (AChE) activity extracted from *Liposcelis bostrychophila* and *L. entomophila* (Psocoptera: Liposcelididae) were compared. The results showed that *L. entomophila* had significantly greater specific activity of carboxylesterase than *L. bostrychophila* (0.045 versus 0.012 $\mu\text{moles/mg/min}$). Moreover, the carboxylesterase of *L. entomophila* showed higher affinity (i.e. lower K_m value) to the substrate 1-naphthyl acetate than *L. bostrychophila* (0.29 versus 0.67 mM). The specific activity and affinity of AChE of the two species were not significantly different. The carboxylesterase of *L. bostrychophila* was more sensitive to the insecticide dichlorvos than that of *L. entomophila*. The I_{50} s values of dichlorvos to carboxylesterase for *L. bostrychophila* and *L. entomophila* were 1.43 and 3.28 μM , respectively, and to AChE were 324 and 612 nM, respectively. Inhibition kinetics revealed that AChE from *L. bostrychophila* was 5.8-fold more sensitive to inhibition than AChE from *L. entomophila*.

Keywords: psocid; susceptibility; carboxylesterase, AChE, DDVP

Abbreviation:

1-NA	1-naphthyl acetate
AChE	Acetylcholinesterase
ATChI	Acetylthiocholine iodide
E3	carboxylesterase isozyme (E.C.3.1.1)

Introduction

The psocids, *Liposcelis bostrychophila* Badonnel and *L. entomophila* (Enderlein) are worldwide and commonly found in various processed and unprocessed dry foods in households, granaries, and warehouses (Turner, 1994). Outbreaks of *L. bostrychophila*, and *L. entomophila* have been reported in humid tropical countries such as Indonesia, Malaysia, Singapore, The Philippines, Thailand, The People's Republic of China and India (Wang et al., 1999). Information on the management of psocid pests, however, is very limited (Rees, 1994). Routine fumigations of warehouses and storage facilities with methyl bromide have failed to control these pests (Ho and Winks, 1995). In addition, the rapid development of resistance to chemical and physical treatments by the psocids has also been reported (Santoso et al., 1996; Wang and Zhao, 2003).

Metabolic resistance to organophosphorus insecticides has been associated with changes in the activity of carboxylesterases in many insect species (Devonshire and Field 1991). In two well-studied cases in which resistance to organophosphorus insecticides is associated with an increase in carboxylesterase activity,

sequestration and slow turnover of the phosphate by an over-expressed esterase are responsible for resistance (Devonshire 1977; Karunaratne et al., 1993; Ketterman et al., 1993; Jayawardena et al., 1994). On the other hand, in some insects, resistance to organophosphorus insecticides is associated with the decrease in carboxylesterase activity, such as in the flies, *Musca domestica*, *Lucilia cuprina*, and *Drosophila melanogaster*, where strains resistant to organophosphorus insecticides have high ali-esterase, low organophosphorus hydrolase, and intermediate malathion carboxylesterase (MCE) activities (Campbell et al., 1997).

Acetylcholinesterase (AChE) is a key enzyme that terminates nerve impulses by catalyzing the hydrolysis of the neurotransmitter acetylcholine in the nervous system. Organophosphorous insecticides, such as diazinon, target AChE and irreversibly inhibit the enzyme by phosphorylating a serine hydroxyl group within the enzyme active site. In China, dichlorvos is commonly used to control the insect pests not only in the field but also in stored products. As in the case of other organophosphate insecticides, dichlorvos exerts its effects by inhibiting esterases, especially AChE. In addition, the production of different forms of carboxylesterases is also reported to be the cause of dichlorvos resistance in several insect species.

As little is known about the mechanisms of insecticide resistance in psocids insects, information on the pesticide biochemistry of *Liposcelis* esterases will certainly prove valuable in formulating strategies in the control of these rapidly proliferating pests (Leong and Ho, 1995). This study was initiated to understand the kinetics of carboxylesterase and AChE inhibition by dichlorvos of two *Liposcelis* species. This is an initial step in elucidating the molecular basis of resistance to organophosphorus insecticides in these species.

Materials and methods

Insects

Stock colonies of *L. bostrychophila* and *L. entomophila* were started with nymphs collected from a wheat warehouse in Chongqing, the People's Republic of China in 1990. The colonies were maintained on an artificial diet consisting of whole wheat flour, skim milk and yeast powder (10:1:1) in a room maintained at $28 \pm 1^\circ \text{C}$ and a scotoperiod of 24 h. Cultures were set up in glass bottles (250 ml) with a nylon screen cover and kept in desiccators (5 liter), in which the humidity was controlled with saturated NaCl solution at 75-80%. After several generations, insects from the stock colonies were used for the tests. All experiments were conducted under the conditions described above with 2- to 5-day old female adults.

Chemicals and insecticide

Acetylthiocholine iodide (ATChI, Sigma, www.sigmaaldrich.com), 5,5'-dithiobis-2-nitrobenzoic acid (DTNB, Sigma, www.sigmaaldrich.com), eserine (Sigma), and 1-naphthyl acetate (1-NA) and other biochemical reagents were of reagent grade or better. The insecticide used was 80% dichlorvos (Shalungda Ltd., Changsha, China).

Bioassay

The efficacy of dichlorvos against the two different liposcelids was determined using the small glass tubes (~6mm x 40mm). Various concentrations of dichlorvos were tested until a satisfactory range (10% - 90% mortality) was ascertained. Six concentrations were used in the final analysis. All the concentrations were diluted with acetone. 30µl of insecticide was pipetted onto the inside of the tubes homogeneously and allowed to dry for 30 min before exposing the insects to it.

Each dichlorvos bioassay consisted of 100 adults per concentration and six concentrations (0.36-367 µg/m²). Control groups received acetone alone. Mortality was assessed after 24 h. Psocids that did not move after stimulation from a camel's hair brush were scored as dead. All tests were run at 25°C and replicated at least three times on three different days. Mortality data were corrected with Abbott's (1925) formula and analyzed by probit analysis (Raymond, 1985) to determine the lethal concentrations (LC₅₀).

Enzyme preparation

For carboxylesterase, fifty female adults were ground in 3 ml of ice-cold 0.04 M, pH 7.0 sodium phosphate buffer in a tissue grinder. The crude homogenates were centrifuged at 10,000g for 15 min at 4°C . For AChE, fifty female adults were prepared in 3 ml

of ice-cold 0.1 M, pH 8.0 phosphate buffer containing 0.1% Triton X-100. The crude homogenates were centrifuged at 20,000g at 4°C for 60 min. The resulting supernatants were used as the enzyme sources.

Carboxylesterase assay

Van Asperen's (1962) method was adapted for the determination of esterase activity. The general buffer was 0.04 M, pH 7.0 phosphate buffer. 1-NA ($3 \times 10^{-4} \text{M}$) was used as substrate. In determining the Michaelis constants for 1-NA, the substrate concentrations of 1.5, 3, 6, 15, 30 and 60 mM were made up in phosphate buffer (0.04 M, pH 7.0). The mixtures were incubated at 37°C for 30 min in a water-bath. The reaction was terminated by adding 1 ml of Fast Blue B salt- sodium dodecylsulphate solution. Absorbance was read in the spectrophotometer after 30 min at 600 nm. The kinetic parameters (K_m and V_{max}) were determined graphically by Lineweaver-Burk plots (Wilkinson, 1961).

The *in vitro* inhibition of carboxylesterase activity was ascertained using $3 \times 10^{-4} \text{M}$ 1-NA as substrate. Concentrations of dichlorvos ranging from $1 \times 10^{-9} \text{M}$ to $1 \times 10^{-3} \text{M}$ in phosphate buffer were tested. Inhibitor (0.1 ml) was added to 5 ml of substrate and the reaction initiated by adding 0.1 ml of enzyme. The relative potency of the inhibitors was investigated by examining their I_{50} values.

Acetylcholinesterase assay

AChE activity and its inhibition by dichlorvos were determined according to the method of Ellman et al. (1961) using ATChI as substrate. Briefly, the reaction mixture consisting of ATChI (1.5 mM), DTNB (1 mM) and enzyme preparation (200 µl) was prepared in a final volume of 2.4 ml with phosphate buffer (0.1 M, pH 8.0). The inhibition of dichlorvos was determined by adding various concentrations (from $1 \times 10^{-9} \text{M}$ to $1 \times 10^{-6} \text{M}$ in phosphate buffer) inhibitor (0.1 ml) to the substrate. Absorbance was recorded at 412 nm after 30 min water-bath.

Values of K_m and V_{max} of AChE were determined at 30°C , with 5 ATChI concentrations (ranging from 15 µM to 6.0 mM). The changes of absorbance were observed at 412 nm for 5 min.

The bimolecular rate constants ($k_i = kp/K_a$), phosphorylation constant kp and affinity constant K_a , with dichlorvos as inhibitor were determined by pre-incubation of the supernatants with varying inhibitor concentrations. Progressive inhibition of AChE activity over time was continuously recorded for 5 min. The activity of AChE at each 30 second interval was measured for fitting the inhibition curve. The k_i value was calculated according to the method of Main (1964). The k_i was determined from the gradient of the linear regression:

$$\frac{1}{i} = \frac{\Delta t}{2.303 \Delta \log v} k_i - \frac{1}{K_a}$$

where i is the initial concentration of inhibitor. Values of ($\Delta t / 2.303 \Delta \log v$) were obtained from a plot of $\log v$ against t at constant i . The slope is k_i , the intercept on the ($1/i$) axis is ($-1/K_a$), and the intercept on the ($\Delta t / 2.303 \Delta \log v$) axis is ($1/kp$).

Assays of protein contents

Protein contents of the enzyme homogenate were

determined according to the method of Bradford (1976) using bovine serum albumin as standard. The measurement was performed with the spectrophotometer at 595 nm.

Results

Bioassays

The exposure concentrations of dichlorvos required obtaining LC₅₀ values for *L. bostrychophila* and *L. entomophila* adults are summarized in Tables 1. The data show that *L. bostrychophila* is more tolerant of dichlorvos than *L. entomophila* based on LC₅₀ values. However, the difference of tolerance between two species is not significant considering the 95% confidence limit ($P > 0.05$).

Activities of carboxylesterase

There was a strong linear relationship between homogenate concentrations and carboxylesterase activity for both *L. bostrychophila* ($R^2 = 0.998$) and *L. entomophila* ($R^2 = 0.999$). Homogenate concentrations of 5 and 1.67 insects equivalents per assay were in the middle portion of the linear regression and were used throughout this study.

L. bostrychophila and *L. entomophila* differed significantly in the amount of protein per individual. It averaged 37.4 and 66.1µg/insect for *L. bostrychophila* and *L. entomophila*, respectively (Table 2).

Carboxylesterase from *L. entomophila* showed a significantly higher affinity (i.e. lower *Km* value) to the substrate 1-NA than that of *L. bostrychophila* ($P < 0.05$). In contrast, the catalytic activity of 1-NA towards carboxylesterase in *L. entomophila* was higher (i.e. higher *Vmax* value) than that in *L. bostrychophila* (Table 4). The higher activity of *L. entomophila* esterase towards 1-NA than that of *L. bostrychophila* is observed (Table 2).

The relative susceptibility of the esterase from the two liposcelids to dichlorvos is shown in Fig 1A. Based on the I₅₀ values (the concentration required to inhibit 50% of esterase activity), *L. bostrychophila* esterase (1.43 µM) showed higher susceptibility to dichlorvos than *L. entomophila* (3.28 µM). Inhibition kinetics of carboxylesterase indicated that the inhibition type of dichlorvos were competitive for both *L. bostrychophila* and *L. entomophila*.

Activity of AChE

A strong linear relationship between homogenate concentrations and AChE activities for both *L. bostrychophila* ($R^2 = 0.998$) and *L. entomophila* ($R^2 = 0.997$) was obtained. Homogenate concentrations of 4.8 and 6.4 insects per assay were in the middle portion of the linear regression and so 5 insects per assay were used throughout this study.

There was no significant difference ($P > 0.05$) between the affinities (*Km*) of the AChE from either *Liposcelis* spp. The catalytic activities (*Vmax*) of AChE toward ATChI were also similar. This implies that similar degrees of substrate protection was afforded in both species in the inhibition studies.

The activity of AChE per insect in *L. entomophila* was significantly higher than that of *L. bostrychophila*, but the specific activity (nanomoles/min) from both species was similar. The protein content differed in both insects ($P < 0.05$) (Table 3).

The effects of dichlorvos on AChE activity are shown in Fig. 1B. The inhibition of enzyme activity was between 10% and 90%. The efficiencies of the AChE inhibitors were compared based on their I₅₀s. *Liposcelis bostrychophila* AChE was more sensitive to the inhibitory action of dichlorvos than that of *L. entomophila*.

Table 4 shows estimates of the kinetic constants, *Ka*, *kp* and *ki* (*kp/Ka*) for the reaction between AChE and the organophosphorus insecticide, dichlorvos. The *ki* values of *L. bostrychophila* and *L. entomophila* are 0.015 and 0.0026 nM⁻¹min⁻¹, respectively, indicating that dichlorvos is a more potent inhibitor of the AChE for *L. bostrychophila* than for *L. entomophila*, largely due to a 4.16-fold lower *Ka*. *kp* did not differ significantly between the two pests. The susceptibility tendency is similar with the inhibition results of the specific activity for esterase.

Discussion

In the survey of possible interactions between insecticides and carboxylesterase from *Aphis gossypii*, Owusu (1996) found significant inhibition of the enzyme by a number of organophosphates

Table 1. The toxicity of DDVP to *Liposcelis bostrychophila* and *L. entomophila*.

Insects	Slope±SE	LC50 (µg/m2) (95%CLa)	2b
<i>L. bostrychophila</i>	1.1±0.2	54.4 (73.1~35.7)	2.7
<i>L. entomophila</i>	0.9±0.1	45.3 (62.0~28.7)	2.4

a. 95% confidence limit. LC50 is considered significantly different when the 95% CI fail to overlap.

b. Chi-square goodness-of-fit test.

Table 2. Carboxylesterase activity in *Liposcelis bostrychophila* and *L. entomophila*.

Insects	Protein (µg / insect)	nmoles / min/ insect	µmoles / min/mg
<i>L. bostrychophila</i>	37.4±0.3a	0.47±0.04a	0.012±0.005a
<i>L. entomophila</i>	66.1±0.6b	2.99±0.05b	0.045±0.006b
Ratio	1.8	6.4	3.8

Within the same row, means followed by the different letters are significantly different ($P < 0.05$).

Table 3. AChE activity in *Liposcelis bostrychophila* and *L. entomophila*.

Insects	Protein (µg/insect)	nanomoles/min
<i>L. bostrychophila</i>	25. 2±0.7a	24.5±0.5a
<i>L. entomophila</i>	35.9±0.8b	25.2±0.7a

Within the same row, means followed by the different letters are significantly different ($P < 0.05$).

Table 4. Affinity constant (*Ka*), phosphorylation rate constant (*kp*) and bimolecular rate constant (*ki*) of *L. bostrychophila* and *L. entomophila*.

Insecticide	Insects	Ki (nM ⁻¹ min ⁻¹)	Ka (nM)	kp (min ⁻¹)
DDVP				
	<i>L. bostrychophila</i>	0.015±0.002	1.80±0.36	0.027±0.009
	<i>L. entomophila</i>	0.0026±0.0004	7.48±1.34	0.019±0.006




Figure 1. Percent inhibition of carboxylesterase activity (A) and AChE (B) in *Liposcelis bostrychophila* (solid line and dots) and *L. entomophila* (dotted line and circles) in relation to the applied dichlorvos concentrations.

and carbamates. Theoretically, the kinetics involved in such inhibition trends based on a Michaelis intermediate reaction mechanism (Aldridge and Reiner, 1972), in which an enzyme and an inhibitor combine to form enzyme-inhibitor complex. The production of different isozymes of carboxylesterase is reported to be the cause of dichlorvos resistance in the cotton aphid, *A. gossipii* (Glover). By isoelectric focusing technique O'Brien et al. (1992) also showed the involvement of different bands in organophosphate resistance of *A. gossipii* on cotton in the United States. However, the role of such isozymes in resistance of this aphid has not yet been clarified. In the green peach aphid, *Myzus persicae* (Sulzer), insecticide hydrolysis by an E4 isozyme in resistant aphids is the cause of organophosphate resistance (Devonshire and Moores, 1982). For psocids, Leong and Ho (1995) reported that a qualitatively good correlation between the results obtained from the *in vitro* carboxylesterase assays and electrophoretic analysis, showing a

higher esterase sensitivity to inhibition in *L. bostrychophila* than in *L. entomophila*. The similar result was also obtained in the present study. However, based on the bioassay, *L. bostrychophila* is more tolerant to dichlorvos than *L. entomophila*. This may be due to the sensitive esterase of *L. bostrychophila* that may preferentially bind dichlorvos, thereby protecting AChE, resulting in greater tolerance to dichlorvos.

Acetylcholinesterase is of interest because it is the target-site for organophosphate and carbamate insecticides in the central nervous system, and its role in cholinergic synapses is essential for life (Fournier and Muterio, 1994). Based on the values of I_{50} s (the concentration required to inhibit 50% of AChE activity) and k_i (bimolecular rate constants for AChE inhibition), *L. bostrychophila*'s AChE was more sensitive to the inhibitory action of dichlorvos than that of *L. entomophila*. In contrast, Leong and Ho (1995) reported that dichlorvos is a more potent inhibitor of AChE in *L. entomophila* than in *L. bostrychophila*. They also found that *L. entomophila* is more tolerant of dichlorvos than *L. bostrychophila* based on bioassay data. The difference might be due to the different geographic population used. Price (1988) pointed out that the sensitivity of AChE does not necessarily mean that insects will be susceptible to the particular chemical when it is used as an insecticide as toxicity of a particular compound depends on many factors including cuticle penetration and metabolic processes (Matsumura, 1985; Hassall, 1990). This suggests that the higher sensitivity of *L. bostrychophila* AChE to inhibition implies that the underlying mechanism of tolerance in insecticide bioassays in this species is largely due to metabolic activities which either detoxify or limit the intoxicating ability of dichlorvos.

The present study has provided some basic information on the esterases of these two psocids that will be useful to understand the mechanisms of insecticide resistance in the psocids. As strains of liposcelid psocids that exhibit varying tolerance to insecticides become available they will be useful for further comparative toxicology and biochemical studies.

Acknowledgements

This research was funded in part by the National Natural Sciences Foundation (39800017) and Fok Ying-Tung Educational Foundation (71022) of P. R. China to Jin-Jun Wang.

References

- Abbott WS. 1925. A method of computing the effectiveness of an insecticide. *Journal of Economic Entomology* 18: 265–267.
- Aldridge W N, Reiner E. 1972. *Enzyme inhibitors as substrates*. vol. 26. North-Holland, Amsterdam.
- Bradford MM. 1976. A rapid and sensitive method for the quantification of microgram quantities of protein utilizing the principle of protein-dry binding. *Analytical Biochemistry* 72: 248–254.
- Campbell PM, Trott JF, Claudianos C, Smyth K, Russell RJ, Oakeshott JG. 1997. Biochemistry of esterases associated with organophosphate resistance in *Lucilia cuprina* with comparisons to putative orthologues in other diptera. *Biochemical Genetics* 35: 17–40.

- Devonshire AL. 1977. The properties of a carboxylesterase from the peach-potato aphid, *Myzus persicae* (Sulz.), and its role in conferring insecticide resistance. *Biochemical Journal* 167: 675-683.
- Devonshire AL, Moores GD. 1982. A carboxylesterase with broad substrate specificity causes organophosphorus, carbamate and pyrethroid resistance in peach-potato aphid (*Myzus persicae*). *Pesticide Biochemistry and Physiology* 18: 235-246.
- Devonshire AL, Field LM. 1991. Gene amplification and insecticide resistance. *Annual Review of Entomology* 36: 1-23.
- Ellman GL, Courtney KD, Andres V, Featherstone RM. 1961. A new and rapid colorimetric determination of acetylcholinesterases activity. *Biochemical Pharmacology* 7: 88-95.
- Fournier D, Mutero A. 1994. Modification of acetylcholinesterase as a mechanisms of resistance to insecticides. *Comparative Biochemistry and Physiology* 108C: 19-31.
- Hassall KA. 1990. *The biochemistry and use of pesticides. Structure, metabolism, mode of action and use in crop protection*. 2nd ed. 536pp, MacMillan, New York.
- Ho SH, Winks RG. 1995. The response of *Liposcelis bostrychophila* Badonnel and *L. entomophila* (Enderlein) (Psocoptera) to phosphine. *Journal of Stored Products Research* 31: 191-197.
- Jayawardena KGI, Karunaratne SHPP, Kettermann AJ, Hemingway J. 1994. Determination of the role of elevated B₂ esterase in insecticide resistance in *Culex quinquefasciatus* (Diptera: Culicidae) from studies on the purified enzyme. *Bulletin of Entomological Research* 84: 39-44.
- Karunaratne SHPP, Jayawardena KGI, Hemingway J, Kettermann AJ. 1993. Characterisation of a B-type esterase involved in insecticide resistance from the mosquito *Culex quinquefasciatus*. *Biochemical Journal* 294: 575-579.
- Kettermann AJ, Karunaratne SHPP, Jayawardena KGI. 1993. Qualitative differences between populations of *Culex quinquefasciatus* in both the esterases A₂ and B₂ which are involved in insecticide resistance. *Pesticide Biochemistry and Physiology* 47: 142-148.
- Leong ECW, Ho SH 1995. *In vitro* inhibition of esterase activity in *Liposcelis bostrychophila* Bad. and *L. entomophila* (End.) (Psocoptera: Liposcelididae). *Comparative Biochemistry and Physiology* 110B: 121-130.
- Main AR. 1964. Affinity and phosphorylation constants for the inhibition of esterases by organophosphates. *Science* 803: 324-332.
- Matsumura F. 1985. *Toxicology of Insecticides*. 2nd ed., 598pp. Plenum Press, New York.
- O'Brien PJ, Abdel-Aal YA, Ottea JA, Graves JB. 1992. Relationship of insecticide resistance to carboxylesterase in *Aphis gossypii* Glover (Homoptera: Aphididae) from mid south cotton. *Journal of Economic Entomology* 85: 651-657.
- Owusu EO, Horiike M, Hirano C. 1996. Polyacrylamide gel electrophoretic assessments of esterase in cotton aphid (Homoptera: Aphididae) resistance to dichlorvos. *Journal of Economic Entomology* 89: 302-306.
- Price NR. 1988. Insecticide-insensitive acetylcholinesterase from a laboratory selected and field strain of housefly (*Musca domestica*). *Comparative Biochemistry and Physiology* 90C: 221-224.
- Raymond M. 1985. Presentation d'un programme basic d'analyse log-probit pour micro-ordinateur. Cahiers ORSTOM: *Serie Entomologie Medicale et Parasitologie* 23: 117-121.
- Rees DP. 1994. *Tiny insects become big problem*. Stored Grain Australia, June 1994, 1 and 5.
- Santoso T, Dharmaputra OS, Halid H, Hodges RJ. 1996. Pest management of psocids in milled rice stores in the humid tropics. *International Journal of Pest Management* 42: 189-197.
- Turner BD. 1994. *Liposcelis bostrychophila* (Psocoptera; Liposcelididae), a stored food pest in the UK. *International Journal of Pest Management* 40: 179-190.
- Van Asperen K. 1962. A study of house fly esterase by means of a sensitive colorimetric method. *Journal of Insect Physiology* 8: 401-416.
- Wang JJ, Zhao ZM, Li LS. 1999. Induced tolerance of the psocid, *Liposcelis bostrychophila* (Psocoptera: Liposcelididae) to controlled atmosphere. *International Journal of Pest Management* 45: 75-79.
- Wang JJ, Zhao ZM. 2003. Accumulation and utilization of triacylglycerol and polysaccharides in *Liposcelis bostrychophila* (Psocoptera, Liposcelididae) selected for resistance to carbon dioxide. *Journal of Applied Entomology* 127: 107-117.
- Wilkinson G.N. 1961. Statistical estimation in enzyme kinetics. *Biochemical Journal* 803: 324-332.