



The effect of development time on the fitness of female *Trichogramma evanescens*

Authors: Doyon, Josée, and Boivin, Guy

Source: Journal of Insect Science, 5(4) : 1-5

Published By: Entomological Society of America

URL: <https://doi.org/10.1673/031.005.0401>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.



The effect of development time on the fitness of female *Trichogramma evanescens*

Josée Doyon^{1,2} and Guy Boivin¹

¹Horticultural Research and Development Center, Agriculture and AgriFood Canada, St-Jean-sur-Richelieu, Québec, Canada

²Département des Sciences Biologiques, Université du Québec à Montréal, Montréal, Québec, Canada
joseedoyon@iphoria.ca

Received 19 September 2003, Accepted 14 August 2004, Published 16 March 2005

Abstract

Trichogramma are facultative gregarious egg parasitoids that attack a wide range of lepidopterous eggs. Because hosts in which parasitoids develop vary in terms of available food, the progeny produced by parasitoid females vary in size and fitness. While one might expect that the developmental rate and emergence rhythm should be similar for all individuals reared under the same environmental conditions, variations in the duration of development of individuals reared under uniform conditions have been found in several insect species. In the Hymenopteran egg parasitoid *Trichogramma evanescens* Westwood (Hymenoptera: Trichogrammatidae) adults emerge at the beginning of the photoperiod on two consecutive days. Longer development may influence the fitness of adults and have an impact on mating opportunities. Size, longevity and daily and lifetime fecundity were measured for female *T. evanescens* that developed in nine and ten days. We observed a significant relationship between size and both longevity and lifetime fecundity. While early emerging females did not live longer and did not produce progeny with a different sex-ratio than females that emerged later, they were larger and produced more progeny than late females. We conclude that early emerging females have a higher fitness than late emerging females *T. evanescens*.

Keywords: emergence, fecundity, longevity, parasitoid, sex-ratio, size

Introduction

Trichogramma species are facultatively gregarious (Rabinovich 1971), polyphagous egg parasitoids that are often used in inundative biological control programs (Smith 1996) against a wide range of Lepidopterous eggs (Corrigan and Laing 1994). *Trichogramma* spp. are moderately synovigenic (Jervis et al. 2001) and females start to lay their eggs shortly after emergence (Bigler et al. 1987; Chassain and Boulétreau 1991). Hosts in which parasitoids develop are variable resources in terms of available food and, as a result, parasitoid adults vary in size with males being generally smaller than females (van den Assem et al. 1989). In a host of a given size, females grow larger than males and males emerge earlier than females (Charnov et al. 1981).

Several studies have found that the female's fitness increases with its size (King 1987; Boivin and Lagacé 1999) indicating that, for an adult parasitoid, being large has consequences on its reproductive success (van den Assem et al. 1989). While body size has no direct relation to fitness, it may be used as a primary proxy when its relationship with another primary proxy (for example fecundity, longevity or mating ability) is known (Roitberg et al. 2001). The size of a female may influence its longevity, its searching efficiency, its egg load and its capacity to locate hosts. The size of a male may influence its longevity, its capacity to locate females and the number of matings (Godfray 1994). In general, the size of *Trichogramma* species seems to be positively related to performance (McDougall and Mills 1997).

Insect developmental rates and emergence rhythms tend to be similar for all individuals of a population reared under the same environmental conditions (Saunders 1976; Beck 1991). However, variations in the developmental period of cohorts reared under the same conditions have been found in many insects (Shaffer 1983). In the species *Trichogramma evanescens*, emergence occur mostly at the beginning of the photoperiod and are spread over two days (nine and ten days at 24 ± 1 °C) with 89% of the males and 95% of the females emerging on the first day (Doyon and Boivin 2004). Because emergence is concentrated at the beginning of the photoperiod, the duration of immature development varies by a few hours between individuals depending on the time of egg deposition. When a female parasitizes a host late in the day, the immature may not be ready to emerge on the morning of the ninth day and have to delay its emergence until the next day. Individuals that delay emergence may experience either an increase or a decrease in fitness, depending on how host resources are used. An increase in fitness is expected if the supplementary developmental time can be used by the individual to either exploit more fully the host resource or to mature. However, a decrease in fitness is more likely if the individual is merely waiting for emergence while burning reserves to maintain its somatic functions. Individuals emerging later can also be penalized by encountering low-quality mates (Carvalho et al. 1998) or by experiencing lower mating opportunities (Waage and Ng 1984).

Emerging early in the day is certainly an advantage for short-lived diurnal species such as *Trichogramma*. Under laboratory conditions, *T. evanescens* females deposit 56% of their eggs during

the first 24 hours (Boivin and Lagacé 1999) and survival in the field probably does not exceed 24–48 hours. It is, therefore, important to emerge at the beginning of a day when host location is possible. Females that are unable to emerge on the morning of the ninth day are thus facing a trade-off between emerging late in the day and missing oviposition opportunities or delaying emergence to the tenth day and possibly having a lower fitness.

In this paper, we compared the longevity, fecundity (daily and lifetime fecundity) and size of females *T. evanescens* that emerged after nine days of development (early females) versus females that emerged after ten days of development (late females).

Material and Methods

Biological material

The *T. evanescens* strain we used originated from Egypt, and was reared on *Ephestia kuehniella* Zeller (Lepidoptera: Pyralidae) eggs at $25 \pm 1^\circ\text{C}$, $40 \pm 10\%$ RH under a light dark photoperiod of 8L:16D. Individuals of *E. kuehniella* were reared at $25 \pm 2^\circ\text{C}$, $40 \pm 10\%$ RH and 8L:16D. The *E. kuehniella* eggs were less than 24 h old and were cold sterilized at -15°C for 24 h.

Longevity, fecundity and size of female *T. evanescens*

Approximately 100 *T. evanescens* females were placed in contact with about 1000 *E. kuehniella* eggs for 30 min and after which the eggs were incubated at $24 \pm 1^\circ\text{C}$. Under these conditions, female *T. evanescens* emerge on the ninth (early females) and tenth (late females) days after oviposition (Doyon 2004). Early ($n = 29$) and late females ($n = 26$) were individually mated at emergence by a male less than 3 h old and were then placed in a 3.7 ml tube (without food) with approximately 250 eggs of *E. kuehniella* that were changed every 24 h during the life of the female. These eggs were incubated in the same rearing chamber, in 3.7 ml tubes, and at emergence the individuals were counted and the sex-ratio calculated. To determine the longevity of early ($n = 30$) and late females ($n = 30$), we monitored their survival every 4 h (± 15 min) from 6:30 to 22:30. The death of females was assigned to the mid-point between two observations. Because overall body size and hind tibia length are highly correlated in *Trichogramma* species (Bai 1986), and because it is less sensitive to distortion than other body parts (Bai 1986), at the end of the experiment, the hind tibia length of all female *T. evanescens* was measured.

Statistical analyses

Linear regressions were used to obtain the relation between size and longevity, and size and lifetime fecundity. An analysis of covariance with size as co-variable was used to compare the lifetime fecundity of early and late females. The longevity and daily fecundity of early and late females *T. evanescens* were compared with *t*-tests. The Fisher's exact test was used to compare the daily sex-ratio of the progeny of the females.

Results

As expected, a significant relationship was found between size and longevity (Fig. 1) and size and lifetime fecundity (Fig. 2) for both early and late females. The mean longevity of early females




Figure 1. Longevity vs. tibia length of (a) early females (linear regression $R = 0.516$, d.f. = 29, $p < 0.01$). (b) late females (linear regression $R = 0.624$, d.f. = 29, $p < 0.01$).




Figure 2. Lifetime fecundity vs. tibia length of (a) early females (linear regression $R = 0.478$, d.f. = 28, $p < 0.01$). (b) late females (linear regression $R = 0.676$, d.f. = 25, $p < 0.01$).

Table 1. Daily fecundity of early and late female *T. evanescens* reared on *E. kuehniella*.

Day	Daily fecundity of early females (±SD)	Daily fecundity of late females (±SD)	t-test
1	35.10 ± 5.17	29.38 ± 7.19	t= 4.14 df= 53 <i>p</i> = 0.01
2	12.59 ± 3.18	10.80 ± 3.46	t= 2.21 df= 52 <i>p</i> = 0.034
3	11.56 ± 3.98	9.60 ± 3.80	t= 2.35 df= 50 <i>p</i> = 0.025
4	7.41 ± 3.15	5.85 ± 3.01	t= 2.30 df= 45 <i>p</i> = 0.029
5	7.84 ± 4.24	6.21 ± 3.57	t=2.04 df= 31 <i>p</i> = 0.057
6	4.86 ± 2.39	7.60 ± 3.44	t= -1.65 df= 10 <i>p</i> = 0.166
7	4.50 ± 1.17	4.83 ± 2.44	t= -2.05 df= 6 <i>p</i> = 0.922

Table 2. Sex-ratio of early and late females *T. evanescens* reared on *E. kuehniella*.

Source	df	Sum of square	Mean square	F value	Pr > F
Model	2	8157.24	4078.62	18.84	< 0.0001
Error	52	11260.15	216.54		
Corrected total	54	19417.38			
Size	1	7386.93	7386.93	34.11	< 0.0001
Lifetime fecundity	1	770.3	770.3	3.56	0.0649

Table 3. Analysis of variance of the lifetime fecundity of early and late females with size as a co-variable.

Source	df	Sum of square	Mean square	F value	Pr > F
Model	2	8157.24	4078.62	18.84	< 0.0001
Error	52	11260.15	216.54		
Corrected total	54	19417.38			
Size	1	7386.93	7386.93	34.11	< 0.0001
Lifetime fecundity	1	770.3	770.3	3.56	0.0649

(112.45 ± 0.25 h) was not significantly different from the longevity of late females (116.25 ± 0.25 h) (*t* = −0.0164; *df* = 58; *p* = 0.630). The longevity varied from a minimum of 43.5 h to a maximum of 162.5 h.

On the first four days of their life, early females had a higher daily fecundity than late females. From day 5 to day 7, there was no significant difference between the fecundity of early and late females (Table 1). The daily (Table 2) and lifetime sex-ratio of the progeny produced by early and late females did not differ (Fisher’s exact test *p* = 0.4284; *df* = 1; *p* > 0.05). However, the sex-ratio of the progeny of females increased with the age of the female in both treatments (Table 2).

Because fecundity is correlated with size in many parasitoid species, the lifetime fecundity of females was analyzed with size as a co-variable. While the complete model showed highly significant differences between durations of development, the individual analysis indicated that lifetime fecundity showed no significant differences, although fecundity was higher for females with short developmental time (early females = 70.18 eggs, late females = 62.57 eggs). Size was highly significantly correlated with fecundity (Table 3). The mean hind tibia length was 0.160 ± 0.007 mm for early females and 0.154 ± 0.012 mm for late females. The smallest hind tibia length was 0.141 mm for early females and 0.124 mm for late females.

The largest hind tibia length was 0.178 mm for early females and 0.173 (±0.012) mm for late females.

Discussion

In *T. evanescens*, a significant relationship was found between the size of females and both longevity and lifetime fecundity. These data confirm the results of Waage and Ng (1984), van den Assem (1989), Bennett and Hoffmann (1998), Boivin and Lagacé (1999), Sagarra et al. (2001) and Rivero and West (2002) who showed that large females parasitoids live longer and produce more progeny than small females, and therefore are at an advantage because they have a higher fitness. Therefore, if differences in size are observed between early and late females, we can predict that these differences will translate into differences in fitness.

Both longevity and the sex-ratio of progeny oviposited by early and late females did not vary significantly. In parasitic Hymenoptera, progeny sex-ratios are known to vary and to be determined at oviposition by arrhenotoky (Luck et al. 1992) where males develop from unfertilized eggs (haploid) and females from fertilized eggs (diploid) (Flanders 1965). Following mating, female parasitoids store sperm and can manipulate the sex-ratio of their progeny by controlling the access of sperm to the egg during

oviposition (Flanders 1965; Suzuki et al. 1984). The gradual increase in sex-ratio that was observed was due to sperm depletion of the female (Boivin and Lagacé 1999, Boivin et al. 2004). Female *T. evanescens* can store up to 50 sperm in their spermatheca (Damien's unpublished data).

Early females were larger and more fecund than late females and they produced 18.8% more progeny. Because each egg deposited contributes to the reproductive success of females, early females had a higher fitness than late females. Thus, emerging early had a considerable effect on the size of females and their production of eggs. Although the effect of emergence day on fecundity was almost significant, most of that effect was due to the difference in size between early and late females. The late females were probably not ready to emerge at the beginning of the photoperiod on day 9, so they had to wait another 24 hours to emerge as adults. This increase in developmental time as pupae or sub-imago could have used part of their energy reserves which resulted in a smaller size and a lower fecundity. In *Trichogramma* parasitoids, oogenesis starts during the pre-pupal stage (Volkoff and Daumal 1994) and continues after adult emergence (Mills and Kuhlmann 2000). *Trichogramma* species have been reported as proovigenic by several authors (Pak and Oatman 1982; Waage and Ng 1984; Bai et al. 1992) but they have since been described as moderately synovigenic (Jervis et al. 2001). In many species, females emerge with an egg load that accounts for a fraction of their potential fecundity (Houseweart et al. 1983; Smith and Hubbes 1986; Bai and Smith 1993; Kuhlmann and Mills 1999). Our hypothesis was that females would spend some of their resources maintaining their somatic functions while waiting for emergence on the tenth day. However, exhausting resources, probably of their fat reserve, should not have influenced the size of the females when measured as hind tibia length. The results suggest that it was not the longer developmental time that produced small individuals but rather that the smaller individuals took longer to develop.

Superparasitism, which is of common occurrence in *Trichogramma* spp. in the laboratory (Yadav et al. 2001), could also be the cause of the increase in developmental time in late females. The eggs of *E. kuehniella* usually support the development of a single adult *Trichogramma* per egg. However, more than one parasitoid egg may be initially deposited per host (Klomp and Terrink 1978). Larval competition results in the death of some individuals and the survival of others (Corrigan et al. 1995). Superparasitism affects subsequent parasitoid development (Ahmad et al. 2002) and has been suggested as a factor leading to extended emergence periods in several *Trichogramma* species (Parra et al. 1988).

Trichogramma spp. are the most widely used parasitoid agents for the biological control of lepidopterous pests (Waage and Ng 1984). They are particularly used in inundative releases projects (Corrigan and Laing 1994). The negative impact of longer development on the size and fecundity of *Trichogramma* females suggests that the quality of the individuals produced in mass rearing could be improved by maximizing the proportion of the population emerging early.

Acknowledgements

We thank Josiane Vaillancourt for rearing *T. evanescens*, Danielle Thibodeau and Marie-Josée Gauvin for technical assistance

and Shahrokh Khanizadeh for advice on statistics. This is contribution 335/2004.09.01R of the Centre de Recherche et de Développement en Horticulture, Agriculture et Agroalimentaire Canada, Saint-Jean-sur-Richelieu, Québec, Canada.

References

- Ahmad M, Ahmad MJ, Mishra RK, Sheel SK. 2002. Superparasitism by *Trichogramma poliae* in the eggs of *Clostera cupreata* (Lepidoptera: Notodontidae) and its effect on offspring. *Journal of Tropical Forest Science* 14:61–70.
- van den Assem J, van Iersel JJA, Los-den Hartogh RL. 1989. Is being large more important for female than for male parasitic wasps? *Behavior* 108:160–195.
- Bai B. 1986. Host effect on quality attributes of *Trichogramma pretiosum* Riley (Hymenoptera: Trichogrammatidae) and host discrimination by *Copidosoma truncatellum* Dalman (Hymenoptera: Encyrtidae). M. Sc. Thesis, University of California at Riverside, Riverside, California, U.S.A.
- Bai B, Smith SM. 1993. Effect of host availability on reproduction and survival of the parasitoid wasp *Trichogramma minutum*. *Ecological Entomology* 18:279–286.
- Bai B, Luck RF, Forster L, Stephens B, Janssen JAM. 1992. The effect of host size on quality attributes of the egg parasitoid, *Trichogramma pretiosum*. *Entomologia Experimentalis et Applicata* 64:37–48.
- Beck SD. 1991. Thermoperiodism. In: Lee RE, Denlinger DL, editors. *Insects at Low Temperature*, 199–228. New York: Chapman & Hall.
- Bennett DM, Hoffmann AA. 1998. Effects of size and fluctuating asymmetry on field fitness of the parasitoid *Trichogramma carverae* (Hymenoptera: Trichogrammatidae). *Journal of Animal Ecology* 67:580–591.
- Bigler F, Meyer A, Bosshart S. 1987. Quality assessment in *Trichogramma maidis* Pintureau et Voegelé reared from eggs of the factitious hosts *Ephestia kuehniella* Zeller and *Sitotroga cerealella* Olivier. *Journal of Applied Entomology* 104:340–353.
- Boivin G, Lagacé M. 1999. Effet de la taille sur la fitness de *Trichogramma evanescens* (Hymenoptera: Trichogrammatidae). *Annales de la Société Entomologique de France* 35:371–378.
- Boivin G, Jacob S, Damien's D. 2004. Spermatogeny as a life history index in parasitoid wasps. *Oecologia*.
- Carvalho MC, Queiroz PCD, Ruszezyk A. 1998. Protandry and females size-fecundity variation in the tropical butterfly *Brassolis sophorae*. *Oecologia* 116:98–102.
- Charnov EL, Los-den Hartogh RL, Jones TW, Assem JVD. 1981. Sex ratio evolution in a variable environment. *Nature* 289:27–33.
- Chassain C, Boulétreau M. 1991. Genetic variability in quantitative traits of host exploitation in *Trichogramma* (Hymenoptera: Trichogrammatidae). *Genetica* 83:195–202.
- Corrigan JE, Laing JE. 1994. Effects of the rearing host species and the host species attacked on performance by *Trichogramma minutum* Riley (Hymenoptera: Trichogrammatidae). *Biological Control* 23:755–760.
- Corrigan JE, Laing JE, Zubricky JS. 1995. Effects of parasitoid to host ratio and time of day of parasitism on development

- and emergence of *Trichogramma minutum* (Hymenoptera : Trichogrammatidae) parasitizing eggs of *Ephesia kuehniella* (Lepidoptera : Pyralidae). *Annals of the Entomological Society of America* 88: 773–780.
- Doyon J. 2004. L'impact de la protandrie sur la capacité d'accouplement des mâles et l'effet de développement sur la valeur adaptative des femelles chez *Trichogramma evanescens* Westwood (Hymenoptera : Trichogrammatidae), M. Sc. Thesis, Université du Québec à Montréal, Montreal, Canada.
- Doyon J, Boivin G. 2004. Impact of protandry on mating capacity of males in *Trichogramma evanescens* Westwood (Hymenoptera : Trichogrammatidae).
- Flanders SE. 1965. On the sexuality and sex ratios of hymenopterous populations. *American Naturalist* 99:489–494.
- Godfray HCJ. 1994. *Parasitoids: Behavioral and Evolutionary Ecology*. Princeton, New Jersey: Princeton University Press.
- Houseweart MW, Jennings DT, Welty C, Southard SG. 1983. Progeny production by *Trichogramma minutum* (Hymenoptera: Trichogrammatidae) utilizing eggs of *Choristoneura fumiferana* (Lepidoptera: Gelechiidae). *The Canadian Entomologist* 115:1245–1252.
- Jervis MA, Heimpel GE, Ferns PN, Harvey JA, Kidd NA. 2001. Life history strategies in parasitoid wasps: A comparative analysis of ovigeny. *Journal of Animal Ecology* 70:442–458.
- King BH. 1987. Offspring sex ratios in parasitoid wasps. *Quarterly Review of Biology* 62:367–396.
- Klomp H, Teerink BJ. 1978. The elimination of supernumerary larvae of the gregarious egg-parasitoid *Trichogramma embryophagum* (Hymenoptera: Trichogrammatidae) in eggs of the host *Ephesia kuehniella* (Lepidoptera: Pyralidae). *Entomophaga* 23:153–159.
- Kuhlmann U, Mills NJ. 1999. Comparative analysis of the reproductive attributes of three commercially-produced *Trichogramma* species and the influence of parasitoid size. *Biocontrol Science and Technology* 9:335–346.
- Luck RF, Stouthamer R, Nunney L. 1992. Sex determination and sex ratio patterns in parasitic Hymenoptera. In: Wrench DL and Ebbert M, editors. *Evolution and diversity of sex ratio in haplodiploid insects and mites*, 442–476. New York: Chapman & Hall.
- McDougall SJ, Mills NJ. 1997. Dispersal of *Trichogramma platneri* (Hymenoptera: Trichogrammatidae) from point-source releases in an apple orchard in California. *Journal of Applied Entomology* 121:205–209.
- Mills NJ, Kuhlmann U. 2000. The relationship between egg load and fecundity among *Trichogramma* parasitoids. *Ecological Entomology* 25:315–324.
- Pak GA and Oatman ER. 1982. Biology of *Trichogramma brevicapillum*. *Entomologia Experimentalis et Applicata* 32:61–67.
- Parra, JRP, Zucchi RA, Silveira Neto S. 1988. Perspectives of biological control using *Trichogramma* and/or *Trichogrammatoidea* in the state of São Paulo (Brazil). In: *Trichogramma and other egg parasites*, 43:527–540. INRA.
- Rabinovich JE. 1971. Population dynamics of *Telenomus fariai* (Hymenoptera: Scelionidae), a parasite of Chaga's disease vector. Parasite size and vital space. *Revista de Biologia Tropical* 19:109–120.
- Rivero A, West SA. 2002. The physiological costs of being small in parasitic wasp. *Evolutionary Ecology Research* 4:407–420.
- Roitberg BD, Boivin G and Vet L. 2001. Fitness, parasitoids, and biological control: An opinion. *The Canadian Entomologist* 133:429–438.
- Sagarra LA, Vincent C, Stewart RK. 2001. Body size as an indicator of parasitoid quality in male and female *Anagyrus kamali* (Hymenoptera : Encyrtidae). *Bulletin of Entomological Research* 91:363–367.
- Saunders DS. 1976. *Insect Clocks*. New York: Pergamon.
- Shaffer PL. 1983. Prediction of variation in developmental period of insects and mites reared at constant temperatures. *Environmental Entomology* 12:1012–1019.
- Smith SM. 1996. Biological control with *Trichogramma*: Advances, successes, and potential of their use. *Annual Review of Entomology* 41:375–406.
- Smith SM, Hubbes M. 1986. Strains of the egg parasitoids *Trichogramma minutum* Riley. II. Utilization for release against the spruce budworm. *Journal of Applied Entomology* 102:81–83.
- Suzuki Y, Tsuiji H, Sasakawa M. 1984. Sex allocation and the effects of superparasitism on secondary sex ratios in the gregarious parasitoid, *Trichogramma chilonis* (Hymenoptera: Trichogrammatidae). *Animal Behavior* 32:478–484.
- Volkoff AN, Daumal J. 1994. Ovarian cycle in immature and adult stages of *Trichogramma cacoeciae* and *T. brassicae* (Hymenoptera: Trichogrammatidae). *Entomophaga* 39:303–312.
- Waage JK, Ming NGS. 1984. The reproductive strategy of a parasitic wasp I. Optimal progeny and sex allocation in *Trichogramma evanescens*. *Journal of Animal Ecology* 53:401–415.
- Yadav RC, Singh SP, Jalali SK, Rao NS. 2001. Effect of host egg density on parasitism and adult emergence in *Trichogramma chilonis* Ishii (Hymenoptera: Trichogrammatidae) in two systems. *Journal of Biological Control* 15:11–14.