BioOne COMPLETE

Species Richness and Host Associations of Lepidoptera-Attacking Tachinidae in the Northeast Ecuadorian Andes

Authors: Stireman, John O., Greeney, Harold F., and Dyer, Lee A.

Source: Journal of Insect Science, 9(39) : 1-19

Published By: Entomological Society of America

URL: https://doi.org/10.1673/031.009.3901

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Downloaded From: https://complete.bioone.org/journals/Journal-of-Insect-Science on 28 Mar 2024
Terms of Use: https://complete.bioone.org/terms-of-use

Species richness and host associations of Lepidoptera-attacking Tachinidae in the northeast Ecuadorian Andes

John O. Stireman, III[1,a], Harold F. Greeney[2,b] and Lee A. Dyer[3,c]

[1] Department of Biological Sciences, 235A, BH, Wright State University, 3640 Colonel Glenn Highway, Dayton, OH 45435

[2] Yanayacu Biological Station & Center for Creative Studies, Cosanga, Ecuador, c/o 721 Foch y Amazonas, Quito, Ecuador

[3] Biology Department, University of Nevada; Reno, Nevada

Abstract

Most of the unknown biological diversity of macro-organisms remaining to be discovered and described lies in the tropical regions of the world and consists primarily of insects. Those insects with parasitoid lifestyles constitute a significant portion of insect diversity, yet parasitoids are among the most poorly known of major insect guilds in the humid tropics. Here we describe and analyze the richness of one diverse taxon of parasitoids, flies in the family Tachinidae, reared from Lepidoptera as part of a biological survey of Lepidoptera and their parasitoids in one mid-elevation (2000 m) area in the northeast Ecuadorian Andes. One hundred fifty-seven separable tachinid "morpho-species" were reared from approximately 160 species of Lepidoptera in 16 families. These tachinid flies were recovered from a sample of over 12,800 successful caterpillar rearing events that resulted in either adult Lepidoptera or parasitoids. Tachinid species accumulation and rarefaction curves exhibit no sign of reaching an asymptote and richness estimators indicate that the community likely consists of nearly twice this number of species (at minimum). Most tachinid species were reared infrequently, with 50% being represented by a single individual. The majority of species appeared to be relatively specialized on one or a few related hosts, but sampling was insufficient to make strong inferences regarding host range. The tribes Blondeliini and Goniini were the best represented, but some tribes that were expected to be common such as Tachinini and Winthemiini were poorly represented. The estimates of tachinid species richness derived here are suggestive of a far more diverse tachinid community than in temperate localities in North America. Additional rearing of Lepidoptera, as well as other herbivorous insect taxa, along with the use of additional collecting methods will be necessary to achieve a more accurate understanding of the richness of tropical Tachinidae and their contribution to broader patterns of tropical biodiversity.

Keywords: parasitoid, Diptera, biodiversity, caterpillars, tropical rainforest, morpho-species, insect rearing
Correspondence: [a]john.stireman@wright.edu, [b]revmmoss@yahoo.com, [c]ldyer@unr.edu
Received: 3 April 2008 | **Accepted:** 26 June 2008 | **Published:** 2 June 2009
Copyright: This is an open access paper. We use the Creative Commons Attribution 3.0 license that permits unrestricted use, provided that the paper is properly attributed.
ISSN: 1536-2442 | Volume 9, Number 39

Cite this paper as:
Stireman JO, Greeney HF, Dyer LA. 2009. Species richness and host associations of Lepidoptera-attacking Tachinidae in the northeast Ecuadorian Andes. 19pp. *Journal of Insect Science* 9:39, available online: insectscience.org/9.39

Downloaded From: https://complete.bioone.org/journals/Journal-of-Insect-Science on 28 Mar 2024
Terms of Use: https://complete.bioone.org/terms-of-use

Introduction

Despite more than a quarter millennium of exploration, categorization and description of organismal diversity, our understanding of the number of living species on earth is relatively poor (Gaston and Spicer 2004). This failure to gain a comprehensive understanding of diversity is due in part to growing use of molecular tools to explore biodiversity, which has opened a window into the hitherto unappreciated magnitude of microbial species richness that we are only beginning to understand (e.g., Rappé and Giovannoni 2003). However, estimates of the diversity of macroscopic organisms, such as insects, are nearly as variable and weakly supported as those of micro-organisms, with estimates of global insect species numbers ranging from as few as 4.8 million (Odegaard 2000) to over 30 million (including other arthropods; Erwin 1982). Much of the uncertainty in estimates of insect diversity is due to our poor understanding of tropical biodiversity (Gaston 1991; Godfray et al. 1999). Although there has been considerable debate regarding the extent of the undescribed richness of tropical insects (e.g., Novotny et al. 2002), one conclusion that researchers tend to agree upon is that we desperately need more surveys and documentation of tropical insects and their ecological interactions (Godfray et al. 1999).

One guild of tropical insects for which our knowledge is perhaps poorest, and which has the greatest potential for influencing estimates of insect diversity upwards, is the insect parasitoids (e.g., Brown 2004). Our poor understanding of tropical (and temperate) parasitoid biodiversity is due in part to their frequently small size, specialized ecological niches, and relatively small population sizes associated with their elevated trophic positions. Godfray (1994) estimated that parasitoids represent 8.5% of described insect species, yet some workers suggest that the fraction of insect species that possess parasitoid lifestyles may exceed 20% (LaSalle and Gauld 1991). Thus, to achieve an accurate appreciation of global insect diversity, we need to obtain reliable estimates of the diversity of parasitoid lineages in the tropics.

The focus of the current study is to document the diversity (species richness) and host associations of one particular group of tropical parasitoids, flies in the family Tachinidae (Figure 1). The family Tachinidae contains approximately 9,200 (Brown 2001) to 10,000 (Irwin et al. 2003) described species and ranks second in species richness among families of Diptera. All species of Tachinidae for which life histories are known are parasitoids of other arthropods, primarily insects. As parasitoids, tachinid flies are second only to the vast "Parasitica" group of Hymenoptera in diversity and ecological importance (Godfray 1994; Stireman and Singer 2003a). They are particularly frequent and apparent as parasitoids of larval Lepidoptera, especially macrolepidopteran familes such as Arctiidae, Noctuidae, Sphingidae and Nymphalidae (Gentry and Dyer 2002, Stireman and Singer 2003a; Janzen and Hallwachs 2008). Although tachinid parasitoids tend to be larger in size and more conspicuous than most of their hymenopteran parasitoid counterparts, the species richness and host-use patterns of tropical tachinids is equally obscure. In fact, despite their diversity and ecological importance as natural enemies, there has been little quantitative exploration of tachinid diversity at local, regional, or global scales (though see Stireman and Singer 2003b; O'Hara 2007; Stireman 2008).

Current estimates of the species richness of Tachinidae among geographical provinces suggest that the Neotropical region harbors the largest number of species and represents a geographic epicenter of tachinid diversification (Stireman et al. 2006; O'Hara 2007). The Neotropical region boasts an estimated 2,864 described species (Guimarães 1971) belonging to an impressive 822 genera, almost twice as species rich as any other geographic realm (O'Hara 2007). Although tachinids are found in a wide variety of habitats in the tropics and from sea level to alpine tundra, their diversity is most apparent at middle elevations (1000–2000m) along the mountain chains of tropical Central and South America, where tachinids are an abundant and conspicuous component of the diurnal insect fauna. Despite the large number of described species, it is generally thought that only a fraction of Neotropical Tachinidae have been described, and for most of these nothing is known about their life histories, host associations, or behavior (Guimarães 1977).

Here, we present preliminary analyses of the species diversity and host affiliations of Tachinidae reared from lepidopteran larvae the northeast Ecuadorian Andes. We present a preliminary list of genera and species reared and provide host–family affiliations for most taxa as well as notes on the taxonomy of the species reared and associations with host genera. In addition, the taxonomic composition of the tachinid community and emergent patterns of host-associations and host specificity are examined.

Materials and Methods

Collection

All tachinids in this study were reared as part of a collaborative biological survey and inventory project initiated in 2001 that is focused on surveying and inventorying plant-caterpillar-parasitoid associations in an Ecuadorian cloud forest (Stireman et al. 2005; Dyer et al. 2007). The survey project is centered at Yanayacu Biological Station & Center for Creative Studies (YBS), located at 2200m in the Quijos Valley, Napo Province, in northeastern Ecuadorian. Much of the ca. 2000 hectare reserve of adjacent Cabana San Isidro and YBS is relatively level cloud forest, some of the only remaining habitat of this type in the Andes (Figure 2). Although most collections of

Downloaded From: https://complete.bioone.org/journals/Journal-of-Insect-Science on 28 Mar 2024
Terms of Use: https://complete.bioone.org/terms-of-use



Figure 1. Examples of some of the Lepidoptera-attacking tachinid taxa commonly encountered around Yanayacu Biological Station. a. Goniini: *Gaediopsis* sp. b. Blondeliini: *Leptostylum* sp. (probably), c. Tachinini: *Epalpus* sp. d. Goniini: *Leschenaultia* sp. (photos by JOS).

caterpillars were made within 1–3 km of the station, collections were also made at higher and lower elevation sites in the surrounding region, primarily within 20 km. Because boundaries between sampling areas are difficult to define and relatively few caterpillars were sampled in these outlying areas, all samples were considered together. Thus, the sampling area should be considered to be on the order of 200 km^2 and encompass elevations from 500–3000 m (although only a small fraction of this area was sampled).

Caterpillars were systematically collected from $10m^2$ plots by Ecuadorian parataxonomists and Earthwatch volunteers. Vegetation within plots was visually scanned and all encountered caterpillars were collected along with material from their host plant. Additional caterpillars were collected opportunistically as they were encountered along trails and streams. Each individual caterpillar was assigned a unique number and the species (or morpho-species) of the caterpillar and its host-plant was recorded. Caterpillars were reared individually in clear plastic bags or glass jars in an open-walled, shaded rearing shed at ambient temperature and humidity. Every two days, bags were cleaned and foliage was replaced. After pupation, individuals were checked regularly and emerged adult Lepidoptera or parasitoids were collected and preserved. Throughout this process life history data were recorded (e.g., host, host plant, collection date, pupation date, eclosion date). Host plants were primarily identified by E. Narvaez (Universidad Central de Ecuador) and C. Chicaiza (Herbario Nacional Ecuatoriano). Caterpillars were assigned temporary morpho-

Downloaded From: https://complete.bioone.org/journals/Journal-of-Insect-Science on 28 Mar 2024
Terms of Use: https://complete.bioone.org/terms-of-use



Figure 2. Yanayacu Biological Station and the surrounding "hanging valley" of montane cloud forest where caterpillar hosts of tachinids were collected and reared (Napo Province, Ecuador).

species names if the species was unknown and later, if an adult was reared, added to the pool of specimens awaiting identification by taxonomic experts associated with the biological inventory project. Each caterpillar reared to either an adult moth/butterfly or a parasitoid is referred to as a "rearing event".

Upon eclosion, tachinids were killed by freezing and subsequently pinned along with their puparia. Initially these collections were stored at YBS in boxes under ambient (indoor) conditions, which resulted in humidity-associated damage to many specimens. When reliable 24-hour electric power became available at YBS, pinned specimens were stored in a −20°C freezer. Accumulated specimens were then sent to the laboratory of JOS at Wright State University for sorting and identification.

Identification

Tachinid specimens were initially sorted to genus using a key to Central American Tachinidae developed by D.M. Wood (Wood, in prep.), which is based, in part, on his generic key of Nearctic Tachinidae (Wood 1987). A number of specimens could not be assigned to genera on this basis for one of three reasons: 1) some reared specimens from Ecuador belong to genera not represented in the Central American fauna, 2) several specimens appeared to belong to undescribed genera (D.M. Wood pers. comm.), 3) some of the morphological distinctions between established genera appear to break down in the Neotropical fauna (e.g., see Wood 1985). Generic determinations for genera not present in Wood's unpublished key were determined in part by inspection by D.M. Wood, comparison with identified specimens in the Canadian National Collection of Insects (CNC) and/or with reference to C.H.T. Townsend's Manual of Myiology (1936–1945). These determinations should be treated as preliminary until further more focused taxonomic work is done with this material. JOS was unable to sort a small number of heavily damaged adult specimens to genus and these are omitted from this study.

After division into genera, specimens were further sorted into morpho-species with reference to external morphology (primarily chaetotaxy, wing patterns, shape and structure of head appendages, and color patterns). For some groups, further identification was made using keys and descriptions from the primary literature, through inspection by D.M. Wood or J.E. O'Hara, and by comparison with identified specimens in the CNC. For most specimens, however, specific determinations could not be made. Sorting of specimens into morpho-species was complicated by the fact that many "species" were represented by only one or a few individuals, and of those represented by multiple individuals, sometimes only a single

Downloaded From: https://complete.bioone.org/journals/Journal-of-Insect-Science on 28 Mar 2024
Terms of Use: https://complete.bioone.org/terms-of-use

sex was available for examination. Furthermore, it was often difficult to determine where intraspecific morphological variation ended and interspecific variation began, especially in several large genera in which there appear to be many undescribed species (e.g., *Siphona*, *Erythromelana*, *Lixophaga*, *Calolydella*). For this reason, estimates of the number of species and their host associations should be interpreted with some caution, and a low estimate (conservative "lumping" approach) of species richness is also provided. Although the divisions of individuals here are properly considered "morpho-species," they are referred to as species for simplicity. All reared and sorted specimens are currently housed in the Wright State University Collection of Insects. As further identifications are made and confirmed, voucher specimens will be returned to Ecuador and be housed in the Museo Ecuatoriana de Ciencias Naturales in Quito, Ecuador.

Relatively few of the host caterpillars from which tachinids have been reared have been identified to species at this time and have been given temporary morphospecies designations. Thus, the exact number of host species is uncertain. Many host species are currently being identified by collaborators on this Biological Survey and Inventory survey of caterpillars and parasitoids in the Ecuadorian Andes.

Analysis

Observed and estimated (rarefaction) species accumulation curves were calculated and plotted using the software EstimateS 7.5 (Colwell 2005). The rarefaction curve was calculated using the Mao Tau estimator (Colwell et al. 2004) with 1000 randomizations. EstimateS was also used to estimate the following species richness estimators and their 95% confidence limits (based on 1000 randomizations without replacement): ACE, ICE, Chao-1, Chao-2, Jack-1, Jack-2, Bootstrap and Michaelis-Menton (see Chao 2005, Colwell 2005, and references therein for detailed descriptions of these estimators). The accumulation of singletons and doubletons over time was also plotted with this software. A least-squares regression was used to assess the relationship between host range and the number of rearings for each tachinid species using PAST 1.8 (Hammer et al. 2001). Significance was assessed with 10,000 permutations. PAST 1.8 was also used to assess which commonly used model (log-series, log-normal, geometric, or broken stick) best fits the observed species abundance distribution.

Results

Four hundred eighty-nine adults representing approximately 157 species and 50 genera of Tachinidae were reared from 357 rearing events of approximately 160 morpho-species of Lepidoptera in 16 families (Table 1). In addition, one sarcophagid species (*Boettcheria* spp.) was reared on two occasions from larvae of an unknown saturniid (Table 1). This estimate of species number represents the number of morphologically differentiable groups of individuals and must be treated with some caution. A secondary examination of the specimens in which individuals deviating only slightly in morphology were "lumped" together results in a more conservative estimate of 139 species. An appreciable number of additional tachinid parasitism events were recorded in which the adult tachinids failed to eclose from observed puparia.

Considering all data generated by the caterpillar rearing project, the overall parasitism frequency (including all parasitoid taxa) is approximately 11% (based on >28,000 caterpillar records). Of the parasitoids identified to family (whether larvae, pupae, or adults), 37% were Tachinidae, with the remainder consisting of Hymenoptera, primarily Ichneumonoidea. Many more tachinids, as well as other parasitoid taxa, likely perished in hosts that succumbed to pathogens or other sources of mortality (e.g., less than 50% of collected Lepidoptera resulted in an adult insect, whether it be moth or parasitoid). If only caterpillar samples that produced an adult moth/butterfly or parasitoid are considered, the estimate of the total parasitism frequency rises to 24.8% (3186 of 12805 records).

Both the empirically observed species accumulation curve and the estimated Mau Tau rarefaction curve suggest that we have sampled only a fraction of the diversity of Lepidoptera-using Tachinidae at YBS and surrounding areas (Figure 3). The observed species accumulation curve is nearly a straight line (with a slight plateau near the center representing a time period in which a large number of the same host caterpillar was reared), with no indication of asymptotic behavior. The shape of the rarefaction curve suggests that the rate of new species accumulation may be decreasing from its initial values due to multiple rearing events of some species, but the rate still remains high (approximately one new species for every four rearing events). The Chao-1 estimator of total species richness predicts that the community of Lepidoptera-attacking Tachinidae in the sampled area consists of approximately 273 species, with a 95% confidence interval between 224 and 361 species (Figure 4; Table 2). Although the estimates stabilized to some extent after approximately 100 rearing events, both the mean estimates and associated confidence intervals continue to increase at a relatively constant rate. Most of the other species richness estimators resulted in similar estimates of 270–300 species (Table 2), although Jack-1 and bootstrap estimates predicted somewhat lower total species richness.

One half of the tachinid "species" reared (79 spp.), were reared only once (i.e., singletons; although in many cases these were gregarious and are represented by multiple individuals; Figure 5; Table 1). Another 20% (31 spp.) were reared on two occasions (doubletons), and only about 2% of species were reared ten or more times, resulting in a

Downloaded From: https://complete.bioone.org/journals/Journal-of-Insect-Science on 28 Mar 2024
Terms of Use: https://complete.bioone.org/terms-of-use

Table 1. A list of the tachinid genera thus far reared from caterpillars from the Yanayacu Biological Station and surrounding areas. The number of rearing events (N), the host families from which they have been reared, and notes about particular taxa are indicated.

Taxon	N	Host Family	Host Species/Notes
DEXIINAE			
CAMPYLOCHETINI			
Camplyocheta nr. *heteroneura* (Brauer & Berg.)	2	Saturniidae	*Gamelia neidhoeferi* Lemaire
Camplyocheta sp. 2	6	Geometridae	most reared from hosts on *Diplazium costale* (Dryopteridaceae)
Camplyocheta nr. sp. 2	2	Geometridae	
Camplyocheta sp. 3	7	Geometridae (6)	most reared from hosts on *Diplazium costale*
		Pyralidae	
Camplyocheta sp. 4	1	Choreutidae	
Camplyocheta sp. 5	1	Geometridae	*Pantherodes colubraria viperaria* Thierry-Mieg
THELAIRINI			
Polygaster sp. 1	2	Geometridae	
URAMYIINI			
Thelairaporia "*brasiliensis*" grp. (Guimarães) sp. 1	2	Arctiidae	(Distinction between this genus and *Uramya* is blurred)
Thelairaporia "*brasiliensis*" grp. (Guimarães) sp. 1a	1	Arctiidae	
Thelairaporia sp. 1b	1	Arctiidae	(1b and 1c are from the same host sp.)
Thelairaporia sp. 1c	1	Arctiidae	
Thelairaporia sp. 2	3	Limacodidae	*Parasa macrodonta* Hering & Hopp,
			Perola sp.
Thelairaporia sp. 2a	2	Arctiidae	
Thelairaporia sp. 3	1	Arctiidae	
Thelairaporia sp. 4	2	Limacodidae	
Thelairaporia sp. 5	1	Geometridae (?)	Questionable record (unusual host)
Uramya sp. nr. *fasciata*	1	Megalopygidae	
Uramya sp. nr. *quadrimaculata*	5	Megalopygidae (4)	*Macara alydda* Druce
		Saturniidae	
Uramya sp. 3	2	Arctiidae	
UNPLACED			
Argyromima mirabilis Tnsd.	3	Nymphalidae	*Pedaliodes montagna* Adams & Bernard
EXORISTINAE			
BLONDELIINI			
Anoxynops sp. 2	1	Nymphalidae	
Anoxynops sp. 3	1	Notodontidae	
Calolydella sp. 1	2	Arctiidae	
Calolydella sp. 1a	12	Nymphalidae (10)	*Actinote diceus* Latreille (2), *A.stratonice* Latrielle (7)
		Geometridae	
		Arctiidae	
Calolydella sp. 1b	2	Nymphalidae	*Actinote* sp.
Calolydella nr *geminata* Tnsd. sp. 2	7	Nymphalidae	*Actinote stratonice* Latreille

Downloaded From: https://complete.bioone.org/journals/Journal-of-Insect-Science on 28 Mar 2024
Terms of Use: https://complete.bioone.org/terms-of-use

Table 1 (con't.)

Taxon	N	Host Family	Host Species/Notes
Calolydella sp. 3	1	Notodontidae	
Calolydella sp. 4	1	Arctiidae	
Calolydella sp. 5	2	Arctiidae, Geometridae	
Calolydella sp. 6	3	Nymphalidae, Noctuidae	*Heliconius congener* Weymer
Calolydella sp. 7	1	Arctiidae	
Eribella sp. 1	4	Geometridae (3), Pyralidae	One pyralid record (on a different host plant) may represent a distinct species.
Erythromelana sp.1	6	Geometridae (5)	*Eois pallidicosta* Warren
		Megalopygidae	
Erythromelana sp.1a	2	Geometridae	*Eois* sp.
Erythromelana sp. 2	1	Geometridae	
Erythromelana sp. 3	5	Geometridae (4)	*Eois pallidicosta* Warren, *Eois olivacea* Feder & Rogenhofer
		Pyralidae	
Erythromelana sp. 4	2	Geometridae	
Erythromelana sp. 5	2	Geometridae	*Eois* sp.
Erythromelana sp. 6	1	Geometridae	*Eois* sp.
Erythromelana sp. 7	1	Geometridae	*Eois* sp.
Erythromelana sp. 8	3	Geometridae	*Eois olivacea* Feder & Rogenhofer
Erythromelana sp. 9	3	Geometridae	*Eois pallidicosta* Warren *Eois* sp. nr. *sagittaria* Snellen
Erythromelana sp. 10	1	Geometridae	*Eois* sp.
Erythromelana (genus nr.) sp. 3	1	Geometridae	
Eucelatoria sp. 1 (*Vibrissina*?)	1	Geometridae	*Eois* sp.
Eucelatoria sp. 2	2	Nymphalidae	(Ithomiinae)
Eucelatoria sp. 3	1	Geometridae	
Eucelatoria sp. 4	3	Pieridae	*Leptophobia* sp.
Eucelatoria sp. 5	4	Pyralidae	
Eucelatoria sp. 6	1	Arctiidae	
Eucelatoria sp. 8	1	Tortricidae	
Eucelatoria sp. 9	3	Geometridae	
Eucelatoria sp. 10	1	unknown	
Eucelatoria sp. 11	2	unknown	
Italispedia sp. 1	3	Notodontidae	*Xenomigia* sp.
Italispedia sp. 2	1	Geometridae	*Pantherodes colubraria viperaria* Thierry-Mieg
Leptomacquartia prob. *planifrons* Tnsd.	1	Noctuidae	
Leptostylum sp. 1	2	Saturniidae	*Gamelia neidhoeferi* Lemaire, *Pseudoautomeris yourii* Lemaire
Leptostylum sp. 1a	2	Saturniidae	*Automeris abdominalis* Felder & Rogenhofer
Leptostylum sp. 2	2	Saturniidae	
Leptostylum sp. 3	1	Saturniidae	*Automeris abdominalis* Felder & Rogenhofer
Leptostylum sp. 4	1	Saturniidae	*Paradirphia geneforti* Bouvier
Lixophaga sp. 1	4	Pyralidae	(species listed in this genus may actually belong to several genera)

Downloaded From: https://complete.bioone.org/journals/Journal-of-Insect-Science on 28 Mar 2024
Terms of Use: https://complete.bioone.org/terms-of-use

Table 1 (con't.)

Taxon	N	Host Family	Host Species/Notes
Lixophaga sp. 3	1	Pyralidae	
Lixophaga sp. 4	2	Limacodidae, Arctiidae	
Lixophaga sp. 5	1	unknown	
Lixophaga sp. 7 (genus nr.)	31	Nymphalidae	*Pteronymia zerlina* Hewitson
Lixophaga sp. 8 (genus nr.)	1	Geometridae	*Pantherodes colubraria viperaria* Thierry-Mieg
Lixophaga sp. 9	2	Pyralidae	
Lixophaga/Eucelatoria sp. 1	1	Noctuidae	
Ptilodegeeria sp. 1	1	Tenthredinidae (Hymenoptera)	
ERYCIINI			
Carcelia prob. *montanus* (Tnsd.)	3	Arctiidae	
Carcelia flavirostris/orellana	2	Arctiidae	*Antichloris ornate* Druce, *Saurita mosca* Dognin
Carcelia sp. 2	5	Arctiidae	
Carcelia sp. 3	1	Arctiidae	*Phaegoptera decrepridoides* Rothschild
Carcelia sp. 5	1	Arctiidae	
Drino sp. 1	1	Sphingidae	
Lespesia sp. 1	4	Nymphalidae	
Lespesia sp. 2	3	Saturniidae	
Eulobomyia (see Woodley and Arnaud 2008a,b)	5	Nymphalidae, Pieridae	*Actinote diceus* Latrielle, *Leptophobia eleone* Doubleday (?)
Zizyphomyia sp. 1	1	Nymphalidae	
GONIINI			
Argyrochaetona sp. 1	1	Pyralidae	
Belapharipa sp. 3	1	Noctuidae	
Chrysoexorista sp. 1	2	Pyralidae	small in size for this genus
Gaediopsis punoenisis Vim. & Souk.	1	Pyralidae	
Gaediopsis sp. 2	3	Nymphalidae	*Actinote stratonice* Latrielle, *A. diceus* Latrielle
Hyphantrophaga sp. 1	1	Hesperiidae	
Hyphantrophaga sp. 2	3	Apatelodidae	
Leschenaultia sp. 1	1	Arctiidae	
Leschenaultia sp. 1a	1	Arctiidae	
Leschenaultia sp. 2 nr. *leucophrys* Wied.	9	Arctiidae (8), Apatelodidae	
Leschenaultia sp. 2a nr. *leucophrys* Wied.	8	Arctiidae (4), Nymphalidae (2), Noctuidae, Sphingidae	*Pseudomaniola loxo* Dognin
Mystacella sp. 1	1	Pyralidae	
Mystacella sp. 2	3	Pyralidae	
Mystacella-Chrysoexorista sp. 3	1	unknown	
Patelloa andina (Tnsd.)	6	Noctuidae, Geometridae	*Hypnea valkeralis* Schaus
Patelloa xanthura/similis sp. 2	5	Hesperiidae (2), Arctiidae (2)	Possibly 2 spp. included here.
Patelloa xanthura (or nr.) sp. 3	1	Arctiidae	

Downloaded From: https://complete.bioone.org/journals/Journal-of-Insect-Science on 28 Mar 2024
Terms of Use: https://complete.bioone.org/terms-of-use

Table 1 (con't.)

Taxon	N	Host Family	Host Species/Notes
Patelloa xanthura (or nr.) sp. 4	1	Arctiidae	
Patelloa sp. 5	2	Saturniidae, Apatelodidae	
WINTHEMIINI			
Winthemia nr. *analisella* Thompson	1	Nymphalidae	*Antirrhea porphyrosticta* Watkins
Winthemia sp. 2	1	Hesperiidae	*Pyrrhopyge papius* Hopffer
Winthemia sp. 3	1	Noctuidae	
TACHININAE			
ERNESTIINI			
Bombylomyia	1	Geometridae	
Fasslomyia nr. *fantastica* Tnsd.	2	Arctiidae, Apatelodidae	
GRAPHOGASTERINI			
Phytomyptera sp. 1	5	Pyralidae	
Phytomyptera sp. 2	1	Pyralidae	
Phytomyptera sp. 3	1	Geometridae	*Eois* sp.
JURINIINI			
Jurinella sp. 1	1	Apatelodidae	
LESKIINI			
Leskia (*leskiopalpus* grp.) sp. 1	1	unknown	
Micronychiops sp. 1	1	unknown	
NEMORAEINI			
Xanthophyto sp. 1	1	Geometridae	
Xanthophyto sp. 2	1	Geometridae	*Pantherodes colubraria viperaria* Thierry-Mieg
Xanthophyto sp. 3	1	Choreutidae	
Macromya sp. 1	3	Arctiidae	
Macromya sp. 2	1	Saturniidae	
POLIDEINI			
Chlorohystricia cussirilis Reinhard	1	unknown	
Chlorohystricia sp. 2	1	Hesperiidae	
Chrysotachina sp. nr. *peruviana* (Tnsd.)	2	Hesperiidae	
Eucheirophaga (?)	1	Hesperiidae	
Hystricia laxa Curran	1	Saturniidae	*Automeris* sp.
Hystricia nr. *browni* Curran	3	Arctiidae	*Dysschema* nr. *joiceyi* Dognin
Hystricia nr. *micans* Curran	2	Apatelodidae	
Hystricia nr. *vultura* Curran	8	Arctiidae	
Hystricia sp. 5	2	Arctiidae	
Hystricia sp. 6	2	Arctiidae	*Bertholdia sp.*
Hystricia sp. 7	1	unknown	
Hystricia sp. 8	3	Saturniidae	
Hystricia sp. 9	1	Saturniidae	

Downloaded From: https://complete.bioone.org/journals/Journal-of-Insect-Science on 28 Mar 2024
Terms of Use: https://complete.bioone.org/terms-of-use

Table 1 (con't.)

Taxon	N	Host Family	Host Species/Notes
Genus nr. *Exoristoides*	1	Pyralidae	
Genus nr. *Chrysotachina*	1	Pyralidae	
Genus nr. *Phobetromyia*	1	Noctuidae	
SIPHONINI			
Actia sp. 1	1	Pyralidae	
Actia sp. 1a	1	Pyralidae	
Actia sp. 2	1	Pyralidae	
Actia sp. 3	1	Pyralidae	
Actia (or genus nr.) sp. 4	1	Saturniidae	*Gamelia neidhoeferi* Lemaire
Siphona sp. 4	2	Geometridae	*Eois* sp.
Siphona sp. 4a	1	Geometridae	
Siphona sp. 5	1	Geometridae	
Siphona sp. 7	3	Pyralidae	
Siphona sp. 7a	3	Pyralidae (2), Geometridae	
Siphona sp. 8	1	Pyralidae	
Siphona sp. 8a	4	Pyralidae	
Siphona (*Siphonopsis*) sp. 9	1	unknown	
Siphona (*Pseudosiphona*) sp. 10	2	Geometridae	*Eois* sp.
Siphona sp. 11	1	Geometridae	*Eois olivacea* Felder & Rogenhofer
Siphona sp. 11a	2	Geometridae	*Eois* sp.
Siphona sp. 12	1	Pyralidae	
Siphona sp. 13	1	Geometridae	
TACHININI			
Rhachoepalpus immaculatus (Macq.)	4	Saturniidae, Apatelodidae	
Trichophora	1	unknown	
TELOTHYRINI			
Telothyria sp.1	2	Pyralidae	
Telothyria sp.2	1	Pyralidae	
Telothyria sp.3	1	Pyralidae	
UNPLACED GENERA			
Unplaced genus # 1	1	Notodontidae	*Xenomigia* sp.
Unplaced genus # 2	1	Geometridae	*Eois* sp.
TOTAL	**357**		
SARCOPHAGIDAE			
BOETTCHERIINI			
Boettcheria	2	Saturniidae	*Dirphia* sp.

highly left-skewed abundance distribution that best fits a log-series distribution ($p = 0.9998$; Figure 5). The number of singletons increases nearly linearly with rearing events with no clear evidence of leveling off, suggesting that many rare species remain to be sampled and that sampling intensity, at this point, is inadequate to characterize the community well (Figure 6). In contrast, numbers of doubletons, are beginning to stabilize and even decline slightly, perhaps due to recent attempts to widen the diversity of host caterpillars sampled.

Downloaded From: https://complete.bioone.org/journals/Journal-of-Insect-Science on 28 Mar 2024
Terms of Use: https://complete.bioone.org/terms-of-use




Figure 3. Accumulation curves of tachinids species against the number of rearing events. Sobs (empirical) equals the observed number of species, Sobs (Mau Tau) is the mean rarefied Mau Tau estimate of observed species based on 1000 randomizations of rearing events (see text).




Figure 4. Mean Chao-1 estimates of total species richness of Tachinidae with sample size (number of rearing events) along with upper and lower 95% confidence limits. The mean estimate at full sample size is 273 species.

Most of the tachinid species (74%) were reared from only a single host species, and no species was reared from more than seven host species (Figure 7). The mean number of host species per tachinid species was 1.38 ± 0.06 (SE) and the mean number of host families was 1.16 ± 0.04. However, these figures reveal little about actual host ranges, as most tachinids were only reared a single time (see above), and therefore could not have been

Downloaded From: https://complete.bioone.org/journals/Journal-of-Insect-Science on 28 Mar 2024
Terms of Use: https://complete.bioone.org/terms-of-use

Table 2. Mean estimates of total species richness of the tachinid community of Yanayacu Biological Station and surrounding areas that use Lepidoptera as hosts, based on 1000 randomizations.

Richess Estimator	Richness Estimate
See Chao (2005) and Colwell (2005) for detailed explanation of estimators and their calculation.	
ACE	279.3
ICE	279.2
Chao-1	273.6
Chao-2	273.6
Jack-1	243.7
Jack-2	297.5
Bootstrap	194.9
MM runs	287.8




Figure 5. The species abundance distribution of tachinids reared in the current study in terms of number of rearing events per species.

reared from multiple host species. A significant relationship exists ($R^2 = 0.354$, $P < 0.0001$; Figure 8) between the inferred host range of tachinid species and the number of rearing events, as has been found in other analyses of tachinid host ranges (e.g., Eggleton and Gaston 1992; Belshaw 1994). This result suggests that the perceived host ranges of the Ecuadorian tachinid species are likely to increase with greater sampling effort. General host associations of tachinid genera obtained from the rearing data are listed in Table 1.

Discussion

Species Richness

Given the linear trajectory of the species accumulation curve and the only slightly declining slope of the rarefaction curve, it is difficult to gauge how many species of Tachinidae might occur in the sampled region. Furthermore, there are few equivalent studies of Tachinidae with which to compare these estimates. In a previous Lepidoptera rearing study in southern Arizona, Stireman and Singer (2003b) reared 64 tachinid species in approximately 1240 rearing events. That study sampled from a considerably larger area than does the current one, but

Downloaded From: https://complete.bioone.org/journals/Journal-of-Insect-Science on 28 Mar 2024
Terms of Use: https://complete.bioone.org/terms-of-use




Figure 6. Observed accumulation of singletons (species recorded only once) and doubletons (species recorded twice) with increasing number of tachinid rearing events.




Figure 7. Frequency histogram of the number of host species from which each tachinid species was reared.

included fewer individuals and species of caterpillar hosts. Nonetheless, rarefaction indicates that an equivalent sampling effort in Arizona, in terms of rearing events (357), would yield a considerably smaller figure of approximately 46 tachinid species (JOS, unpub data), indicating a much larger fauna at the Ecuador site. Recent

Downloaded From: https://complete.bioone.org/journals/Journal-of-Insect-Science on 28 Mar 2024
Terms of Use: https://complete.bioone.org/terms-of-use




Figure 8. Least squares regression of the number of host species against the number of rearing events. (regression equation: $y = 0.273x + 0.76$, $R^2 = 0.354$, $p < 0.0001$).

caterpillar rearing studies in the eastern United States have yielded fewer tachinid species (e.g., ca. 30 by Barbosa and Caldas [2004], and 24 by Stazanac et al. [2001]), but these studies were narrower in scope, limiting the usefulness of comparison.

A recent catalogue places the number of Nearctic tachinids (north of Mexico) at 1345 species (O'Hara and Wood 2004). Assuming that 75% of these species use Lepidoptera as hosts, an estimated 1076 species could be reared from Lepidoptera. This suggests that the relatively small area sampled in and around YBS (ca. 200 km^2) in Ecuador is home to about a quarter as many tachinid species as the entire Nearctic Region north of Mexico (ca. 20 million km^2). This difference in the areas sampled is conservative, as most of the samples at YBS were taken from ca. 10 km^2 surrounding the station. Furthermore, the estimators of total numbers of species at YBS (e.g., ICE, Chao-2) all exhibit signs of continuing increase with more rearing events (e.g., Figure 4). These estimators may not, therefore, be reliable estimates of total richness, but instead represent lower bounds (Longino et al. 2002).

The high diversity of tachinids in this mid-elevation tropical Andean site is also indicated by preliminary examination of tachinid specimens (ca. 1000 individuals) obtained by hand collecting around the YBS and in surrounding areas, which suggest relatively little overlap with reared material (unpub. data). This lack of overlap persists even when considering only those groups that are likely to use Lepidoptera as hosts (e.g. Goniini, Tachinini, Eryciini), supporting the prediction that many additional species remain to be reared from Lepidoptera in this area. The species abundance distribution (Figure 5) suggests a preponderance of rare species (70% singletons and doubletons). However, as this distribution reflects in large part inadequate sampling, it is premature to make conclusions concerning the relative abundance and rarity of species or higher taxa.

It has been suggested that species richness of some hymenopteran parasitoid groups may not increase with decreasing latitude or may increase more slowly than other insect taxa (Janzen and Pond 1975; Janzen 1981; though see Sääksjärvi et al. 2004). This trend may be due to greater resource fragmentation and/or increasingly toxic chemical defenses in potential hosts in the tropics (e.g. Janzen 1981; Gauld et al. 1992). The high species richness of Tachinidae at this site relative to similar Lepidoptera-rearing studies in more temperate areas (e.g., Stireman and Singer 2003b) suggests that tachinids exhibit a marked latitudinal gradient in species richness similar to their (primarily) phytophagous insect hosts. One speculative reason for this difference is that tachinids may be relatively less affected by resource fragmentation and host chemical defenses.

Accuracy of species delimitations

As discussed in the Methods, there are several factors which may have resulted in an overestimation of tachinid diversity in the current study. For example, extensive intraspecific variation or sex-specific morphological

Downloaded From: https://complete.bioone.org/journals/Journal-of-Insect-Science on 28 Mar 2024
Terms of Use: https://complete.bioone.org/terms-of-use

variation could lead to splitting of one species into multiple morpho-species. These problems are somewhat mitigated by the approach employed here of rearing hosts to collect parasitoids, where multiple individuals (often both sexes) are frequently reared from a single host. In addition, associations of parasitoids with particular host taxa can provide ecological evidence that may support morphological divisions of the parasitoids, as exemplified in a recent study of the tachinid genus *Belvosia* in Costa Rica (Smith et al. 2006). However, if understanding patterns of host association and host range is a goal, using this information in defining parasitoid species may result in tautological reasoning.

It is also possible that the number of distinct morpho-species identified here is actually an underestimate of the true species diversity. Observed variation among species may be easily ascribed to intra-specific variation and considerable existing variation may go unrecognized. Given recent evidence of sympatric, morphologically cryptic, but apparently reproductively isolated populations of tropical tachinids that differ in host use (Smith et al. 2006; Smith et al. 2007), it seems likely that the species richness reported here is underestimated. The great diversity and confusing taxonomy of Neotropical tachinids suggests that the region may be an "epicenter" of recent tachinid diversification (Stireman et al. 2006; supporting the general pattern of elevated speciation rates in the tropics; Mittelbach et al. 2007). Furthermore, tachinid diversity appears to peak at intermediate elevations in the tropics (Wood, pers. comm.). Thus, areas such as YBS, at middle-elevations on the eastern equatorial Andes, may be particularly rich with morphologically similar (i.e., cryptic) tachinid species.

It should be noted that currently fewer than 10% of the reared species have been assigned to a named species (or "near" a named species), and it could be conservatively estimated that perhaps 75% or more have not been described. Similarly, it is probable that >90% of the species have previously never been associated with a host in the scientific literature. This figure may be revised downward as we gain a better understanding of the overlap in taxa between this Ecuadorian site and other major Lepidoptera rearing efforts in Costa Rica (Gentry and Dyer 2002; Janzen and Hallwachs 2007). Many of the tachinids reared in this study are currently featured in a preliminary online guide to the tachinids of Ecuador with photos, taxonomic notes, and life history information (see http://www.caterpillars.org).

Distribution of Taxa

Certain subfamilies and tribes of Tachinidae were extremely well represented in the collection of reared species and others poorly so (Table 1). In part, this is due to the focus on lepidopteran hosts, such that higher taxa containing species that primarily attack non-lepidopteran hosts such as Phasiinae and Dexiini are not well represented. However, relative frequencies of taxa reared from caterpillars are generally consistent with observed frequencies of taxa in hand-collected samples from the same area. For example, extensive hand collections made along roadsides and trailsides contained no representatives of Phasiinae. Dexiini were also rare in the hand collections, despite extensive collecting effort on flowers (e.g., Asteraceae) where many dexiine genera (and Phasiinae) often take nectar and/or pollen in other regions. Voriini were also markedly absent in collections obtained by hand and through host rearing, despite the general use of Lepidoptera as hosts by members of this tribe (e.g., Geometridae; Arnaud 1978).

Over one half of all species reared (and one half of all tachinid rearing events) belong to the subfamily Exoristinae. Blondeliini was the most frequently reared tribe and was represented by a diverse assemblage of closely related genera, most of which were represented by several species (e.g., *Calolydella, Eucelatoria, Lixophaga, Erythromelana*). This tribe accounted for over 30% of both rearing events and tachinid species reared. Goniini (Figure 1a, 1d) and Eryciini were also responsible for appreciable numbers of parasitism events (12.6% and 5.7% of species reared respectively), although certain taxa well represented in D.H. Janzen and W. Hallwach's (2007) caterpillar rearing database, such as *Drino*, *Lespesia*, and *Belvosia* were noticeably rare or absent. The genus *Winthemia*, which is commonly reared from macrolepidoptera in the temperate zone and Central American tropics (Arnaud 1978; Janzen and Hallwachs 2007), was also conspicuously underrepresented.

The subfamily Tachininae was well represented (33.5% of reared species), but in this case there was incongruence between reared and hand-collected taxa. Field observations and hand-netting indicate an inordinately diverse fauna of Tachinini, particularly the "big fuzzy" taxa such as *Epalpus* (Figure 1c) and allied genera (e.g., *Lindigepalpus, Parepalpus, Eulasiopalpus*). Despite these observations, few Tachinini were reared (only 1.3% of total species). This may be due to the relatively low rearing numbers of large-bodied caterpillar taxa, such as Sphingidae and Saturniidae, capable of hosting these bulky tachinids. These large and active tachinids may also be relatively more apparent in the field than other tachinid taxa (i.e., their abundance is overestimated due to their conspicuousness). In contrast, Siphonini were common and diverse in both reared (11.4% of species) and netted collections. The relatively large number of polideine species reared (15 species) was somewhat surprising, given the limited representation of this tribe in other Lepidoptera rearing studies (e.g., Stireman and Singer 2003b). Much of this diversity of Polideini was a function of the large number of species (9) of the tachinine-like genus *Hystricia*.

Downloaded From: https://complete.bioone.org/journals/Journal-of-Insect-Science on 28 Mar 2024
Terms of Use: https://complete.bioone.org/terms-of-use

Host associations

Most of the family-level host associations of taxa reported here (Table 1), are consistent with previously established host-associations of particular tachinid groups (Guimarães 1977; Arnaud 1978). For example, *Carcelia* and *Leschenaultia* (Figure 1d) species were associated primarily with hosts in the family Arctiidae; uramyines were associated with Megalopygidae, Arctiidae, and Limacodidae; *Eucelatoria* species were primarily associated with Geometridae; and siphonines were associated with hosts of small size such as Geometridae and Pyralidae. Additional strong associations that were observed included *Leptostylum* with Saturniidae (Figure 1b) and *Erythromelana* with Geometridae. It is interesting to note that many *Lixophaga* spp. were reared from shelter-building Pyralidae larvae, reflecting the ability of their host searching larvae to attack of concealed hosts.

Although a thorough evaluation of the parasitism frequency of various ecological guilds or phylogenetic clades of hosts is beyond the scope of this paper, a preliminary examination of parasitism frequency by host family suggests that tachinids do not use all hosts with equal frequency. For example, the families Limacodidae, Megalopygidae, Sphinigidae, Saturniidae, and Arctiidae are attacked at relatively high frequencies by tachinids (<10%), while Geometridae, Noctuidae, and Choreutidae were parasitized less frequently. This variation probably reflects host ecology more than phylogenetic or physiological interactions between host and parasitoid (Stireman and Singer 2003a). Interestingly, despite their shelter building habit and small size, Pyralidae experienced relatively high parasitism by tachinids, possibly due to the attraction of parasitoids by the copious frass built up inside shelters (Roth et al. 1978).

It has been suggested that the host associations of tachinid species may depend more upon the microhabitat and diet of potential host species than the taxonomic/phylogenetic affiliations of the hosts (Crosskey 1980; Feener & Brown 1997; Stireman and Singer 2003a). This may reflect the importance of host location mechanisms in shaping patterns of host use and the indirect strategies by which many tachinids parasitize hosts (Feener & Brown 1997; Stireman et al. 2006). Although the current study provides several clear examples of broad associations between tachinid and host taxa (see above), many associations also support an important role of host-plants in shaping patterns of tachinid host use. For example, *Uramya* nr. *quadrimaculata* was reared primarily from Megalopygidae (one of the typical host families for this tachinid genus; Arnaud 1978); however, this species was also reared from a completely unrelated species of Saturniidae that fed on the same host plant (*Acalypha*, sp.; Euphorbiaceae). As another example, *Erythromelana* species tended to be associated with Geometridae, especially *Eois* sp. on plants in the genus *Piper*, but at least one species was also found to attack unrelated Pyralidae on the same *Piper* host plants.

Due to the relatively low sampling intensity and infrequent rearing events of particular tachinid species, little can be surmised concerning the breadth of host use for the tachinid species reared in this study. Observed host associations suggest, however, that there are few, if any, true generalists in this fauna and that most species are likely specialized on a few taxonomically or ecologically related host species. This contrasts with some analyses of host range in temperate tachinid taxa (Eggleton and Gaston 1992; but see Stireman and Singer 2003b), suggesting that there may exist a latitudinal gradient in specialization in Tachinidae as has been indicated for their lepidopteran hosts (Dyer et al. 2007).

Future Directions

Considerably more host rearing will be necessary to achieve a robust understanding of the diversity and taxonomic composition of this local community of Tachinidae. Despite the relatively large number of species reared thus far, it is clear that only a fraction of tachinid species in this diverse community have been sampled. Focused collection and rearing of caterpillars from the forest canopy may be one strategy to obtain greater proportion of the community of tachinid species that attack Lepidoptera. Furthermore, rearing of additional herbivorous insect taxa such as Hemiptera, Symphyta, and Chrysomelidae would significantly expand the number of species and taxonomic diversity of tachinids recorded from the region. The caterpillar rearing program described here is currently being augmented by hand-netting as well as pan-trapping and Malaise trapping, which will allow more exhaustive estimation of the total richness of the tachinid community, as well as a more thorough examination of biases associated with different sampling methods.

In addition to establishing the patterns of diversity and host associations of Tachinidae, much of the material reared from this project will contribute to future species descriptions, taxonomic revisions, and phylogenetic analyses of this poorly known Neotropical fauna. One benefit of the rearing approach to sampling employed here is that it provides basic natural history and life history information such as host-associations, plant-associations and developmental phenology. In addition to the inherent worth of this information, it may prove useful in establishing species limits and phylogenetic relationships. Another benefit of the approach is that in gregarious species (i.e., those in which multiple individuals can develop in a single host) sexes can be associated. This limits the possibility of each sex being described as a distinct species. As recent studies of Costa Rican tachinids have shown (Smith et al. 2006; 2007), DNA data can aid substantially in delineating species. DNA samples are now being gathered for mtDNA "barcode" sequencing of the reared tachinids, which may also be useful comparing

Downloaded From: https://complete.bioone.org/journals/Journal-of-Insect-Science on 28 Mar 2024
Terms of Use: https://complete.bioone.org/terms-of-use

this Andean tachinid fauna to that in Costa Rica to assess species turnover and geographic population structure of species.

Our current lack of understanding of tropical biodiversity is particularly alarming when one considers the rate of tropical deforestations and estimated rates of species extinctions measured on the scale of species loss per week or month (Pimm and Raven 2000). Given their specialized life histories and elevated trophic position, tropical parasitoids may be particularly prone to extinction via habitat loss. This problem is compounded due to a current taxonomic crisis (Brown 2005), wherein the rate of description of new species from the Neotropical Region is extremely low, suggesting that many species are quietly disappearing without ever being recorded. A major goal of this study and associated research is to provide baseline documentation of the diversity and natural history of caterpillars and parasitoids in the Ecuadorian Andes. The results reported here, on the richness and host associations of Tachinidae, represent a small but important step towards understanding tropical insect diversity.

Acknowledgments

We would like to thank Monty Wood and Jim O'Hara for their valuable assistance in identifying a number of the tachinid taxa listed herein. Hosts were identified by G. Gentry, J. Miller, G. Brehm, M. Epstein, K. Willmott, M. Pogue, S. Rab-Green, A. Brower, and G. Lamas. The caterpillar-parasitoid biological inventory project from which this data is derived was supported by the U.S. National Science Foundation (DEB0346729 and DEB0717092) and Earthwatch. We would like to acknowledge our collaborators T. R. Walla, Grant Gentry, and Genoveva Rodriguez Castaneda who have overseen the rearing program. We would also like to thank Yanayacu Biological Station for its support, Hilary Devlin for specimen curation, the Ecuadorian "gusaneros", Wilmer Simbaña, Rafael Granizo, Marco Gualinga Licuy, and Santiago Gualinga Licuy, who reared caterpillars and prepared material, and the many Earthwatch volunteers who helped to collect and rear the caterpillars.

Editor's note

Paper copies of this article will be deposited in the following libraries. Senckenberg Library, Frankfurt Germany; National Museum of Natural History, Paris, France; Field Museum of Natural History, Chicago, Illinois USA; the University of Wisconsin, Madison, USA; the University of Arizona, Tucson, Arizona USA; Smithsonian Institution Libraries, Washington D.C. USA; The Linnean Society, London, England.

References

Arnaud PH. 1978. *A Host-Parasite Catalog of North American Tachinidae (Diptera)*. USDA Misc. . *Publ* 1319:

Belshaw R. In: Hawkins BA, Sheehan W, editors. 1994. Life history characteristics of Tachinidae (Diptera) and their effect on polyphagy. *Parasitoid Community Ecology*145-62. Oxford University Press

Barbosa P, Caldas A. 2004. Patterns of parasitoid-host associations in differentially parasitized macrolepidopteran assemblages on Black Willow *Salix nigra* (Marsh) and Box Elder *Acer negundo* L. *Basic and Applied Ecology* 5: 75-85.

Brown BV. 2001. Diversity of flies, gnats and mosquitoes. In: Levin SA, editor. *Encyclopedia of Biodiversity* 2: , pp. 815-826. Academic Press

Brown BV. 2004. Diversity of ant-decapitating flies from the ALAS project: new results and projections. *Sociobiology* 44: 683-688.

Brown BV. 2005. Malaise trap catches and the crisis in Neotropical Dipterology. *American Entomologist* 51: 180-183.

Chao A. In: Balakrishnan N, Read CB, Vidakovic B, editors. 2005. Species richness estimation. *Encyclopedia of Statistical Sciences*7909-7916. Wiley

Colwell RK, Mao CX, Chang J. 2004. Interpolating, extrapolating, and comparing incidence-based species accumulation curves. *Ecology* 85: 2717-2727.

Colwell RK. 2005. EstimateS: Statistical estimation of species richness and shared species from samples. Version 7.5. *User's Guide and application*. Published at: http://purl.oclc.org/estimates.

Crosskey RW. 1980. Family Tachinidae. In: Crosskey RW, editor. *Catalogue of the Diptera of the Afrotropical Region*, pp. 822-82. British Museum of Natural History, London British Museum of Natural History, London

Dyer LA, Singer MS, Lill JT, Stireman JO, Gentry GL, Marquis RJ, Ricklefs RE, Greeney HF, Wagner DL, Morais HC, Diniz IR, Kursar TA, Coley PD. 2007. Host specificity of Lepidoptera in tropical and temperate forests. *Nature* 448: 606-700.

Eggleton P, Gaston KJ. 1992. Tachinid host ranges: a reappraisal (Diptera: Tachinidae). *Entomologist's Gazette* 43: 139-143.

Erwin T. 1982. Tropical forests: their richness in Coleoptera and other species. *Coleopterists' Bulletin* 36: 74-75.

Feener DH, Brown BV. 1997. Diptera as parasitoids. *Annual Review of Entomology* 42: 73-97.

Gauld ID, Gaston KJ, Janzen DH. 1992. Plant allelochemicals, tritrophic interactions and the anomalous diversity of tropical parasitoids: the 'nasty' host hypothesis. *Oikos* 65: 353-57.

Gaston KJ, Spicer JI. 2004. *Biodiversity: an introduction*, 2nd Ednd edition. Blackwell Publishing

Gaston KJ. 1991. The magnitude of global insect species richness. *Conservation Biology* 5: 283-296.

Gentry G, Dyer LA. 2002. On the conditional nature of neotropical caterpillar defenses against their natural enemies. *Ecology* 83: 3108-3119.

Downloaded From: https://complete.bioone.org/journals/Journal-of-Insect-Science on 28 Mar 2024
Terms of Use: https://complete.bioone.org/terms-of-use

Godfray HCJ. 1994. *Parasitoids: Behavioral and Evolutionary Ecology*. Princeton University Press

Godfray HCJ, Lewis OT, Memmott J. 1999. Studying insect diversity in the tropics. *Philosophical Transactions of the Royal Society of London Series B- Biological Sciences* 354: 1811-1824.

Guimarães JH. 1971. Family Tachinidae (Larvaevoridae). *A catalogue of the Diptera of the Americas south of the United States* 104: São Paulo. Museu de Zoologia, Universidade de São Paulo 333

Guimarães JH. 1977. Host-parasite and parasite-host catalogue of South American Tachinidae (Diptera). *Arquivos de Zoologia* 28(3): 1-131.

Hammer Ø, Harper DAT, Ryan PD. 2001. PAST: Paleontological statistics software package for education and data analysis. *Palaeontologia Electronica* 4: 1

Irwin ME, Schlinger EI, Thompson FC. In: Goodman SM, Benstead JP, editors. 2003. Diptera, true flies. *The Natural History of Madagascar*692-702. University of Chicago Press

Janzen DH, Pond CM. 1975. Comparison, by sweep sampling, of arthropod fauna of secondary vegetation in Michigan, England, and Costa Rica. *Transactions of the Royal Entomological Society of London* 127: 33-50.

Janzen DH. 1981. The peak in North American ichneumonid species richness lies between 38° and 42° N. *Ecology* 62: 532-37.

Janzen DH, Hallwachs W. 2006. Dynamic database for an inventory of the macrocaterpillar fauna, and its food plants and parasitoids, of the Area de Conservacion Guanacaste (ACG), northwestern Costa Rica. http://janzen.sas.upenn.edu.

Lasalle J, Gauld ID. 1991. Parasitic Hymenoptera and the biodiversity crisis. *Redia* 74: 315-334.

Longino JT, Coddington J, Colwell RK. 2002. The ant fauna of a tropical rain forest: estimating species richness three different ways. *Ecology* 83: 689-702.

Mittelbach GG, Schemske DW, Cornell HV, Allen AP, Brown JM, Bush MB, Harrison SP, Hurlbert AH, Knowlton N, Lessios HA, McCain CM, McCune AR, McDade LA, McPeek MA, Near TJ, Price TD, Ricklefs RE, Roy K, Sax DF, Schluter D, Sobel JM, Turelli M. 2007. Evolution and the latitudinal diversity gradient: speciation, extinction and biogeography. *Ecology Letters* 10: 315-331.

Novotny V, Basset Y, Miller SE, Weiblen GD, Bremer B, Cizek L, Drozd P. 2002. Low host specificity of herbivorous insects in a tropical forest. *Nature* 416: 841-844.

Odegaard F. 2000. How many species of arthropods? Erwin's estimate revised. *Biological Journal of the Linnaean Society* 71: 583-597.

O'Hara JE, Wood DM. 2004. Catalogue of the Tachinidae (Diptera) of America north of Mexico. *Memoirs Entomology International* 18: 1-410.

O'Hara JE. 2007; World genera of the Tachinidae (Diptera) and their regional occurrence. *Version* 3: http://www.nadsdiptera.org/Tach/Genera/Gentach ver2.pdf.

Pimm SL, Raven P. 2000. Biodiversity: Extinction by numbers. *Nature* 403: 843-845.

Rappé MS, Giovannoni SJ. 2003. The uncultured microbial majority. *Annual Review of Microbiology* 57: 369-394.

Roth J, King E, Thompson A. 1978. Host location behavior by a tachinid, *Lixophaga diatraeae*. *Environmental Entomology* 7: 794-798.

Sääksjärvi IE, Haataja S, Neuvonen S, Gauld ID, Jussila R, Salo J, Burgos AM. 2004. High local species richness of parasitic wasps (Hymenoptera : Ichneumonidae; Pimplinae and Rhyssinae) from the lowland rainforests of Peruvian Amazonia. *Ecological Entomology* 29: 735-743.

Smith MA, Woodley NE, Janzen DH, Hallwachs W, Hebert PDN. 2006. DNA barcodes reveal cryptic host-specificity within the presumed polyphagous members of a genus of parasitoid flies (Diptera : Tachinidae). *Proceedings of the National Academy of Sciences USA* 103: 3657-3662.

Smith MA, Wood DM, Janzen DH, Hallwachs W, Hebert PDN. 2007. DNA barcodes affirm that 16 species of apparently generalist tropical parasitoid flies (Diptera: Tachinidae) are not all generalists. *Proceedings of the National Academy of Sciences USA* 104: 4967-4972.

Stireman JO. 2008. alpha and beta diversity of a tachinid parasitoid community over space and time. *Annals of the Entomological Society of America* 101: 362-370.

Stireman JO, Singer MS. 2003a. What determines host range in parasitoids? An analysis of a tachinid parasitoid community. *Oecologia* 135: 629-638.

Stireman JO, Singer MS. 2003b. Determinants of parasitoid-host associations: Insights from a natural tachinid-lepidopteran community. *Ecology* 84: 296-310.

Stireman JO, Dyer LA, Janzen DH, Singer MS, Lill JT, Marquis RJ, Ricklefs RE, Gentry GL, Hallwachs W, Coley PD, Barone JA, Greeney HF, Connahs H, Barbosa P, Morais HC, Diniz IR. 2005. Climatic unpredictability and caterpillar parasitism: implications of global warming. *Proceedings of the National Academy of Sciences USA* 102: 17384-17387.

Stireman JO, O'Hara JE, Wood DM. 2006. Tachinidae: evolution, behavior, and ecology. *Annual Review of Entomology* 51: 525-555.

Strazanac JS, Plaugher CD, Petrice TR, Butler L. 2001. New tachinidae (Diptera) host records of eastern North American forest canopy Lepidoptera: Baseline data in a *Bacillus thuringiensis* variety *kurstaki* nontarget study. *Journal of Economic Entomology* 94: 1128-1134.

Townsend CHT. 1934–42. *Manual of Myiology in Twelve Parts*. Townsend and Filhos

Wood DM. 1985. A taxonomic conspectus of the Blondeliini of North and Central America and the West Indies (Diptera: Tachinidae). *Memoirs of the Entomological Society of Canada* 132: 1-130.

Wood DM. In: McAlpine JF, Peterson BV, Shewell GE, Teskey HJ, Vockeroth JR, Wood DM, editors. 1987. Chapter 110. Tachinidae. *Manual of Nearctic Diptera* 2: 1193-1269. Agriculture Canada Monograph 28-1269.

Wood DM. In preparation. In: Brown BV, Borkent A, Cumming JM, Wood DM, Woodley NE, Zumbado M, editors. Chapter 113 Tachinidae. *Manual of Central American Diptera*. National Research Council, Press, Ottawa, Canada.

Downloaded From: https://complete.bioone.org/journals/Journal-of-Insect-Science on 28 Mar 2024
Terms of Use: https://complete.bioone.org/terms-of-use

Woodley NE, Arnaud PH. 2008a. *Lobomyia neotropica*, a new genus and species of Tachinidae (Diptera) from the Neotropical Region. *Zootaxa* 1783: 31-39.

Woodley NE, Arnaud PH. 2008b. *Eulobomyia/, a new replacement name for* Lobomyia Woodley & Arnaud (Diptera: Tachinidae). *Zootaxa* 1856: 67-68.

Downloaded From: https://complete.bioone.org/journals/Journal-of-Insect-Science on 28 Mar 2024
Terms of Use: https://complete.bioone.org/terms-of-use