



## **Modified Artificial Diet for Rearing of Tobacco Budworm, *Helicoverpa armigera*, using the Taguchi Method and Derringer's Desirability Function**

Authors: Assemi, H., Rezapanah, M., Vafaei-Shoushtari, R., and Mehrvar, A.

Source: Journal of Insect Science, 12(100) : 1-18

Published By: Entomological Society of America

URL: <https://doi.org/10.1673/031.012.10001>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.



## Modified artificial diet for rearing of tobacco budworm, *Helicoverpa armigera*, using the Taguchi method and Derringer's desirability function

H. Assemi<sup>1a\*</sup>, M. Rezapana<sup>2b</sup>, R. Vafaei-Shoushtari<sup>1c</sup> and A. Mehrvar<sup>3d</sup>

<sup>1</sup>Department of Agricultural Entomology, Arak Branch, Islamic Azad university, Arak, Iran

<sup>2</sup>Center of Excellence for Organic Agriculture and Insect Virology Laboratory, Biocontrol Department, Iranian Research Institute of Plant Protection, Iran

<sup>3</sup>Department of Plant Protection, Azarbaijan Shahid Madani University-5375171379, East-Azarbaijan, Tabriz, Iran

### Abstract

With the aim to improve the mass rearing feasibility of tobacco budworm, *Helicoverpa armigera* Hübner (Lepidoptera: Noctuidae), design of experimental methodology using Taguchi orthogonal array was applied. To do so, the effect of 16 ingredients of an artificial diet including bean, wheat germ powder, Nipagin, ascorbic acid, formaldehyde, oil, agar, distilled water, ascorbate, yeast, chloramphenicol, benomyl, penicillin, temperature, humidity, and container size on some biological characteristics of *H. armigera* was evaluated. The selected 16 factors were considered at two levels (32 experiments) in the experimental design. Among the selected factors, penicillin, container size, formaldehyde, chloramphenicol, wheat germ powder, and agar showed significant effect on the mass rearing performance. Derringer's desirability function was used for simultaneous optimization of mass rearing of tobacco budworm, *H. armigera*, on a modified artificial diet. Derived optimum operating conditions obtained by Derringer's desirability function and Taguchi methodology decreased larval period from 19 to 15.5 days (18.42 % improvement), decreased the pupal period from 12.29 to 11 days (10.49 % improvement), increased the longevity of adults from 14.51 to 21 days (44.72 % improvement), increased the number of eggs/female from 211.21 to 260, and increased egg hatchability from 54.2% to 72% (32.84 % improvement). The proposed method facilitated a systematic mathematical approach with a few well-defined experimental sets.

**Keywords:** mass rearing, parameter optimization

**Correspondence:** <sup>a</sup> [hoda\\_assemi@yahoo.com](mailto:hoda_assemi@yahoo.com), <sup>b</sup> [rezapana@yahoo.com](mailto:rezapana@yahoo.com), <sup>c</sup> [r-vafaei@iau-arak.ac.ir](mailto:r-vafaei@iau-arak.ac.ir),

<sup>d</sup> [ac.mehrvar@azaruniv.ac.ir](mailto:ac.mehrvar@azaruniv.ac.ir). \* Corresponding author

**Received:** 26 June 2011, **Accepted:** 17 February 2012

**Copyright :** This is an open access paper. We use the Creative Commons Attribution 3.0 license that permits unrestricted use, provided that the paper is properly attributed.

**ISSN:** 1536-2442 | Vol. 12, Number 100

#### Cite this paper as:

Assemi H, Rezapana M, Vafaei-Shoushtari R, Mehrvar A. 2012. Modified artificial diet for rearing of tobacco budworm, *Helicoverpa armigera*, using the Taguchi method and Derringer's desirability function. *Journal of Insect Science* 12:100. Available online: <http://www.insectscience.org/12.100>

## Introduction

*Helicoverpa armigera* (Hübner) (Lepidoptera: Noctuidae) is a major pest affecting a wide range of economically important plants, including field and horticultural crops in Iran (Naseri et al. 2011) and throughout the world (Jallow et al. 2004; Reddy et al. 2004; Mironidis and Savopoulou- Soultani 2008; Yu et al. 2008). In northern Iran, tobacco plants are subjected to attack by several insect pests, the most serious of which is the budworm, *H. armigera*. Recently, it has been observed in increasing numbers. In tobacco cultivated fields, *H. armigera* predominantly damages young leaves and bolls, causing whole leaves to be destroyed during the crop's reproductive stage. The destruction of the leaves results in considerable loss of yield at harvest.

It is important to be able to economically rear important insects in order to study their life history, behavior, feeding habits, and susceptibility and resistance to chemical pesticides and biological control agents such as insect viruses (Rezapanah et al. 2008).

Rearing insects on artificial diets is an expensive process, and is difficult for developing countries where sufficient research funds are unavailable. As a result, the economic threats imposed by insect pests on agriculture will be poorly studied (Ahmed 1983).

Despite large expenditures for development of mass-rearing/artificial diet-based technology over the past five decades, successful commercialization of biological control systems based on such technology has been sparse. Also, many of the shortcomings of diet development are attributable to a failure to recognize the tremendous difficulty in the problems of diet and rearing system

development (Cohen 2003). Although diets are incredibly dynamic and complex mixes of multiple interactive functional components, diet development is often treated in the most simplistic manner (Cohen 2003). Cohen (2003) has explained that there are four equally important functional domains of diets: nutritional value, sensory qualities, availability, and stability. However, diet researchers often pay exclusive attention to the nutritional functions of diet components, without considering phagostimulation, textural factors (hydrocolloid characteristics, particle size, particle shape, viscosity, and other rheological matters), meta-nutritional antioxidant functions, anti-nutrients, bioavailability, water activity, antimicrobial potential, over-all stability, microbial deterioration, and numerous other features that must be juggled skillfully to achieve development of fully functional, successful diets (Cohen 2003). He has also explained that a most unfortunate confusion has developed regarding the terms “insect nutrition” and “insect dietetics,” with an even more unfortunate tacit assumption that nutrition is a “real science,” and dietetics an extension of fast food cooking. As a result of this confusion, and the prejudice against dietetic inquiries, researchers neglect most of the key issues that would explain how diets and certain diet components work or fail to work (Cohen 2003).

So far, various artificial diets have been developed and proposed for the maintenance and continuous rearing of economically important insects (Ahmed et al. 1998; Cohen, 2001; Castane and Zapata 2005). Although there is some success in efforts to rear successive generations of these insects entirely on an artificial diet, in many cases there is loss of both fitness and reproductive

potential, which causes longer developmental period and lower fecundity rate (Coudron et al. 2002). As a result, the cost-saving ratio is diminished. For those who work on life and fecundity tables of insects, it is important to know on what type of diet the insect has been reared. This factor affects the accuracy of biological characteristics of the insect, e.g. growth, viability, reproduction and population density, and consequently the accuracy of the life table (Bellows et al. 1992). Moreover, the shifting emphasis in insect control, using biological entities such as natural enemies and insect pathogens (bacteria, viruses and fungi), has increased the demand for constant reliable sources of such insects.

Laboratory-reared larvae can be used for the study of insect pathogens, plant resistance factors, effects of insecticides and radiation on fecundity and growth, as and insect life cycle (Ahmed et al. 1998). Beck et al. (1949) reported the first successful aseptic rearing of phytophagous moth, the European corn borer, *Ostrinia nubilalis* Hübner (Lepidoptera: Pyralidae), on a meridic diet. Ishii and Urushibara (1954) were able to rear the rice stem borer, *Chilo suppressalis* Walker (Lepidoptera: Crambidae) on a modification of the diet reported by Beck et al. (1949). Vanderzant and Reiser (1956) were the first to formulate a meridic diet for the pink bollworm, *Pectinophora gossypiella* Saunders (Lepidoptera: Gelechiidae), without a plant adjuvant. Later, Adkinsson et al. (1960) developed a satisfactory artificial medium supplemented with wheat germ for rearing pink bollworm. The wheat germ medium has been used to rear several species of plant-feeding lepidopterans, including the bollworm *Helicoverpa zea* Boddie (Lepidoptera: Noctuidae) (Vanderzant et al., 1962); the cabbage looper, *Trichoplusia ni* Hübner (Lepidoptera: Noctuidae) (Getzin, 1962); the

white large butterfly, *Pieris brassicae* Linnaeus (Lepidoptera: Pieridae) (David, 1965); and the codling moth, *Carpocapsa pomonella* Linnaeus (Lepidoptera: Tortricidae) (Rock et al., 1964).

Ahmed et al. (1998) reported that agar is a vital ingredient for insect-rearing diet, and is acquired from marine algae such as the *Gracilaria* and *Geladiella* species (Nene 1996). Abbasi et al. (2007) showed that when larval stage *H. armigera* were fed with a tapioca-based artificial diet, larval and pupal developmental period, percent pupating, pupal weight, emergence rate of male and female, longevity, fecundity, and hatching were non-significantly different from those of the control agar-based artificial diet.

Different models can be used for various design of experiments. For example, Hare (1992) used simple models to study trace metal contamination, and its effect on both distribution and abundance of aquatic insects. He found choosing a predictive model will facilitate the relationship between nature and laboratory data.

Lapointe et al. (2010) employed response surface methodology (RSM) in order to determine the effects of proportionality and amount of diet ingredients on larval survival, weight gain, and development rate in *Diaprepes abbreviatus* (L.). In his research, all measured larval responses led to the conclusion that cottonseed meal is the only major nutritive component, in addition to standard vitamin and salt mixes, required for a successful artificial diet for rearing larvae of *D. abbreviatus* to pupation. The response surface models demonstrated that larval survival to pupation and larval weight gain for *D. abbreviatus* are responses that can be independently manipulated. Diets that produce

large insects do not necessarily produce the greatest number of insects (Lapointe et al. 2008). Their experience in previous experimentation (Lapointe et al. 2008) was that survival of pupae to adult emergence was invariably close to 100%.

There are many statistical procedures used to design and analyze an experiment, but the most frequently used approach is a full factorial experiment. However, for full factorial experiments, there are  $2^f$  possible combinations that must be tested ( $f$  = the number of factors each at two levels). Therefore, it is very time-consuming when there are many factors involved (Ross, 1989). In order to minimize the number of required tests, fractional factorial experiments were developed. However, a shortcoming of this method is only a few of the total possible combinations are used in order to estimate the effects of main factors, and the effects of some of the interactions.

Taguchi developed a family of fractional factorial matrices that could be utilized in various situations. These matrices reduce the number of experiments, while providing valuable conclusions. The conclusions can be associated with a statistical level of confidence (Ryan 1988; Ross 1989; Peace 1992). The major steps of implementing the Taguchi method are: (1) identify the factors/interactions, (2) identify the levels of each factor, (3) select an appropriate orthogonal array, (4) assign the factors/interactions to columns of the orthogonal array, (5) conduct the experiments, (6) analyze the data and determine the optimal levels, and (7) conduct the confirmation experiment. The Taguchi method uses the signal to noise ratio to measure the quality characteristics deviating from the desired value. The signal to noise ratios differ

according to the type of characteristics. The "smaller the better" and "the bigger the better" responses are considered, aiming to minimize or maximize responses which are calculated using the following equations (Akhyar et al. 2008):

The smaller the better response category:

$$S/N = -10 \log \frac{1}{N} (\sum Y^2) \quad (1)$$

The bigger the better response category:

$$\frac{S}{N} = \frac{-10 \log \left( \frac{1}{y_1^2} + \frac{1}{y_2^2} + \frac{1}{y_3^2} + \dots + \frac{1}{y_n^2} \right)}{n} \quad (2)$$

where  $y_i$  is the characteristic properties: mass rearing,  $n$  is the replication number of the experiments.

In our study, we tried to determine the impact of the most critical ingredients in an artificial diet (such as agar, bean, ascorbic acid, ascorbate, formaldehyde, yeast, wheat germ powder, etc.), all of which have significant influence on the mass rearing of *H. armigera*. One strategy used in order to achieve the most effective performance is the optimization of operating parameters to create an optimum mass rearing of *H. armigera*. In order to establish better conditions by relating all the factors considered, numerous experiments need to be carried out with all the possible parameter combinations, which is not practical. Conventional experimental procedures involve altering of one factor at a time, and keeping all other factors constant, and assessing the impact of the individual factors. This procedure is, however, time consuming, requires more experimental sets, and is unable to provide information about how the individual factors interact with each other. Design of experiments is an approach

that helps to gain information about the optimized levels by taking a large number of variables into account. In our study, by performing Taguchi's method, and by making some modifications on a series of parameters, we were able to define the best combination of parameters for mass rearing feasibility of the *H. armigera* larvae.

## Materials and Methods

A colony of *H. armigera* was raised from two pairs of adults collected from a tobacco field at Tirtash Research and Education Center, Behshahr, Iran, and transferred to the laboratory. Then, it was reared on the agar-based diet (Ahmed et al. 1998), and used as a control. Experiments were conducted in the controlled laboratory conditions of  $65 \pm 5\%$  RH,  $27 \pm 2^\circ\text{C}$ , and 14:10 L:D. A homogenous stock of different larval instars, eggs, pupae, and adults were obtained from F1 generation for testing. The experiments were carried out with 16 factors at two levels, as shown in Table 1. The ingredients used in this study were selected according to Teakle and Jensen's work (1985) with an increase of about 14.28-50.00% in the quantities of all the components.

The dry ingredients of the diet were weighed carefully, and kept in separate containers. The wet ingredients were measured, and also kept in separate containers. The entire quantity of agar was suspended and brought to a boil. The total quantity of bean was added to the boiled agar. Then, all the dry and wet ingredients were added to this mixture. The prepared diet was then poured into the desired number of sterilized Petri plates, and allowed to cool and harden.

## Rearing procedures

**a) Petri plate for Individual larval development.** In our study, two standard-sized Petri plates (6 and 9 cm diameter, 1cm high) were used in the Taguchi design. Initially, these sterilized dishes containing food were infested with 20-30 newly hatched first instar larvae, with the help of a sterilized camel hair brush. Then, to prevent larval cannibalism, the larvae were divided into 10 sets, each comprising 30-50 dishes (one larva/dish).

**b) Adult emergence.** The larvae that were pupated in Petri plates were taken out and placed in the plastic dishes with a thin layer of soil at the bottom, and covered with muslin cloth. After emergence, the adults were collected individually into oviposition dishes. The emerged adults were checked and removed daily.

**c) Mating- oviposition cage.** 50 boxes ( $8 \times 15 \times 20$  cm) made of acrylic sheet were fabricated into an oviposition cage. A pair of adults were placed in each oviposition cage. These cages were covered with muslin cloth for ventilation. In order to provide food for the gravid females, several cotton wool pads, soaked in 10% sucrose solution, were placed on muslin cloth. The eggs deposited on the layers of the cotton wool oviposition pads were checked and removed daily, and kept in polyethylene bags for incubation and hatching.

**d) Egg incubation.** The eggs were incubated at two temperatures & humidity, i.e.,  $20^\circ\text{C}$  and  $25^\circ\text{C}$ , 60 % and 70% RH. The hatched larvae were then transferred into Petri plates containing their diet. The percent of egg hatching was recorded daily.

### Statistical analysis

In our study, the obtained experimental data was processed using Qualitek-4 (Nutek Inc. Bloomfield Hills, MI) software (Venkata-Mohan et al. 2007). Also, an analysis of variance (ANOVA) was applied to the data in order to conduct an analysis of the relative importance of each factor more systemically.

### Results

#### Mass rearing by employing Taguchi method

In Taguchi method, if the factors and interactions are more than 15, a possible matrix is a thirty two-trial orthogonal array, which is labeled as L32 matrix (Growley 1988; White 1991; Speidell et al. 1994). As seen in Table 2, mass rearing experiments were performed using selected 32 experimental trials, in combination of 16 factors at 2 levels. All sets of experiments, as per assigned factors, and the results obtained from signal to noise ratio for mass rearing using equations 1 and 2 in the case of "the bigger and the smaller, the better" response categories, are shown in Table 2. As seen in Table 2, some responses, such as average larval period, larval mortality, average pupal period, and pupal mortality, were supposed to be minimized, and the other responses, such as longevity of adults, the number of eggs laid, and egg hatchability, were considered to be maximized. Also, the raw data for average value of signal to noise ratio for each parameter at two levels are given in Table 3. The difference between values at levels 2 and 1 (L2-L1) of each factor indicates the relative influence of the effect. The larger the difference, the stronger the influence. The negative value has been ignored in assessing the main effect as the placement order of levels assigns either positive or negative values (Venkatan-Mohan et al. 2007).

ANOVA results for signal to noise ratio data with response characteristics are given in Table 4. The main objective of ANOVA is to extract from the results how much variation each factor causes relative to the result. The larger the variance of each factor, the stronger the influences on the results. According to the results of ANOVA (Table 4), the relative influences of these factors on "the bigger the better" response categories are in the order of: temperature > penicillin > ascorbate > Nipagin > formaldehyde > agar > oil > yeast > distilled water > ascorbic acid > humidity > container size > bean > wheat germ powder > chloramphenicol > benomyl and in the case of "the smaller the better" responses are in order of yeast > ascorbic acid > ascorbate > container size > benomyl > formaldehyde > penicillin > wheat germ powder > water sterile > agar > oil > bean > humidity > Nipagin > temperature > chloramphenicol.

#### Optimum levels and estimation of optimum response characteristics

Table 5 indicates that, in the case of "the bigger the better" responses, the first level of agar, bean, ascorbate, yeast, formaldehyde, distilled water, chloramphenicol, wheat germ powder, penicillin, humidity, container size, and the second level of other factors has a large signal to noise ratio (28.40). In the case of "the smaller the better" response categories, the first level of bean, ascorbic acid, ascorbate, yeast, formaldehyde, distilled water, Nipagin, benomyl, temperature, humidity, and the second level of other factors has a small signal to noise ratio (-20.00). For simultaneous optimization of the two different sets of responses (the smaller, the better, and the bigger, the better) partial desirability ( $d_i$ ) and global desirability (D) functions were used, and are described in the following section.

### Partial Desirability ( $d_i$ ) and Global Desirability (D) Functions

The procedure followed in this work for simultaneous optimization of two different sets of responses (the smaller, the better, and the bigger, the better) is a method developed by Derringer and Suich (1980). The method involves transformation of each predicted response,  $\hat{y}$ , to a dimensionless partial desirability function,  $d_i$ , which includes the researcher's priorities and desires when building the optimization procedure. One or two-sided functions are used, depending on whether each of the responses has to be maximized or minimized, or has an allotted target value. If the response  $i$  is to be maximized, the quantity  $d_i$  is defined as:

$$d_i = \left[ \frac{\hat{y} - A}{B - A} \right]^{w_i} \quad \text{when } A \leq \hat{y} \leq B, \quad (3)$$

$$d_i = \left[ \frac{\hat{y} - A}{B - A} \right]^{w_i}$$

$$d_i = 1, \hat{y} > B \text{ and } d_i = 0, \hat{y} < A$$

Likewise,  $d_i$  can be defined when the response is to be minimized, or if there is a target value for the response. In Eq. (3),  $A$  and  $B$  are, respectively, the lowest and the highest values obtained for the response  $i$ , and  $w_i$  is the weight.  $d_i$  ranges between 0, for a completely undesired response, and 1, for a fully desired response. In both cases,  $d_i$  will vary non-linearly while approaching the desired value. But with a weight of 1,  $d_i$  varies linearly. In our work, we chose weights equal to 1 for all six responses. The partial desirability functions are then combined into a single composite response, the so-called global desirability function  $D$ , defined as the geometric mean of the different  $d_i$ -values:

$$D = \left[ d_1^{p_1} \times d_2^{p_2} \times d_3^{p_3} \times \dots \times d_n^{p_n} \right]^{\frac{1}{n}} \quad (4)$$


A value of  $D$  different from zero implies that all responses are in a desirable range simultaneously, and, consequently, for a value of  $D$  close to 1, the combination of the different criteria is globally optimum, so the response values are near the target values. In Eq. (4),  $p_i$  is the relative importance assigned to the response  $i$ . The relative importance of  $p_i$  is a comparative scale for weighing each of the resulting  $d_i$  in the overall desirability product, and it varies from the least important ( $p_i = 1$ ) to the most important ( $p_i = 5$ ). It is noteworthy that the outcome of the overall desirability  $D$  depends on the  $p_i$  value that offers users flexibility in the definition of desirability functions. According to Table 6, in row 1,  $Y_{min}$  (average larval period, larval mortality, average pupal period and pupal mortality) and  $Y_{max}$  (longevity of adults, the number of eggs laid and percent hatched of eggs) were measured 9.87 and 94.96, respectively. Also, according to equations 3 and 4,  $d_{min}$ ,  $d_{max}$  and  $D$  were defined 0.1, 0.73 and 0.14, respectively. Then, the obtained global desirability functions ( $D$ ) are used as new responses of Taguchi design, and the other steps of analysis were again carried out by Qualitek-4 software. In this case, predicted results are shown in Table 7. So, It can be concluded that in simultaneous optimization of two different response categories (the smaller, the better, and the bigger, the better), the first level of container size, yeast, oil, chloramphenicol, Nipagin, benomyl, penicillin, agar, and the second level of the other factors have important roles.

According to the final analysis of variance (Table 8), the  $F$ -ratio indicates the variation in the response can be explained by the regression. The  $p$  value is used to estimate whether  $F$  is large enough to indicate statistical significance. If  $p$  value is lower than 0.05, it indicates the factor is statistically



significant (Segurola et al. 1999). The larger the variance of each factor, the stronger the influence on the results. Hence, from the results of the final ANOVA displayed in Table 8, the order of relative influence of these selected factors on yield of mass rearing in the case of simultaneous optimization of two different set of responses (the smaller, the better, and the bigger, the better) are as follows: penicillin > bean > container size > formaldehyde > agar > ascorbate > chloramphenicol > wheat germ powder > benomyl > distilled water > temperature > humidity > oil > Nipagin > yeast > ascorbic acid. As seen in Table 8, the results showed that, in comparison with other factors, penicillin, bean, container size, formaldehyde, agar, and ascorbate are significant ( $p$  value is lower than 0.05). For further investigation, six final significant factors (with  $P$  value less than

0.05) in our final artificial diet formula (Table 8) were chosen as a control. Then, a new ANOVA table was provided, with these six main factors, three levels (high, medium and low) and, keeping the other factors constant (Table 9, 10). As shown in Table 10, these 6 factors were meaningful at these three levels as well ( $p < 0.05$ ). Moreover, in order to be sure of the meaningfulness of the primary selected two levels, the Pair  $t$ -test was used. In our test, when  $t$ -calculated >  $t$ -critical, it indicates that there is a significant difference between the levels. Table 11 expresses this claim. Eventually, it is concluded that, based on Tables 8, 10 and 11, these results are in good agreement with each other. In other words, the main factors are the meaningful factors, and there is a significant difference between the selected levels.



**Figure 1.** Plots of Individual factors performance at different levels for optimization of modified artificial diet. High quality figures are available online.

The response curves for the individual effects of parameters on average value of responses are given in Figure 1. In Figure 1, an increase in the levels of factors such as bean, ascorbate, formaldehyde, distilled water, wheat germ powder, humidity, and temperature resulted in an increase in the response values. This is discussed further below.

Two conformation experiments were performed at selected optimal levels of the process parameter. The predicted result obtained for the final artificial diet is  $D=0.789$ , with a confidence level of  $\pm 0.148$  via Taguchi method. It means that if our confirmatory experiment ( $\mu$ ) is  $0.641 < \mu < 0.937$ , we can conclude that predicted and confirmatory experiment results are in agreement with each other. In our study, confirmatory experiment was obtained  $\mu=0.916$ . Note that this result is valid within the specified range. Table 12 shows the comparison of confirmatory experiments and predicted values of different objective functions under the optimum conditions, and it can be concluded that they are in good agreement with each other.

## Discussion

This investigation revealed that using Taguchi and Derringer's Desirability Function approach facilitated the process of optimization by understanding the role of factors involved in the ingredients for the preparation of diet. The final results showed, in comparison with other parameters, penicillin (Level 1, 2 g), bean (Level 2, 305 g), container size (Level 1; 6×1 cm), formaldehyde (Level 2; 1 cc), agar (Level 1; 14 g), and ascorbate (Level 2; 5 g) were the most effective factors for the yield of mass rearing (Table 8). An increase in the amount

of bean, ascorbate, formaldehyde 37%, and wheat germ powder in the artificial diet resulted in higher mass rearing (Figure 1), because these factors acted as nutrients for insects, and infection reducers for diet.

The results showed a significant decrease in yield of mass rearing with increasing agar, penicillin, and container size. The maximum effect of penicillin for reduction of fungal contamination was obtained at level 1 (2 g), whereas subsequent increases in this factor resulted in the reverse effect due to mortality of larvae. Small size of container can provide desired growth of insect, and even decreased fungal and bacterial contaminations.


According to Figure 1, the highest yield of mass rearing was obtained at the smaller amount of agar (level 1, 14 g), which is in agreement with the results of other studies on artificial diet showing its economical usefulness for mass rearing (Shorey and Hall 1965; Ahmed 1983; Abbasi et al. 2007). In our study other factors had no significant effects on optimization of production. Table 8 shows that, in comparison with other factors, penicillin, bean, container size, formaldehyde, agar, and ascorbate are significant ( $p < 0.05$ ), and we should be careful about amounts of these components. However, it does not mean that the other factors were non-influential; rather, all the factors together help to create the best modified artificial diet. Hence, it can be concluded that final artificial diet in first order are penicillin (Level 1; 2 g), bean (Level 2; 305 g), container size (Level 1; 6×1 cm), formaldehyde (Level 2; 1 cc), agar (Level 1; 14 g) and ascorbate (Level 2; 5 g) that were significant factors, following chloramphenicol (Level 1; 2 cc), wheat germ powder (Level 2; 45 g), benomyl (Level 1; 3 g), distilled water (Level 2; 1000 cc), temperature (Level 2; 25 °C), humidity (Level 2, 70%), oil (Level 1; 5

cc), Nipagin (Level 1; 3 g), yeast (Level 2; 50 g) and ascorbic acid (Level 2; 2.2 g).

Moreover, the experiments carried out under optimized conditions showed that growth was enhanced. The process variation at improved and current conditions, with the function of frequency distribution, is shown in Figure 2. It can be concluded that a substantial increase in the frequency distribution was observed with the optimal factors. Therefore, it can be concluded that our design was successful.

Derived optimum operating conditions obtained by the Taguchi methodology and Deringer's desirability functions showed effective yield of mass rearing of tobacco budworm so that the larval and pupal periods decreased from 19 to 15.5 days (over all 18.42% improvement), and from 12.29 to 11 days (over all 10.49 % improvement), respectively at 20-25 °C. Under the same conditions, however, Ahmed et al (1998) reported that the average larval and pupal periods were 22.9 and 17.1 days. In this study, the longevity of adult was enhanced from 14.51 to 21 days (over all 44.72 % improvement). The number of eggs/female increased from 211.21 to 260 eggs and egg hatchability increased from 54.2 to 72% (over all 32.84 % improvement).

Compared with what Abbasi et al. (2007) found, our results show the impact of the combined ingredients of different levels on longevity of adult, number of eggs/ female, and egg hatchability, which were 15.03 days, 201%, and 59.01%, respectively. These results are in agreement with Gelman et al. (2001). They found out the highest egg hatchability for leaf-fed beetles fed the Lettuce diet ( $70.0 \pm 8.0\%$ ), and lowest for beetles fed the Potato diet ( $35.4 \pm 3.6\%$ ). Also, Hamed and Nadeem (2008) studied seven artificial diets in



laboratory conditions. They prepared a diet using basic ingredients such as flour of chickpea, mungbean, soybean, wheat, maize, cotton seed, and water chestnut. They reported that larval duration of *H. armigera* was shortest (14.5 days) in the chickpea flour diet, but was longest (42.8 days) in the water chestnut flour diet. In other diets, larval duration was 15.3, 15.5, 15.6, 16.0, 16.5 and 21.3 days on mungbean, soybean, chickpea leaves and pods (control), wheat, maize, and cotton seed, respectively. It was concluded that larval duration (15.5 days) in our study was close to larval duration of soybean, chickpea leaves and pods (control), wheat, maize diets in Hamed and Nadeem's (2008) study. Average larval and pupal mortality in our findings were 4.4% and 7.7%, respectively. Hamed and Nadeem (2008) suggested that the minimum mortality was on chickpea flour (1.1%), while maximum mortality was on cotton seed flour (21.0%). Based on our finding, pupal duration (11.0 days) was equal to the chickpea flour diet investigated by Hamed and Nadeem (2008).

In conclusion, the Taguchi optimization methodology provided a systematic and efficient mathematical approach to

understanding complex processes for optimization of mass rearing parameters with only a few well-defined experimental sets. Hence, this diet is recommended as a cost-effective production for rearing of *H. armiger*. Note that our findings from this work are relevant to the two test levels used, and could give different results at other test levels, or at greater extremes than tested in this experiment.

## Acknowledgements

The authors would like to thank Ms. Seyed Afshin Sajjadi and Dr. Majid Ghaninia for assistance and comments on the first draft of the manuscript. The financial support for this investigation, given by Tirtash Reaserch and Education Center Behshahr, Iran, is gratefully acknowledged.

## References

- 
- Abbasi BH, Ahmed K, Khalique F, Ayub N, Liu HJ, Kazami SAR, Nauman M. 2007. Rearing the cotton bollworm, *Helicoverpa armigera*, on a tapioca-based artificial diet. *Journal of Insect Science* 7:35. Available online, <http://insectscience.org/7.35>
- Adkinsson PL, Vanderzant ES, Bull DL, Allison WE. 1960. A wheat germ medium for rearing the pink bollworm. *Journal of Economic Entomology* 53(5): 759-762.
- Ahmed K. 1983. *Development of artificial diets for mass breeding of Lepidopterous insects*. M. Phil. Thesis, University of Karachi, Karachi Pakistan.
- Ahmed K, Khalique F, Malik BA. 1998. Modified artificial diet for mass rearing of Chickpea Pod borer, *Helicoverpa armigera* (H.). *Pakistan Journal of Biological Sciences* 1: 183-187.
- Akhyar G, Che Haron CH, Ghani JA. 2008. Application of Taguchi method in the optimization of turning parameters for surface roughness. *International Journal of Science Engineering and Technology* 1(3): 60-66.
- Beck SD, Lilly JH, Stauffer JF. 1949. Nutrition of European corn borer, *Pyrausta nubilais* (Hubn.). Development of satisfactory purified diet for larval growth. *Annals of the Entomological Society of America* 42: 483-496.
- Bellows TS, Van Driesche RG, Elkinton JS. 1992. Life table construction and analysis in the evaluation of natural enemies. *Annual Review in Entomology* 37: 587-614.
- Castane C, Zapata R. 2005. Rearing the predatory bug *Macrolophus caliginosus* on a meat based diet. *Biological Control* 34: 66-72.
- Cohen AC. 2001. Formalizing insect rearing and artificial diet technology. *American Entomologist* 47: 198-206
- Cohen AC. 2003. *Insect Diets: Science and Technology*. pp. 429. CRC Press.
- Coudron TA, Wittmeyer J, Kim Y. 2002. Life history and cost analysis for continuous rearing of *Podisus maculiventris* (Heteroptera: Pentatomidae) on a Zoophytophagous artificial diet. *Journal of Economic Entomology* 95: 1159-1168.
- Derringer G, Suich R. 1980. Simultaneous Optimization of Several Response ariables. *Journal of Quality Technology* 12: 214-219.

- Gelman DB, Bell RA, Liska LJ, Hu JS. 2001. Artificial diet for rearing the Colorado potato beetle, *Leptinotarsa desmolineata*. *Journal of Insect Science* 1:7. Available online, <http://insectscience.org/1.7>.
- Getzin, LW. 1962. Mass rearing of virus free Cabbages looppers on an artificial diet. *Journal of Insect Pathology* 4: 486- 487.
- Growley RT. 1986. Laser fabrication of photomasks for hybrid circuits, *Proceedings of SPIE* 611: 18-22.
- Hare L. 1992. Aquatic insects and trace metals: bioavailability, bioaccumulation, and toxicity. *Critical Reviews in Toxicology* 22: 327-369.
- Jallow MFA, Cunningham JP, Zalucki MP. 2004. Intra-specific Variation for Host Plant Use in *Helicoverpa armigera* (Hübner) (Lepidoptera: Noctuidae): implications for management. *Crop Protection* 23: 955-964.
- Hamed M, Nadeem S. 2008. Rearing of *Helicoverpa armigera* (Hub.) on Artificial Diets in Laboratory. *Pakistan Journal of Zoology* 40(6): 447-450.
- Ishii S, Urushibara H. 1954. On fat soluble and water soluble growth factors required by the rice stem borer, *Chilo simplex* Buttl. *Bulletin of the National Institute of Agricultural Sciences Series C* 4: 109-133.
- Lapointe SL, Evens TJ, Niedz RP. 2008. Insect diets as mixtures: optimization for a polyphagous weevil. *Journal of Insect Physiology* 54: 1157-1167.
- Lapointe R, Stephen LP, Niedz RP, Evens TJ. 2010. An artificial diet for Diaprepes abbreviatus (Coleoptera: Curculionidae) optimized for larval survival. *Florida Entomologist* 93(1): 56-62.
- Mironidis, G. K. and Savopoulou-Soultani, M. 2008. Development, Survivorship and Reproduction of *Helicoverpa armigera* (Lepidoptera: Noctuidae) under Constant and Alternating Temperatures. *Environmental Entomology* 37:16-28.
- Naseri B, Fathipouri Y, Moharramipour S, Hosseiniaveh V. 2011. Comparative Reproductive Performance of *Helicoverpa armigera* (Hübner) (Lepidoptera: Noctuidae) Reared on Thirteen Soybean Varieties. *Journal of Agricultural Science and Technology* 13: 17-26.
- Nene YL. 1996. Tapioca a potential substitute for agar in tissue culture media. *Current Science* 70: 493-494.
- Peace GS. 1992. *Taguchi Methods: a Hands-on Approach*, Addison-Wesley Publishing Company, Inc.
- Reddy KS, Rao GR, Rao PA, Rajasekhar P. 2004. Life Table Studies of the Capitulum Borer, *Helicoverpa armigera* (Hübner) Infesting Sunflower. *Journal of the Entomological Research Society* 28:13-18.
- Rezapanah M, Shojai-Estrabragh S, Jehle JA, Huber J. 2008. Molecular and biological characterization of new isolates of *Cydia pomonella* granulovirus (CpGV) from Iran. *Journal of Pest Science* 81: 187-191.
- Rock GC, Glass EH, Patton RL. 1964. Axenic rearing of the redbanded leaf roller *Argyrotaenia velutinana* on meridic diet. *Annals of the Entomological Society of America* 57: 617-621.



Ross R. 1989. *Taguchi Techniques for Quality Engineering*. McGraw-Hill.

Ryan N E. 1988. *Taguchi Methods and QFD: Hows and Whys for Management*. ASI Press.  
Segurola J, Allen NS, Edge M, Mahon AM. 1999. Design of eutectic photo initiator blends for UV/curable acrylated printing inks and coating. *Progress in Organic Coatings* 37: 23-37.

Shorey HH, Hale RL. 1965. Mass rearing of the larvae of nine noctuid species on a simple artificial medium. *Journal of Economic Entomology* 58(3): 522-524.

Speidell JL, Cordes SA, Ferry A. 1994. An improved excimer laser pattern generator for photomask fabrication. *Proceedings of SPIE* 2322: 66 -78.

Taguchi G. 1987. *System of Experimental Design*. UNIPUB Kraus, International Publications.

Teakle RE, Jensen JM. 1985. *Heliothis armigera*. In: Singh P, Moore RF, Editors. *Handbook of insect rearing* 2: 313-322. Elsevier.

Vanderzant E S, Rieser R. 1956. Studies on the nutrition of the pink bollworm using purified casein media. *Journal of Economic Entomology* 49: 454-458.

Vanderzant E S, Rechardson CD , Fort SW. 1962. Rearing of bollworm on artificial diet . *Journal of Economic Entomology* 55: 140.

Venkata-Mohan S, Sirisha K, Sreenivasa R, Sarma PN. 2007. Bioslurry phase remediation of chlorpyrifos contaminated soil: Process evaluation and optimization by Taguchi design of experimental (DOE) methodology.

*Ecotoxicology and Environmental Safety* 68: 252-262.

WAL David , Gardiner BOC. 1965. Rearing *Peiris brassicae* L. larvae on a semi-synthetic diet . *Nature* 207: 882-883.

White, A. T. 1991. Evolution of the photo mask industry, *Proceedings of SPIE* 1604: 2 - 23.

Yu, F. L., Wu, G., Liu, T. J., Zhai, B. P. and Chen, F. J. 2008. Effects of Irrigation on the Performance of Cotton Bollworm, *Helicoverpa armigera* (Hübner) during Different Pupal Stages. *International Journal of Pest Management* 54(2): 137-142.

**Table 1.** Factors and their levels chosen for optimization of modified artificial diet using Taguchi method.

Factors	Levels	
	1	2
bean (g)	205	305
wheat germ powder (g)	30	45
Nipagin (g)	2.2	4.4
ascorbic acid (g)	1.1	2.2
formaldehyde 37% (cc)	0.5	1
oil (cc)	5	10
agar (g)	14	21
water sterile (cc)	700	1000
ascorbate (g)	3.5	5
yeast (g)	35	50
chloramphenicol (cc)	2	3
benomyl (g)	3	4.5
penicillin (g)	2	4
temperature (°C)	20	25
humidity (%)	60	70
container size (cm)	6 × 1	9 × 1

**Table 2.** Column assignment for various factors in the Taguchi's L<sub>32</sub> orthogonal array (OA) and experimental values for optimization of modified artificial diet.

Exp. run	Factors																Responses								
	A	B	C	D	E	F	G	H	I	J	K	L	M	N	O	P	Ymin (the smaller the better)					Ymax(the bigger the better)			
																	Larval period	Larval mortality	Pupal period	Pupal mortality	S/N Ratio	Longevity of adult	N. of eggs/	% hatched of eggs	S/N Ratio
1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	16.2	5.1	10.1	8.1	-20.57	14.8	211	59	27.89
2	1	1	1	1	1	1	1	1	2	2	2	2	2	2	2	2	17.3	4.8	11.2	7.4	-20.99	14.9	200	55	27.9
3	1	1	1	1	2	2	2	2	1	1	1	1	2	2	2	2	21.2	3.2	12.3	7	-22.17	15.2	216	54	28.05
4	1	1	1	1	2	2	2	2	2	2	2	2	2	1	1	1	16.4	5.1	13	8.3	-21.24	14.3	215	58	27.6
5	1	1	2	2	1	1	2	2	1	1	2	2	1	1	2	2	17.4	5.2	10.2	8.5	-21.02	15	200	57	27.97
6	1	1	2	2	1	1	2	2	2	2	1	1	2	2	1	1	18.1	4.8	11	7.4	-21.19	14.5	210	55	27.68
7	1	1	2	2	2	2	1	1	1	1	2	2	2	2	1	1	23.2	4.7	13	6.9	-22.88	14.4	214	50	27.57
8	1	1	2	2	2	2	1	1	2	2	1	1	1	1	2	2	21.4	5.3	12.2	8.1	-22.43	13.9	221	54	27.33
9	1	2	1	2	1	2	1	2	1	2	1	2	1	2	1	2	19.6	3.4	12.3	8	-21.84	14	220	58	27.43
10	1	2	1	2	1	2	1	2	2	1	2	1	2	1	2	1	17.9	3.4	11.9	7.8	-21.25	14.5	211	55	27.68
11	1	2	1	2	2	1	2	1	1	2	1	2	2	1	2	1	19.2	5.5	12.1	7.5	-21.77	13.5	213	59	27.13
12	1	2	1	2	2	1	2	1	2	1	2	1	1	2	1	2	21.9	4.8	13.1	8.4	-22.7	15.2	215	53	28.04
13	1	2	2	1	1	2	2	1	1	2	2	1	1	2	2	1	22.1	5.3	14.2	8	-22.91	15.3	219	58	28.15
14	1	2	2	1	1	2	2	1	2	1	1	2	2	1	1	2	18	4.1	13.9	7.7	-21.71	14.2	200	50	27.45
15	1	2	2	1	2	1	1	2	1	2	2	1	2	1	1	2	17.9	3.4	12.8	8.2	-21.48	14.5	205	53	27.66
16	1	2	2	1	2	1	1	2	2	1	1	2	1	2	2	1	23.1	4.1	10.2	8.5	-22.59	15.3	211	54	28.1
17	2	1	1	2	1	2	2	1	1	2	2	1	2	1	1	2	16.9	5.1	11.8	8.8	-21.2	13.8	211	52	27.25
18	2	1	1	2	1	2	2	1	2	1	1	2	1	2	2	1	21.2	3.4	12.2	7.9	-22.25	14	215	56	27.41
19	2	1	1	2	2	1	1	2	1	2	2	1	1	2	2	1	18.2	5.1	13.1	7.5	-21.65	15	217	54	27.95
20	2	1	1	2	2	1	1	2	2	1	1	2	2	1	1	2	20.2	4.2	13.5	8.1	-22.26	14.3	210	53	27.55
21	2	1	2	1	1	2	1	2	1	2	1	2	2	2	1	2	19	5.1	14	8.6	-22.15	14.6	215	50	27.68
22	2	1	2	1	1	2	1	2	1	2	2	1	1	2	1	2	16.4	5	11.2	8	-20.82	14.9	200	58	27.93
23	2	1	2	1	2	1	2	1	1	2	1	2	1	2	1	2	17.4	4.9	14.1	7.4	-21.61	15.2	202	54	28.05
24	2	1	2	1	2	1	2	1	2	1	2	1	2	1	2	1	20.4	3.4	13.9	7.1	-22.24	13.8	216	51	27.24
25	2	2	1	1	1	1	2	2	1	1	2	2	2	2	1	1	18.4	4.5	12.4	6.2	-21.39	15.1	220	50	27.95
26	2	2	1	1	1	1	2	2	2	2	1	1	1	1	2	2	16.2	3.2	11.8	7.2	-20.64	14	214	59	27.43
27	2	2	1	1	2	2	1	1	1	1	2	2	1	1	2	2	19	5.1	12.3	7.4	-21.71	14.2	200	54	27.5
28	2	2	1	1	2	2	1	1	1	2	2	1	1	2	2	1	16	5.3	10.5	6.3	-20.35	14.5	210	50	27.62
29	2	2	2	2	1	1	1	1	1	1	1	1	2	2	2	2	15	4.9	11.2	8.4	-20.46	13.9	212	53	27.32
30	2	2	2	2	1	1	1	1	2	2	2	2	1	1	1	1	23	3.9	11.8	7.6	-22.67	15.2	213	55	28.06
31	2	2	2	2	2	2	2	2	2	1	1	1	1	1	1	1	22	4.2	14.1	8.4	-22.85	14	219	54	27.39
32	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	20	3.9	12	7.6	-21.88	14.4	204	50	27.57

**Table 3.** Response (average value of S/N ratio) for Taguchi analysis for "the bigger, the better responses" (average longevity of adults, the number of eggs laid and egg hatchability) and "the smaller the better responses" (average larval period, larval mortality, average pupal period and pupal mortality).

No.	Factors	The bigger, the better			The smaller, the better		
		Level 1	Level 2	L2- L1	Level 1	Level 2	L2- L1
1	agar	27.73	27.62	-0.1	-21.8	-21.63	0.61
2	bean	27.69	27.66	-0.03	-21.67	-21.76	-0.09
3	ascorbic acide	27.65	27.7	0.49	-21.5	-21.93	-0.43
4	ascorbate	27.76	27.58	-0.18	-21.53	-21.89	-0.35
5	yeast	27.7	27.65	-0.05	-21.44	-21.99	-0.54
6	formaldehyde	27.75	27.6	-1.45	-21.58	-21.85	-0.27
7	distilled water	27.7	27.65	-0.05	-21.63	-21.8	-0.16
8	oil	27.72	27.73	0.1	-21.78	-21.65	0.12
9	chloramphenicol	27.68	27.66	-0.02	-21.73	-21.7	0.02
10	wheat germ powder	27.69	27.66	-0.03	-21.8	-21.62	0.17
11	Nipagin	27.6	27.75	0.15	-21.68	-21.75	-0.07
12	benomyl	27.66	27.68	0.01	-21.56	-21.87	-0.31
13	penicillin	27.76	27.58	-0.18	-21.84	-21.59	0.25
14	temperature	2.55	27.79	0.24	-21.7	-21.73	-0.02
15	humidity	27.7	27.65	-0.04	-21.67	-21.76	-0.08
16	container size	27.69	27.65	-0.04	-21.87	-21.56	0.31



**Table 4.** Analysis of variance (ANOVA) for the "bigger, the better responses" (average longevity of adults, the number of eggs laid and egg hatchability) and "the smaller the better responses" (average larval period, larval mortality, average pupal period and pupal mortality).

No.	Factors	DOF (f)	The bigger, the better			The smaller, the better		
			Sum of Sqr. (S)	Variance (V)	F - Ratio (F)	Sum of Sqr. (S)	Variance (V)	F - Ratio (F)
1	agar	1	0.094	0.094	1.39	0.21	0.21	0.36
2	bean	1	0.009	0.009	0.134	0.073	0.073	0.125
3	ascorbic acide	1	0.018	0.018	0.276	1.499	1.499	2.567
4	ascorbate	1	0.257	0.257	3.775	1.027	1.027	1.759
5	yeast	1	0.019	0.019	0.287	2.393	2.393	4.098
6	formaldehyde	1	0.165	0.165	2.427	0.609	0.609	1.044
7	distilled water	1	0.019	0.019	0.288	0.223	0.223	0.382
8	oil	1	0.09	0.09	1.32	0.13	0.13	0.223
9	chloramphenicol	1	0.003	0.003	0.044	0.004	0.004	0.008
10	wheat germ powder	1	0.007	0.007	0.111	0.255	0.255	0.437
11	Nipagin	1	0.194	0.194	2.844	1.045	1.045	0.078
12	benomyl	1	0.002	0.002	0.039	0.793	0.793	1.359
13	penicillin	1	0.265	0.265	3.884	0.527	0.527	0.903
14	temperature	1	0.471	0.471	6.903	0.007	0.007	0.012
15	humidity	1	0.016	0.016	0.242	0.057	0.057	0.098
16	container size	1	0.015	0.015	0.221	0.795	0.795	1.363

**Table 5.** Predicted Optimum Conditions and Contribution for four values of mass rearing using "the bigger, the better" responses (average longevity of adults, the number of eggs laid and egg hatchability) and the "smaller the better" responses (average larval period, larval mortality, average pupal period and pupal mortality) via Taguchi method.

No.	Factors	The bigger, the better			The smaller, the better		
		Level Description	Level	Contribution	Level Description	Level	Contribution
1	agar	14	1	0.05	21	2	0.08
2	bean	205	1	0.01	205	1	0.04
3	ascorbic acide	2.2	2	0.02	1.1	1	0.21
4	ascorbate	3.5	1	0.08	3.5	1	0.17
5	yeast	35	1	0.02	35	1	0.27
6	formaldehyde	0.5	1	0.07	0.5	1	0.13
7	distilled water	700	1	0.02	700	1	0.08
8	oil	10	2	0.05	10	2	0.06
9	chloramphenicol	2	1	0.01	3	2	0.01
10	wheat germ powder	30	1	0.01	45	2	0.08
11	Nipagin	4.4	2	0.07	2.2	1	0.03
12	benomyl	4.5	2	0.09	3	1	0.15
13	penicillin	2	1	0.09	4	2	0.12
14	temperature	25	2	0.12	20	1	0.01
15	humidity	60	1	0.02	60	1	0.04
16	container size	6	1	0.02	9	2	0.15
Expected Result At Optimum Condition		28.4			-20		



**Table 6.** Partial ( $d_i$ ) and global (D) desirability Functions for analysis of seven values of mass rearing (average longevity of adults, the number of eggs laid, egg hatchability, larval period, larval mortality, average pupal period and pupal mortality).

No.	Average $Y_{\min}^a$		Average $Y_{\max}^b$		$d_{i\min}$		$d_{i\max}$		D		Weight
	I	II	I	II	I	II	I	II	I	II	
1	9.875	11.62	94.96	96.13	0.1	0.73	0.73	0.86	0.14	0.24	1
2	10.175	10.25	89.93	90.46	0.23	0.25	0.18	0.24	0.003	0.006	1
3	10.65	10.92	95.13	96.4	0.39	0.5	0.75	0.89	0.17	0.34	1
4	10.7	11.72	95.93	95.26	0.41	0.82	0.84	0.76	0.2	0.4	1
5	10.325	10.7	90.83	95.93	0.28	0.41	0.28	0.84	0.01	0.2	1
6	10.325	10.65	93.2	95.13	0.28	0.39	0.54	0.75	0.05	0.17	1
7	11.95	10.7	92.73	95.93	0.84	0.42	0.49	0.84	0.1	0.24	1
8	11.75	12.32	96.26	96.26	0.77	0.97	0.87	0.87	0.52	0.67	1
9	10.825	11.97	97.4	97.4	0.45	0.85	1	1	0.44	0.84	1
10	10.25	9.5	93.6	94.96	0.25	0	0.58	0.73	0.06	0.14	1
11	11.075	10.92	95.2	97.33	0.54	0.49	0.76	0.99	0.26	0.46	1
12	12.05	12.1	94.5	96.24	0.87	0.89	0.68	0.87	0.31	0.61	1
13	12.4	12.4	97.4	97.4	1	1	1	1	1	1	1
14	10.925	10.92	88.2	88.2	0.49	0.49	0	0	0	0	1
15	10.65	11.95	90.93	92.73	0.39	0.84	0.29	0.49	0.001	0.1	1
16	11.475	11.17	93.4	95.16	0.68	0.57	0.56	0.75	0.15	0.26	1
17	10.65	11.95	92.3	92.73	0.39	0.84	0.44	0.49	0.03	0.1	1
18	11.175	10.92	95.16	97.33	0.57	0.49	0.75	0.99	0.26	0.46	1
19	10.975	10.92	95.43	97.33	0.5	0.49	0.78	0.99	0.26	0.46	1
20	11.5	11.95	92.5	92.73	0.68	0.84	0.46	0.49	0.09	0.1	1
21	11.675	10.97	93.26	95.43	0.75	0.5	0.55	0.78	0.15	0.26	1
22	10.15	11.62	90.9	92.73	0.22	0.73	0.29	0.49	0.009	0.11	1
23	10.95	11.95	90.46	92.73	0.5	0.84	0.24	0.49	0.01	0.12	1
24	11.2	10.7	93.6	95.93	0.58	0.41	0.58	0.84	0.1	0.2	1
25	10.375	10.7	95.1	95.93	0.3	0.42	0.75	0.84	0.14	0.24	1
26	9.6	11.95	95.73	92.73	0.03	0.84	0.81	0.49	0.01	0.1	1
27	9.7	11.95	89.5	92.73	0.5	0.84	0.14	0.49	0.003	0.1	1
28	9.52	11.5	91.63	92.5	0	0.68	0.37	0.46	0	0.09	1
29	9.87	11.95	92.9	92.73	0.1	0.84	0.51	0.49	0.01	0.1	1
30	11.57	10.82	94.36	97.4	0.71	0.45	0.66	1	0.24	0.44	1
31	12.17	11.92	95.73	96.26	0.83	0.83	0.81	0.87	0.47	0.57	1
32	10.87	11.92	89.6	92.73	0.47	0.84	0.15	0.49	0.004	0.1	1

**Table 7.** Final predicted results in the case of simultaneous optimization of two different sets of responses (the bigger, the better) using Derringer's desirability function and Taguchi method.

Run	Factors	Level Description	Level	Contribution
1	agar	14	1	0.045
2	bean	305	2	0.032
3	ascorbic acide	2.2	2	0.004
4	ascorbate	5	2	0.048
5	yeast	50	2	0.007
6	formaldehyde	1	2	0.05
7	distilled water	1000	2	0.027
8	oil	5	1	0.018
9	chloramphenicol	2	1	0.042
10	wheat germ powder	45	2	0.04
11	Nipagin	2.2	1	0.009
12	benomyl	3	1	0.029
13	penicillin	2	1	0.106
14	temperature	25	2	0.026
15	humidity	70	2	0.019
16	container size	6	1	0.062

**Table 8.** Final Analysis of variance (ANOVA) for optimization of modified artificial diet.

No	Factors	DOF (f)	Sum of Sqr. (S)	Variance (V)	F - Ratio (F)	P	Remark
1	agar	1	0.152	0.037	4.645	0.0017	Significant
2	bean	1	0.66	0.034	4.028	0.042	Significant
3	ascorbic acide	1	0.001	0.061	0.044	0.7998	---
4	ascorbate	1	0.152	0.009	4.657	0.019	Significant
5	yeast	1	0.003	0	0.12	0.0913	---
6	formaldehyde	1	0.164	0.093	4.997	0.0026	Significant
7	distilled water	1	0.048	0.028	1.485	0.7553	---
8	oil	1	0.022	0.026	0.697	0.9368	---
9	chloramphenicol	1	0.113	0.02	3.455	0.061	---
10	wheat germ powder	1	0.102	0.018	3.134	0.1501	---
11	Nipagin	1	0.005	0.003	0.17	0.4413	---
12	benomyl	1	0.054	0.064	1.647	0.0719	---
13	penicillin	1	0.72	0.193	21.939	<0.0001	Significant
14	temperature	1	0.045	0.012	1.379	0.0864	---
15	humidity	1	0.023	0.01	0.709	0.242	---
16	container size	1	0.25	0.067	7.615	0.0042	Significant

**Table 9.** Six main factors with their high, medium and low levels chosen for further investigation of factors significance.

Factors	Levels		
	1	2	3
A: bean (g)	205	255	305
B: formaldehyde 37% (cc)	0.5	0.75	1
C: agar (g)	14	17	21
D: ascorbate (g)	3.5	4.25	5
E: penicillin (g)	2	3	4
F: container size (cm)	3 × 1	6 × 1	9 × 1

**Table 10.** ANOVA table for six main factors with their high, medium and low levels.

Factors	DOF (f)	Sum of Sqr. (S)	Variance (V)	F - Ratio (F)	P	Remark
formaldehyde	2	579.583	289.791	0.185	0.0034	Significant
ascorbate	2	50.929	25.464	0.016	0.0256	Significant
agar	2	59.271	29.635	0.018	0.0114	Significant
container size	2	1097.04	548.518	0.351	<0.001	Significant
penicillin	2	3343.8	1671.9	1.071	<0.001	Significant
bean	2	1178.93	589.466	0.377	<0.001	Significant

**Table 11.** Paired t-test for six main factors with two levels chosen for further investigation of factors significance.

Response	penicilin		bean		container size		formaldehyde		agar		ascorbate	
	Level 1	Level 2	Level 1	Level 2	Level 1	Level 2	Level 1	Level 2	Level 1	Level 2	Level 1	Level 2
Larval period	15.5	22.1	21.2	16.2	16.4	22.1	22.1	16.8	17.3	23.1	17.3	21.4
Larval mortality	3.9	5.1	3.4	5.1	3.2	5.5	5.8	3.4	3.2	5.5	3.4	5.2
Pupal Pupal	10.1	14.1	10.1	12.2	10.1	14.2	13.4	10.8	10.2	13.5	11.2	13.5
Pupal mortality	7.1	8.4	7.9	8.1	6.2	8.8	8.2	6	6.9	8.6	6.2	8.5
Longevity of adult	14	15.2	14.8	14	14	15.3	13.5	15	14.2	15.3	14	15.2
N. of eggs/ female	217	200	211	215	221	200	215	220	220	200	221	200
% hatched of eggs	59	51	56	59	59	50	53	59	58	53	58	54
t calculated> t critical (Significant)	6.87 > 2.447		9.8 > 2.447		6.70 > 2.447		6.70 > 2.447		5.96 > 2.447		5.22 > 2.447	



**Table 12.** Comparison of confirmatory experiments and predicted values of different objective functions under the optimum conditions.

Optimum conditions	Desirability Value (D)	Larval period (days)	Larval mortality (%)	Pupal period	Pupal mortality (%)	Longevity of adult (days)	N. of eggs/female	Hatching (%)
Predicted	0.789	19	4.48	12.29	7.75	14.51	211.21	54.2
Experimental	0.916	15.5	4.4	11	7.7	21	260	72