

Diversity of Non-Biting Midge Larvae Assemblages in the Jacuí River Basin, Brazil

Authors: Floss, Elzira Cecília Serafini, Kotzian, Carla Bender, Spies, Márcia Regina, and Secretti, Elisangela

Source: Journal of Insect Science, 12(121) : 1-33

Published By: Entomological Society of America

URL: <https://doi.org/10.1673/031.012.12101>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.



Diversity of non-biting midge larvae assemblages in the Jacuí River basin, Brazil

Elzira Cecília Serafini Floss^{1a}, Carla Bender Kotzian^{2b*}, Márcia Regina Spies^{3c}, Elisângela Secretti^{4d}

¹Programa de Pós-Graduação em Biodiversidade Animal, Centro de Ciências Naturais e Exatas, Universidade Federal de Santa Maria. Faixa de Camobi km 9, 97105-900, Santa Maria, RS, Brazil

²Departamento de Biologia e PPG Biodiversidade Animal, Centro de Ciências Naturais e Exatas, Universidade Federal de Santa Maria. Faixa de Camobi km 9, 97105-900, Santa Maria, RS, Brazil

³Universidade Federal do Pampa (Unipampa - Campus São Gabriel) Av. Antônio Trilha 1847, São Gabriel, RS. CEP 97300-000, Brazil

⁴Curso de Graduação em Ciências Biológicas, Centro de Ciências Naturais e Exatas, Universidade Federal de Santa Maria. Faixa de Camobi km 9, 97105-900, Santa Maria, RS, Brazil

Abstract

The richness and composition of a mountain-river chironomid larvae assemblage in the Jacuí River basin, Brazil were studied, and compared with other riverine non-biting midge larvae assemblages previously studied in the country. Additionally, the influence of some regional-scale environmental characteristics on the spatial distribution of these assemblages was tested. The specimens were collected at 12 sites in the middle course of the Jacuí River basin (in the state of Rio Grande do Sul) between April 2000 and May 2002. Around 100 taxa were recorded. The dominant taxa belonged to the genera *Rheotanytarsus*, *Cricotopus*, *Polypedilum*, and *Pseudochironomus*. Twenty-two rare taxa were found, representing 22% of the total of taxa inventoried. Fourteen genera (*Aedokritus*, *Axarus*, *Endotribelos*, *Kiefferulus*, *Manoa*, *Oukuriella*, *Phaenopsectra*, *Stenochironomus*, *Xenochironomus*, *Xestochironomus*, *Cardiocladius*, *Metriocnemus*, *Paracladius*, and *Rheocricotopus*) represent new occurrences in Rio Grande do Sul. The similarity analysis of the chironomid larvae assemblages inventoried in 32 regions of Brazil indicated five groups with similarity higher than 50%. The groups, when the effects of spatial autocorrelation were removed, displayed a weak positive correlation between the assemblage composition and the aquatic system or hydraulic conditions and the hydrographic basin, and a weak negative correlation in relation to the biome. The altitude showed no correlation with the composition of the assemblage. The relatively high richness of the region surveyed in relation to other Brazilian regions corroborates some tendencies already noted in other parts of the world, such as: i) lotic systems may constitute an exception to the rule that diversity is greater in tropical regions, ii) regions of transitional relief may contain the greatest richness of Chironomidae, and iii) in rivers, the group might have its spatial distribution influenced to a greater extent by local environmental characteristics than by regional ones.

Keywords: Chironomidae, environmental variables, inventory, Neotropics, regional scale

Correspondence: ^a elzira_bio@yahoo.com.br, ^b modrizralok@hotmail.com, ^c marciaspies@gmail.com, ^d e-secretti@hotmail.com, *Corresponding author

Editor: Henry Hagedorn was editor of this paper.

Received: 28 September 2011, **Accepted:** 14 March 2012

Copyright : This is an open access paper. We use the Creative Commons Attribution 3.0 license that permits unrestricted use, provided that the paper is properly attributed.

ISSN: 1536-2442 | Vol. 12, Number 121

Cite this paper as:

Floss ECS, Kotzian CB, Spies MR, Secretti E. 2012. Diversity of non-biting midge larvae assemblages in the Jacuí River basin, Brazil. *Journal of Insect Science* 12:121. Available online: <http://www.insectscience.org/12.121>

Introduction

The family Chironomidae is one of the most diversified aquatic insect groups, occurring in all zoogeographical regions, including Antarctica (Coffman 1995). On a global scale, 4147 species with an obligatory aquatic phase are known, attributed to 339 genera (Ferrington 2008), and grouped in 22 tribes and 11 subfamilies (Epler 2001). However, the true diversity must be still higher; estimates suggest that 8000 to 20,000 species should occur (Coffman 1995).

The non-biting midges are important inhabitants of freshwater aquatic ecosystems, where their larvae reach high densities (Trivinho-Strixino and Strixino 1999; Trivinho-Strixino 2011). They constitute an important item in the trophic chain, representing the main food of many fish and birds (Porinchu and MacDonald 2003; Sánchez et al. 2006; Fagundes et al. 2007). Additionally, they are an important tool in ecological (Armitage et al. 1995) and paleoecological (Walker 1998) studies, as well as in environmental evaluation (Rosenberg 1992), agricultural entomology (Ferrarese 1993), and public-health research (Cranston 1995a).

The chironomids are important elements of the faunas of lacustrine and fluvial biotopes (Trivinho-Strixino and Strixino 1999). They

also occur in reservoirs, although in lower richness and abundance (McKie et al. 2005; Jacobsen et al. 1997), and with different species composition (Rossaro et al. 2006). Studies analyzing the environmental factors that influence the spatial distribution of their assemblages are being developed. Landscape factors such as altitude, area size, order of river segments, drainage basin, phytogeographical unit, riparian vegetation, environmental preservation of the area, temperature, and hydraulic conditions (lentic/lotic) (e.g., Rossaro 1991; Jacobsen et al. 1997; Kleine and Trivinho-Strixino 2005; McKie et al. 2005; Morrone 2006; Rossaro et al. 2006; Roque et al. 2007; Luoto 2009; Marziali et al. 2009; Sonoda et al. 2009; Roque et al. 2010) are mentioned among the variables affecting assemblage distributions. However, studies conducted at larger spatial scales, with a geographical or regional (*sensu* Sandin and Johnson 2004) approach, are rare.

The Neotropical region, third in number of species of Chironomidae, harbors 618 species comprising 154 genera (Ferrington 2008). However, the diversity of the family is far from being well known. In Brazil, only 354 species have been described, but estimates suggest that there are about 1500 in total (Trivinho-Strixino 2011). Additionally, inventories and ecological studies about the assemblages of Chironomidae have concentrated on the northern and southeastern

regions of the country (ie.g., Callisto and Esteves 1998; Sanseverino et al. 1998; Serrano et al. 1998; Henriques-Oliveira et al. 1999; Siqueira and Trivinho-Strixino 2005; Sanseverino and Nessimian 2007; Roque et al. 2010). In the southern region, which has a climate tending toward temperate (Maluf 2000), studies are incipient, and focus primarily on lentic environments such as lakes and wetlands (e.g., Wiedenbrug et al. 1997; Stenert et al. 2004; Panatta et al. 2006, 2007). The few studies on the spatial distribution of the assemblages of larvae of Neotropical Chironomidae have focused mainly on local or small regional scales (e.g., Corbi and Trivinho-Strixino 2008a; Sonoda et al. 2009).

The middle course of the Jacuí River (southern Brazil) contains a very diversified macroinvertebrate community, as evidenced during the environmental program “Monitoring of mollusk vectors of human diseases” conducted for the construction of the Dona Francisca Hydroelectric Power Station (see Neri et al. 2005 for Heteroptera; Spies et al. 2006 for Trichoptera; Siegloch et al. 2008 for Ephemeroptera; and Pires 2011 for Odonata). Chironomidae was the most abundant family found in the river. Thus, this study presents the first survey on the diversity of a non-biting midge assemblage in a temperate climate, of a montane river in southernmost Brazil. Additionally, the similarity between this assemblage and other chironomid larvae assemblages previously studied in Brazilian rivers was also analyzed, testing the influence of regional-scale environmental factors such as altitude, biome, hydrographic basin, and hydraulic conditions, on the spatial distribution patterns of chironomid larvae assemblages, by means of four hypothesis matrices.

Materials and Methods

Study area

The study was carried out in the Jacuí River basin, in the state of Rio Grande do Sul, southern Brazil. Its main headwaters are in the Planalto (central plateau) region, at a mean altitude of approximately 730 meters above sea level, and its mouth is in the Central Depression, where it contributes to formation of the Jacuí delta, in Guaíba Lake. The river is 710 km long, and its 71,600 km² drainage basin is characterized by intense land use for agriculture, livestock, energy generation, and urban supply (FEPAM 2010).

The middle course of the Jacuí River is located in the transition zone between the physiogeographic regions of the lower Northeast Slope and Central Depression, with altitudes from 50 to 500 meters above sea level (Pereira et al. 1989). The climate is humid subtropical (Cfa, according to Köppen’s classification) with warm and rainy summers, but considered temperate locally (Maluf 2000). Rainfall is regularly distributed throughout the year, with mean precipitation varying from 1500 to 1708 mm, and the monthly mean temperature ranges from 13° C during winter to 18–22° C during summer (Pereira et al. 1989; Maluf 2000). The native vegetation is mainly Seasonal Semideciduous Forest (*sensu* Prado 2000), currently vastly altered, and represented only by small portions of secondary forest, sparsely distributed along rivers and slopes (Longhi et al. 1982; Marchiori et al. 1982).

In 2000, the middle course of the Jacuí River was dammed in its final section by the construction of the Dona Francisca Hydroelectric Power Station (29° 26’ 50” S; 53° 16’ 50” W). The water reservoir inundated six municipalities, covering an area

of 1337 ha, which contributed to the environmental changes in the region.

Sampling methods

Sampling was conducted irregularly between April 2000 and May 2002. Twelve sites were selected for the study, two in the main channel of the Jacuí River, and the remaining sites in its tributaries (seven in three tributaries of the right bank, and three in two tributaries of the left bank) (Figure 1). The environmental characterization of these sites is presented in Table 1. The hydrological classification of stream orders was done on a cartographic scale of 1:50,000, according to Strahler (1957).

The collection was carried out using a Surber-type sampler (area = 0.36 m², mesh = 1 mm). At each site, three subsamples were collected, one in midriver and one near each bank, except in the Jacuí River, where only one of the banks was sampled. The subsamples were pooled in a single sample per site. The sampling took place in shallow water, never deeper than 1 m, in the river. Macrophytes present on the substrate were scraped and collected. The material obtained was fixed in 80% ethanol. The material was sorted, and specimens were counted under a stereoscopic microscope. For taxonomic identification, the specimens were cleared in 10% KOH (potassium hydroxide), mounted on semi-permanent slides using Hoyer's medium, and analyzed under an optical microscope (Trivinho-Strixino and Strixino 1995; Epler 2001; Trivinho-Strixino 2011).

Because of the large numbers of specimens in some samples (> 2000), in samples with over 100 larvae, a subsample with 100 specimens was randomly selected for identification purposes. The number of individuals per taxon was estimated afterwards, according to the

percentage recorded in the subsample. An ongoing study conducted by the authors, in which all the specimens of chironomids of each sample are identified, showed that there is no difference in the diversity of chironomids in the area studied, if all or only 100 of the specimens are classified. The specimens were identified to genus and/or species level or morphotype using the taxonomic keys of Wiederholm (1983), Trivinho-Strixino and Strixino (1995), Saether et al. (2000), and Epler (2001). Later, the identifications were confirmed by experts.

Voucher specimens were deposited in the Coleção de Zoologia, of the Departamento de Biologia, of the Universidade Federal de Santa Maria, Rio Grande do Sul state, and in the Laboratório de Hidrobiologia of the Universidade Federal de São Carlos, São Paulo state.

Data analysis

The sampling effectiveness was evaluated by a species accumulation curve (collector curve), which was obtained by the mean of 500 curves generated by the random addition of samples using EstimateS 7.5 software (Colwell 2006). This method was chosen because it calculates the fluctuations around the mean curve when the data sets are added, so it is the best method to evaluate the similarity of the inventory to the total richness of the area (Colwell and Coddington 1994).

The composition of the chironomid larvae assemblage recorded in the middle course of the Jacuí River and its tributaries was compared to the assemblages found by inventories conducted at 32 other locations in Brazil (Table 2). Among the inventories found in literature, those with sampling methods favoring benthic fauna were selected. In cases when an area was surveyed more than once, or

in cases when several surveys were made in areas close to each other (< 10 km), only the studies with the most complete and/or up-to-date lists were included. In the studies carried out in wide areas, such as along the Iguaçu River, in the state of Paraná (Takeda et al. 2005), the taxa list of the assemblage was divided into sectors (i.e., sections of the river course), if the article provided the taxa list by sampling sites. In order to compare the assemblages of different studies, taxa with uncertain identifications were discarded (i.e., Genus A, Genus 1), as well as identifications below genus level, because in many cases such an identification is based on morphotypes (e.g., Suriano and Fonseca-Gessner 2004; Siqueira and Trivinho-Strixino 2005; Rosin et al. 2009), and is not suitable for comparison between different studies.

The similarity between the chironomid larvae assemblages was calculated using the Coefficient of Geographic Resemblance (Duellman 1990). This coefficient is expressed by Coefficient of Geographic Resemblance = $2N_s / (N_a + N_b)$, where N_s = number of species in both areas, N_a = number of species in area A. and N_b = number of species in area B. It is equivalent to Sørensen's, Dice's, and Czekanowski's indices (Wolda 1981; Krebs 1999; Magurran 2004), and was performed using NTSYS PC 2.10s software. The matrix of similarity (Coefficient of Geographic resemblance) was later represented by means of clustering analysis with the weighted pair-group method with an arithmetic average (Sokal and Michener 1958) to avoid the effect of sample size (richness of genera in different assemblages) on the analyses (Valentin 1995). Possible distortions in the graphical representation of the matrix of similarity by clustering analysis were evaluated by means of the cophenetic correlation coefficient (r)

(Romesburg 1984). The closer the value of r is to 1, the smaller is the distortion (*sensu* Rohlf 2000).

The similarity among taxa lists of assemblages of inventories can be affected by differences in sampling effort, such as the size of the area surveyed, periodicity of sampling, etc. However, this tendency can be minimized if studies carried out with a low sampling effort on the temporal axis show higher effort on the spatial axis (and vice-versa) (Santos et al. 2009). The differences between these studies are considered here, and reported in Table 2.

Data sets based on a spatially structured schedule may show relationships due to geographical proximity, which is known as spatial autocorrelation (Legendre and Legendre 1998). In order to test this relationship, a matrix of geographical distances between locations was constructed, obtained from the geographical coordinates cited in the publications or found by using Google Earth (<http://www.google.com/earth>). The correlation between the matrix of geographical distance and the matrix of similarity in the composition of the chironomid larvae assemblages was determined by the Mantel Test (Manly 2000). This test indicates the correlation between matrices based on Z-statistics, where Z depends on the number or magnitude of elements in the matrix to be compared (Valentin 1995). Thus, normalization is needed to transform Z into a coefficient (r) that varies from 1 to -1 (Valentin 2000). The significance of Z was determined by the Monte Carlo permutation test (Smouse et al. 1986), using 5000 permutations. The Mantel and Monte Carlo tests were performed using the NTSYS PC 2.10s software (Rohlf 2000).

The observed pattern of the spatial distribution of chironomid larvae assemblages was correlated with four hypothesis matrices, based on regional environmental characteristics designed to explain the pattern that was found:

1) Altitude hypothesis matrix: this postulates that the assemblages of Chironomidae at the locations with similar altitudes are more similar to each other. This hypothesis is based on the altitudes of the different locations of the studies compared (as provided in the reports and/or obtained using the geographical coordinates in the online version of Google Earth). The matrix of similarity of altitude among the locations was obtained through the Euclidean Distance index (Magurran 2004).

2) Biome hypothesis matrix: this postulates that the compositions of the chironomid assemblage at locations in the same biome are more similar to each other than to those of other biomes. This hypothesis matrix is based on the locations of the inventories, and on the Brazilian biome classification (i.e., Cerrado, Amazon Forest, Atlantic Forest, Pantanal, and Pampa, according to the Instituto Brasileiro de Geografia e Estatística (2003)). A binary matrix was constructed in which pairs of locations from different biomes (i.e., Cerrado and Atlantic Forest) received a similarity value of 0, and pairs of locations from the same biome received a similarity value of 1.

3) Hydraulic conditions (lotic or lentic) hypothesis matrix: this postulates that the chironomid assemblages at locations with the same hydraulic conditions are more similar to each other than to those with different conditions. Reservoirs and other impoundments were considered lentic environments. In order to obtain the matrix of similarity, pairs of locations with different

hydraulic conditions received the value 0 (i.e., rivers x lakes), pairs with the same conditions received the value 1, and pairs of locations in which one inventory involved lotic and lentic environments, and another only one of these conditions, received the value 0.5.

4) Hydrographic basin hypothesis matrix: this postulates that the assemblages from locations within the same hydrographic basin are more similar to each other than to those in different hydrographic basins. The hydrographic basin of each study was obtained by plotting (geographical coordinates) the locations on the map of the major hydrographic basins of Brazil. These basins, shown in Figure 2, are: Rio Amazonas (Amazon River), Rio da Prata (Plate River), Costeira Sudeste (Southeast Coastal), and Costeira Sul (South Coastal), according to the Instituto Brasileiro de Geografia e Estatística (2003). In order to obtain the matrix of similarity, pairs of locations in different hydrographic basins received the value 0, and pairs of locations in the same hydrographic basin received the value 1.

Because of the occurrence of spatial autocorrelation ($r = 0.257$; $p = 0.016$), a partial Mantel test (Smouse et al. 1986) was run to test the correlation between the matrix of similarity for the composition of the chironomid larvae assemblages and each of the four hypothesis matrices, using the matrix of geographical distance to avoid the effect of geographical proximity. This test consists of the comparison of two matrices (A and B), removing the effects of a third matrix (C, in the present study, corresponds to the geographical distance matrix) on the first two, using a regression of C on A and B. Thus, the residual matrix obtained represents the variations of matrices A and B that cannot be explained by matrix C (Smouse et al. 1986).

Then, the two residual matrices can be compared freely. For similarity and grouping analysis, Mantel and partial Mantel tests were performed using NTSYSpc 2.10S software (Rohlf 2000).

Results

Taxonomic composition and richness in the middle course of the Jacuí River

In total, 12,346 larvae of Chironomidae were collected, classified in 99 taxa (84 genera and/or species and 15 morphospecies), attributed to three subfamilies (Table 3). The subfamily that showed the greatest richness was Chironominae (68 taxa). Other subfamilies showed lower richness (Tanypodinae, 11 taxa; Orthocladiinae, 20 taxa). The dominant taxa were *Rheotanytarsus* sp. 1 (18.3%), *Cricotopus* sp. 2 (16.3%), *Cricotopus* sp. 1 (11.2%), *Polypedilum* (*Polypedilum*) sp. 1 (7.2%), *Rheotanytarsus* sp. 2 (7%), *Pseudochironomus* (6.6%), and *Polypedilum* (*Polypedilum*) sp. 2 (6.4%). Of these, only *Rheotanytarsus* sp. 1 occurred at all sites. Twenty-two taxa were rare (up to three larvae), which corresponds to 22% of the total of taxa, and each taxon occurred at only one site (Table 3).

The species accumulation curve for the 12 sampling sites (88 samples in total) in the middle course of the Jacuí River basin was stable, with little variation along the mean curve, showing that the asymptote was reached (Figure 3).

Regional distribution pattern

The cluster analysis of the locations with inventories of chironomid larvae assemblages in Brazil showed the formation of five groups with similarity higher than 50%. Group (i) clustered mainly assemblages of locations in the Rio da Prata basin and the Cerrado and

Atlantic Forest biomes, although some locations within the Costeira Sul basin and the Pantanal biome were also included (BPCR1, BPCR6, BPCR5, BPMT1, BPCR2, BPCR7, BPMT14, BPMT13, BCSMT, and BPPAN2). Group (ii) clustered mainly assemblages of locations in the Rio da Prata basin and the Atlantic Forest biome, but also included some locations in the Costeira do Sudeste basin and the Cerrado biome (BCSeMT3, BPMT5, BPMT6, BPCR3, BPMT7, BPMT8, BCSeMT7, BPMT12, and BPMT11). Group (iii) clustered assemblages of locations exclusively in the Atlantic Forest and, except for one location, the Costeira do Sudeste basin (BCSeMT1, BCSeMT5, BCSeMT4, BCSeMT6, BCSeMT2, and BPMT3). Group (iv) clustered locations in two different basins and biomes (BAAM and BCSPM). Group (v) clustered two locations in the Rio da Prata basin and the Atlantic Forest biome (BPMT2 and BPMT10).

The main taxa responsible for clustering the five groups of chironomid larvae assemblages are presented in Table 4. In group i, 21 taxa were frequent, i.e., occurred in 70% or more of the locations compared, such as *Beardius*, *Caladomyia*, *Endotribelos*, *Goeldichironomus*, *Stenochironomus*, *Clinotanytus*, *Coelotanytus*, and *Procladius*; furthermore, 11 taxa did not occur in group ii. Group ii contained nine frequent taxa, which were also frequent in locations of group i, as well as taxa that were frequent only within group ii, such as *Democritus*, *Cladopelma*, *Dicrotendipes*, and *Fissimentum*. In group iii, 12 taxa were frequent, of which five were not frequent in groups i and ii (*Nimbocera*, *Oukurriella*, *Pelomus*, *Stempellinella*, and *Parametriocnemus*), and five were exclusive (*Nilotanytus*, *Thienemanniella*, *Mesosmitia*, *Pseudosmittia*, and *Rheocricotopus*). Group iv shared the genera *Chironomus*, *Ablabesmyia*,

and *Larsia*, and contained one exclusive taxon (*Macropelopia*). In group 'v', the genera *Axarus*, *Chironomus*, *Cryptochironomus*, *Glyptotendipes*, *Nimbocera*, *Tanytarsus*, and *Djalmabatista* were shared, and *Glyptotendipes* and *Micropsectra* were exclusive.

The partial Mantel tests (i.e., without spatial autocorrelation effect) indicated weak positive correlations between the distribution of the chironomid larvae assemblages and the hydraulic condition matrix (lotic or lentic), and the matrix of their hydrographic basins ($r = 0.153$, $p = 0.05$; $r = 0.149$, $p = 0.05$ respectively). The correlation between the assemblage distribution and the biome matrix was weakly negative ($r = -0.136$, $p = 0.05$). On the other hand, altitude did not show a significant correlation with the distribution of the chironomid larvae assemblages ($r = 0.005$, $p = 0.454$).

Discussion

Taxonomic composition and richness in the middle course of the Jacuí River

The stability and asymptote reached by the collection curve of the present study suggest that little or no increase would be expected with greater sampling effort. Hence, considering that the occurrence of over 100 species of Chironomidae, most of them rare, in a single river is common (Coffman 1995; Cranston 1995a; Roque et al. 2007), and that a high level of endemism is expected for different biogeographical regions (Coffman 1989), it is possible that many of the 99 taxa found in the middle course of the Jacuí River, of which 20% were rare, are new species. However, as the identification of larvae of Chironomidae at species level is related to the prior description of adults and their associated larvae (Oliver 1971; Pinder 1986; Raunio

2008), it is not possible to tell how many of these taxa might represent new species.

Fourteen genera found in the study area are new occurrences for Rio Grande do Sul (*Aedokritus*, *Axarus*, *Endotribelos*, *Kiefferulus*, *Manoa*, *Oukuriella*, *Phaenopsectra*, *Stenochironomus*, *Xenochironomus*, *Xestochironomus*, *Cardiocladius*, *Metriocnemus*, *Paracladius*, and *Rheocricotopus*). Adding the 99 taxa recorded here to the 13 other genera recorded by Wiedenbrug et al. (1997) and Panatta et al. (2007) for the lakes and wetlands of the Coastal Plain, and by Hepp et al. (2008) and Wiedenbrug et al. (2009) for small streams (*Apedilo*, *Clinotanypus*, *Coelotanypus*, *Onconeura*, *Alotanypus*, prox. *Macropelopia*, prox. *Adenopelopia*, *Procladius*, *Monopelopia*, *Psectrocladius*, *Fitkauimya*, *Paralauterborniella*, and *Stempellina*), the total richness of Chironomidae in Rio Grande do Sul reaches at least 112 taxa.

In terms of geographical patterns, the richness of Chironomidae tends to be greater and to show higher endemism rates in tropical and subtropical climates (Fittkau 1971; Coffman 1989; Cranston 1995b). However, inventories conducted in Brazilian rivers have recorded lower richness levels of genera or species (between 11 and 71) in both tropical (Callisto and Esteves 1998; Marques et al. 1999; Sanseverino and Nessimian 2001; Amorim et al. 2004; Suriano and Fonseca-Gessner 2004; Siqueira and Trivinho-Strixino 2005; Aburaya and Callil 2007; Corbi and Trivinho-Strixino 2008a, b; Silva et al. 2008; Siqueira et al. 2009; Simião-Ferreira et al. 2009; Siqueira et al. 2009) and subtropical (Takeda et al. 2005; Resende and Takeda 2007; Rosin et al. 2009; Sonoda et al. 2009; Rosin et al. 2009) climates than in the cooler temperate climate of this study. Some workers have recorded high

richness (200 species) of Chironomidae in temperate lotic environments (e.g., Raunio 2008, in rivers of Finland). Therefore, it is possible that lotic ecosystems constitute an exception to the tendency for richness to be highest in tropical areas, although the available data are somewhat contradictory (McKie et al. 2005). However, other environmental factors can favor high richness of Chironomidae in rivers.

Environmental gradients have been related to the diversity of non-biting midges (Pinder 1995; McKie et al. 2005). The distribution patterns of chironomid subfamilies vary according to the relief. The Orthocladiinae are common in streams located on plateaus, because they are more adapted to cool and well oxygenated waters, while the Chironominae are well adapted to live in lowlands, dwelling in fine sediments, and are tolerant of high temperatures and variations in oxygen content (Pinder 1995). Thus, in regions of transition from rithral to potamic areas, some species live near their limits of ecological tolerance (Statzner and Higler 1986), and so these species can overlap in their distributions (Principe et al. 2008), allowing more species to coexist. The area studied here is a slope, between the upper course (in the uplands of the Planalto) and the lower course (in the lowlands of the Central Depression) of the Jacuí River. Hence, the high richness observed in the middle course of the Jacuí River could be related to its transitional relief.

Environmental heterogeneity is another factor that may promote high richness of macroinvertebrates (e.g., Beisel et al. 2000; Voelz and McArthur 2000; Lencioni and Rossaro 2005). Structurally complex substrates (wood, leaves, stones and gravels, macrophytes, and moss) can provide more

niches, with refuges and food resources, as well as protection from predation (Principe and Corigliano 2006), than structurally simple ones (sand or mud). This condition can also facilitate the colonization of middle and lower courses of rivers by taxa characteristic of rithral areas (Tokeshi and Pinder 1985). Many sites of the middle course of the Jacuí River have gravels with encrusting aquatic macrophytes (i.e., *Podostemon*), generating a layer of macrophytes and fine sediments over a gravel substrate. These macrophyte patches increase the heterogeneity of the environment, and probably contributed to the high richness recorded.

The stream order can also influence the composition and richness of chironomid assemblages, especially if the order covaries with altitude, substrate granulometry, and land use (e.g., Lindegaard 1995; Principe et al. 2008; Puntí et al. 2009). In this study, sampling was conducted in stretches from the first to seventh orders. However, altitude (70 to 140 meters above sea level) and granulometry (boulders and cobbles) did not vary among sites, showing no longitudinal gradient. In the middle course of the Jacuí River, the highest richness levels were recorded at four sites (1, 3, 7, 11) ranging from third to seventh order. This result was strongly influenced by the larger sampling effort (higher abundance of larvae) used at those segments (12 samples in each one). In other words, the sampling design does not allow discussion of the role of stream order in affecting the richness of Chironomidae assemblages. The richness levels previously recorded in other Brazilian rivers do not follow a pattern. The highest numbers of species were found in large-order stretches (Takeda et al. 2005; Rosin et al. 2009), but large-order stretches can also show lower richness (Callisto and Esteves 1998; Aburaya

and Callil 2007). Thus, climate, relief (slope), and heterogeneity of the substrate possibly play more important roles in affecting the diversity of chironomids in the region studied.

Because sampling in the middle course of the Jacuí River was originally planned to collect mollusks, the abundance and richness of many macroinvertebrates could be underestimated. The mesh size of 1 mm may have allowed many minute specimens to be lost. Thus, the diversity of chironomids might increase if a smaller mesh size were used. However, the sampling area of 0.36 m², and the tangles of *Podostemum* scraped from the stones and added to the samples, in which many chironomid larvae were attached, may have counteracted the large mesh-size effect. In any event, Chironomidae was the most abundant family in the area, and reached the highest richness among the insects found in the river (Neri et al. 2005; Spies et al. 2006; Siegloch et al. 2008; Pires 2011).

The predominant taxa in the middle course of the Jacuí River belong to genera that are characteristic of potamic or backwater environments, with stony substrate and litter deposition. *Polypedilum*, *Rheotanytarsus*, and *Cricotopus* have been recorded in several other Brazilian rivers with stony bottoms (Sanseverino and Nessimian 2001; Roque et al. 2003; Suriano and Fonseca-Gassner 2004) and litter deposition (Sanseverino and Nessimian 2001). *Polypedilum* sp. and *Rheotanytarsus* sp. are characteristic of rapids, with a coarse substrate and turbulent flow (Principe et al. 2008). Even though species of *Polypedilum* are usually associated with fine sediments, some of its species can be found in coarse substrates (Pinder and Reiss 1983).

Regional distribution pattern

Studies conducted with freshwater macroinvertebrates and other animal groups showed that environmental factors of large spatial scale, such as climate (Bonada et al. 2008), altitude (Maltchik et al. 2010), phytogeographical unit (Santos et al. 2009), and hydrographic basin (Martel et al. 2007), can affect the spatial distribution of the communities. However, the hydraulic conditions of the environments (i.e., lentic or lotic) seem to be the most important factor influencing the macroinvertebrates at the regional scale (Buffagni et al. 2009, 2010). Studies on large-scale patterns of diversity and distribution in lotic environments are few, and are restricted to temperate regions of the Northern Hemisphere (Vinson and Hawkins 1998).

Studies of the regional-scale spatial distribution of chironomid assemblages are also rare. The taxonomic composition of families is more similar in tropical and subtropical areas (Cranston and Naumann 1991; Pinder 1995; Rossaro et al. 2006). The influence of other factors, such as temperature, altitude, phytogeographic unit, hydrography, etc., is little investigated, and the few data available are incipient and/or contradictory. Changes in chironomid assemblages were observed along altitude gradients in European lakes (Bitusík and Svitok 2006), but not in rivers of the bioregion of the Humid Tropics in Australia (McKie et al. 2005). Changes were also observed according to the mean July air temperature in Finland lakes (Luoto 2009). However, chironomid genera have been used to identify the hydraulic conditions of ancient aquatic environments in paleoecological studies (Porinchu and MacDonald 2003), because lotic settings usually support a greater

diversity of larvae (Lindergaard 1995; Pinder 1995).

In the present study, the absence of influence of altitude on the spatial pattern of the assemblages, as well as the small degree of influence shown by hydrographic basins, biomes, and hydraulic conditions, probably indicate the relatively small influence of regional environmental factors on the distribution patterns of lotic chironomids. McKie et al. (2005) suggested that the Chironomidae have such a wide tolerance to many environmental variables that they are not affected by regional-scale factors. Additionally, studies conducted with communities of macroinvertebrates and focusing on multiple scales have demonstrated that a large part of the variation in community structure may be influenced by local factors (Rios and Bailey 2006). Hence, the spatial pattern of distribution of chironomid larvae observed in this study may have been more influenced by local factors. Previous studies have shown that local factors, such as oxygen, water velocity and temperature, pH, solid material in suspension, phosphorus, sulfate, presence of algae and macrophytes, type of sediment or substrate, calcium and ferrous ions, and electrical conductivity commonly affect the richness, abundance, and/or composition of communities (Ali et al. 2002; Bisthoven et al. 2005; Inoue et al. 2005; Woodcock et al. 2005; Principe et al. 2008; Siqueira et al. 2008; Al-Shami et al. 2010; Luoto 2011).

Among the taxa that were frequent only in group i, the genera *Beardius*, *Caladomyia*, *Endotribelus*, *Goeldichironomus*, *Stenochironomus*, *Clinotanypus*, *Coelotanypus*, and *Procladius*, and some exclusive genera (*Goeldichironomus*, *Polypedilum* (*Asheum*)) are characteristic of

lentic environments associated with litter, higher temperatures, changes in the hydrological regime, and the presence of macrophytes (Epler 2001). Many of the locations where this group was found show these characteristics, such as the Rio Paraná, Rio Ivinhema, and Saracacá and Carnã creeks. The most frequent and exclusive taxa of group ii (*Aedokrytus*, *Cladopelma*, *Dicrotendipes*, *Fissimentum*, *Tanytarsus*, and *Ablabesmyia*) are typical of lentic environments and sandy substrates, and are resistant to certain types of environmental degradation, such as the absence of riparian forest (Pinder and Reiss 1983; Epler 2001). Human activities including agriculture, dams, deforestation, erosion and silting in streambeds, and discharge of industrial and domestic wastes are prevalent in group ii locations.

Among the taxa that were frequent only in group iii or exclusive to it, *Mesosmittia*, *Pseudosmittia*, *Rheocricotopus*, and *Thienemanniella* are characteristic of montane rivers and streams with good environmental preservation (Cranston et al. 1997), conditions shown by the locations of this group (Sanseverino and Nessimian 1998; Roque et al. 2007). The most frequent or exclusive taxa of group iv, such as *Nilothauma*, *Alotanypus*, *Labrundinia*, *Larsia*, and *Macropelopia*, are resistant to environmental degradation (Spies and Reis 1996; Epler 2001). These conditions are found in Porto Trombeta creek and the Bela Vista and Ouro streams (Callisto and Esteves 1998; Panatta et al. 2006).

Among the taxa exclusive to and frequent only in group v, *Axarus*, *Glyptotendipes*, *Micropsectra*, and *Paracladopelma* are characteristic of littoral or sublittoral environments, in shallow lentic and lotic environments with slow flow, high temperatures, organic-matter concentration,

and fine sediment (mud), in mesotrophic and oligotrophic environments (Pinder and Reiss 1983). The two locations where the genera of group v were found had these characteristics (Takeda et al. 1997; Santos and Henry 2001).

The chironomid larvae assemblage of the middle course of the Jacuí River was most similar to the assemblages found in group i locations, and shared certain frequent genera, such as *Rheotanytarsus*, *Cricotopus*, *Polypedilum*, *Rheotanytarsus*, and *Pseudochironomus*. However, the most abundant taxa found in the area studied here, as previously discussed, are characteristic of lotic environments with rocky substrates and litter deposition, and not of lentic environments, which characterize the environmental preferences of the majority of the frequent species of group i.

Furthermore, the Jacuí River harbored several taxa that were exclusive to this group, such as *Kiefferulus*, *Xestochironomus*, *Metriocnemus*, *Monopelopia*, *Onconeura*, and *Paracladius*. It is necessary to consider that taxa typical of both uplands and lowlands coexist in the transition regions of the middle course of the Jacuí River, and that the similarity analysis considered only the occurrence of genera.

Local factors such as granulometry, presence of macrophytes, and leaf litter might have influenced the groupings that were formed. However, factors associated with the terrestrial environment, which are usually not taken into account in studies on aquatic communities, might also have influenced the spatial distribution of the Chironomidae. The occurrence of adults of this family is influenced primarily by typically terrestrial factors such as humidity, insolation, shade, air temperature, and predation (Armitage et al. 1995; Tokeshi 1995). Furthermore, it is

possible that species, even more than genera, might respond better to the factors tested in our study. For instance, larvae of different species of a single genus (such as *Orthocladius*, *Rheotanytarsus*, *Thienemanniella*, and *Polypedilum*) have different environmental preferences in relation to granulometry and water velocity (Pinder and Reiss 1983; Pinder 1995; Epler 2001).

Final considerations

The richness of larvae of Chironomidae recorded in the middle course of the Jacuí River was higher than those recorded in the warmer tropical and subtropical regions of Brazil. This high richness constitutes even more evidence that lotic ecosystems provide a general exception to the tendency of richness of species to be greater in tropical regions (McKie et al. 2005). This study also corroborates two patterns observed in previous studies of the spatial distribution of lotic Chironomidae: 1) the existence of greater diversity in mountain regions, and in transition zones of rithral and potamic areas, and 2) the possibility that environmental regional-scale factors related to altitude and temperature exert little influence on the distribution of Chironomidae. Moreover, it is important to emphasize the need for additional studies, conducted on local and large spatial scales, especially in Brazil, in order to fill the existing lacunae in knowledge, and to understand the spatial distribution patterns recorded.

Acknowledgments

The authors thank Dr. Susana Trivinho-Strixino (UFSCAR) for giving the opportunity of an internship to E. Floss, and for helping with or confirming the identification of taxa; Dr. Ana Emília Siegloch (UFSC) and Dr. Rosemary Davanzo for their helpful

suggestions that significantly improved this manuscript; Biologist Mateus Marques Pires for helping with the execution of the statistical tests and graphing programs, and information exchange; Dr. Marlise L. Bartholomei-Santos (UFES) for lending the microscope and camera that were used to identify the taxa; DFESA (Dona Francisca Energética) for financial support; and CAPES for awarding a Master's scholarship to E. Floss. Finally, we thank the anonymous referees for constructive suggestions, and Dr. Janet W. Reid for revising the English text.

References

- Aburaya FH, Callil CT. 2007. Variação temporal de larvas de Chironomidae (Diptera) no Alto rio Paraguai (Cáceres, Mato Grosso, Brasil). *Revista Brasileira de Zoologia* 24(3): 565-572.
- Al-Shami SHA, Rawi CSM, Ahmad AH, Nor SAM. 2010. Distribution of Chironomidae (Insecta: Diptera) in polluted rivers of the Juru River Basin, Penang, Malaysia. *Journal of Environmental Sciences* 22(11): 1718-1727.
- Ali A, Frouz J, Lobinske RJ. 2002. Spatio-temporal effects of selected physico-chemical variables of water, algae and sediment chemistry on the larval community of nuisance Chironomidae (Diptera) in a natural and a man-made lake in central Florida. *Hydrobiologia* 470: 181-193.
- Amorim M, Henriques-Oliveira AL, Nessimian JL. 2004. Distribuição espacial e temporal das larvas de Chironomidae (Insecta: Diptera) na seção ritral do Rio Cascatinha, Nova Friburgo, Rio de Janeiro, Brasil. *Lundiana* 5(2): 119-127.
- Armitage PD, Cranston PS, Pinder LCV. 1995. *The Chironomidae. The Biology and Ecology of Non-Biting Midges*. Chapman and Hall.
- Beisel JN, Usseglio-Polatera P, Moreteau JC. 2000. The spatial heterogeneity of a river bottom: a key factor determining macroinvertebrate communities. *Hydrobiologia* 422(423): 163-171.
- Bisthoven LJ, Gerhardt A, Soares AMVM. 2005. Chironomidae larvae as bioindicators of an acid mine drainage in Portugal. *Hydrobiologia* 532: 181-191.
- Bitusík P, Svitok M. 2006. Structure of Chironomid assemblage along environmental and geographical gradients in the Bohemian Forest lakes (Central Europe): An exploratory analysis. *Biologia* 61(20): 467-476.
- Bonada N, Rieradevall M, Dallas H, Davis J, Day J, Figueroa R, Resh VH, Prat N. 2008. Multi-scale assessment of macroinvertebrate richness and composition in Mediterranean-climate rivers. *Freshwater Biology* 53: 772-788.
- Buffagni A, Armanini DG, Erba S. 2009. Does the lentic-lotic character of rivers affect invertebrate metrics used in the assessment of ecological quality? *Journal of Limnology* 68(1): 92-105.
- Buffagni A, Erba S, Armanini DG. 2010. The lentic-lotic character of Mediterranean rivers and its importance to aquatic invertebrate communities. *Aquatic Sciences* 72: 45-60.
- Callisto M, Esteves FA. 1998. Biomonitoramento da macrofauna bentônica de Chironomidae (Diptera) em dois igarapés amazônicos sob influência das atividades de

uma mineração de bauxita. In: Nessimian JL, Carvalho AL, Editors. *Ecologia de Insetos Aquáticos. Série Oecologia Brasiliensis* 5: 299-309.

Coffman WP. 1989. Factors that determine the species richness of lotic communities of Chironomidae. *Acta Biologica Debrecina* 3: 95-100.

Coffman WP. 1995. Conclusions. In: Armitage EPD, Cranston PS, Pinder LCV, Editors. *The Chironomidae. The Biology and Ecology of Non-Biting Midges*. pp. 438-447. Chapman and Hall.

Colwell RK. 2006. Estimates: Statistical estimation of species richness and shared species from samples. Version 7.5, Available online: <http://viceroy.eeb.uconn.edu/estimates>.

Colwell RK, Coddington JA. 1994. Estimating terrestrial biodiversity through extrapolation. *Philosophical Transaction of the Royal Society of London* 345(1311): 101-118.

Corbi JJ, Trivinho-Strixino S. 2008a. Effects of land use on lotic Chironomid communities of Southeast Brazil: Emphasis on the impact of sugar cane cultivation. *Boletim do Museu Municipal do Funchal* 13: 93-100.

Corbi JJ, Trivinho-Strixino S. 2008b. Relationship between sugar cane cultivation and stream macroinvertebrate communities. *Brazilian Archives of Biology and Technology* 51(4): 769-779.

Cranston PS. 1995a. Introduction. In: Cranston PS, Organizers. Chironomids: from genes to ecosystems. *Proceedings of XII International Symposium on Chironomidae*. pp. 1-2.

Cranston PS. 1995b. Introduction. In: Armitage PPS, Cranston PS, Pinder LCV, Editors. *The Chironomidae. The Biology and Ecology of Non-biting Midges*. pp. 1-7. Chapman and Hall.

Cranston PS, Cooper PD, Hardwick RA, Humphrey CL, Dostine PL. 1997. Tropical acid streams: The Chironomid (Diptera) response in northern Australia. *Freshwater Biology* 37: 473-483.

Cranston PS, Naumann I. 1991. Biogeography. In: Cranston PS, Naumann I, Editors. *Insects of Australia*. pp. 181-197. CSIRO: Melbourne University Press.

Dorfeld CB, Fonseca-Gessner AA. 2005. Fauna de Chironomidae (Diptera) associada à *Salvinia* sp. e *Myriophyllum* sp. em um reservatório do Córrego do Espirado, São Carlos, São Paulo, Brasil. *Entomologia y Vectores* 12(2): 181-182.

Duellman WE. 1990. Herpetofauna in Neotropical rainforest: comparative composition history and resource use. In: Gentry AH, Editor. *Four Neotropical Rainforests*, 4th edition. Yale University Press.

Epler JH. 2001. *Identification Manual for the Larval Chironomidae (Diptera) of North and South Carolina. A guide to the taxonomy of the midges of the southeastern United States, including Florida*. Special Publication SJ2001-SP13. North Carolina Department of Environment and Natural Resources, Raleigh, NC, and St. Johns River Water Management District.

Fagundes CK, Behr ER, Kotzian CB. 2007. Alimentação de *Rhinodoras dorbignyi*

(Kröyer, 1855) (Siluriformes: Doradidae) no rio Ibicuí, Rio Grande do Sul, Brasil. *Acta Scientiarum Biological Sciences* 29(2): 137-143.

FEPAM (Fundação Estadual de Proteção Ambiental) 2010. *Qualidade das Águas da Bacia Hidrográfica do Rio Jacuí*, Available online: http://www.fepam.rs.gov.br/qualidade/qualidade_jacui/jacui.asp.

Ferrarese U. 1993. Chironomids of Italian rice fields. *Netherlands Journal of Aquatic Ecology* 26: 341-346.

Ferrington JL. 2008. Global diversity of non-biting midges (Chironomidae; Insecta-Diptera) in freshwater. *Hydrobiologia* 595: 447- 455.

Fittkau EJ. 1971. Distribution and ecology of Amazonian Chironomids (Diptera). *The Canadian Entomologist* 103: 407-413.

Freire CF, Fonseca-Gessner AA. 2001. Larvas de Chironomidae (Diptera) na Microbacia do Ribeirão Cachim, São Carlos, São Paulo, Brasil. *Entomología y Vectores* 8(4): 417-429.

Henriques-Oliveira AL, Nessimian JL, Dorvillé LFM. 2003. Distribution of Chironomidae larvae fauna (Insecta: Diptera) on different substrates in a stream at Floresta da Tijuca, RJ, Brazil. *Acta Limnologica Brasiliensia* 15(2): 69-84.

Henriques-Oliveira AL, Sanseverino AM, Nessimian JL. 1999. Larvas de Chironomidae (Insecta: Diptera) de substrato rochoso em dois rios em diferentes estados de preservação na Mata Atlântica, RJ. *Acta Limnologica Brasiliensia* 11(2): 17-28.

Hepp LU, Biasi C, Milesi SV, Veiga FO, Rastello RM. 2008. Chironomidae (Diptera) larvae associated to *Eucalyptus globulus* and *Eugenia uniflora* leaf litter in a subtropical stream (Rio Grande do Sul, Brazil). *Acta Limnologica Brasiliensia* 20(4): 345-350.

Higuti J, Takeda AM. 2002. Spatial and temporal variation in densities of Chironomid larvae (Diptera) in two lagoons and two tributaries of the upper Paraná River floodplain, Brazil. *Brazilian Journal of Biology* 62(4B): 807-818.

Instituto Brasileiro de Geografia e Estatística. 2003. *Divisão Hidrográfica Nacional*. Available online: http://www.ibge.gov.br/ibgeteen/atlasescolar/mapas_pdf/brasil_bacias.pdf

Inoue E, Kawai K, Imabayashi H. 2005. Species composition and assemblage structure of chironomid larvae (Diptera: Chironomidae) attaching to the artificial substrates in a Japanese temperate basin, in relation to the longitudinal gradient. *Hydrobiologia* 543: 119-133.

Jacobsen D, Shultz R, Encalada A. 1997. Structure and diversity of stream invertebrate assemblages: the effect of temperature with altitude and latitude. *Freshwater Biology* 38: 247-261.

Kleine P, Trivinho-Strixino S. 2005. Chironomidae and other macroinvertebrates of a first order stream: community response after habitat fragmentation. *Acta Limnologica Brasiliensia* 17(1): 81-90.

Krebs CJ. 1999. *Ecological Methodology*, 2nd edition. Benjamin Cummings.

Legendre P, Legendre L. 1998. *Numerical Ecology*, 2nd edition. Elsevier.

Lencioni V, Rossaro B. 2005. Microdistribution of chironomids (Diptera: Chironomidae) in Alpine streams: an autoecological perspective. *Hydrobiologia* 533: 61-76.

Lindegaard C. 1995. Classification of water-bodies and pollution. In: Armitage PD, Cranston, PS, Pinder LCV, Editors. *The Chironomidae. The Biology and Ecology of Non-Biting Midges*. pp. 385-304. Chapman and Hall.

Longhi SJ, Durlo MA, Marchiori JNC. 1982. A vegetação da mata ribeirinha no curso médio do rio Jacuí, RS. *Ciência & Natura* 4: 151-161.

Luoto TP. 2009. Subfossil Chironomidae (Insecta: Diptera) along a latitudinal gradient in Finland: development of a new temperature inference model. *Journal of Quaternary Science* 24: 150-158.

Luoto TP. 2011. The relationship between water quality and chironomid distribution in Finland: A new assemblage-based tool for assessments of long-term nutrient dynamics. *Ecological Indicators* 11: 255-262.

Magurran AE. 2004. *Measuring Biological Diversity*, 2nd edition. Blackwell Sciences.

Maltchik L, Lanés LEK, Stenert C, Medeiros E. 2010. Species-area relationship and environmental predictors of fish communities in coastal freshwater wetlands of southern Brazil. *Environmental Biology of Fishes* 88: 25-30.

Maluf JRT. 2000. Nova classificação climática do Estado do Rio Grande do Sul. *Revista Brasileira de Agrometeorologia* 8(1): 141-150.

Manly BFG. 2000. *Multivariate Statistical Methods: A Primer*, 2nd edition. Chapman and Hall.

Marchiori JC, Longhi SJ, Durlo MA. 1982. A vegetação de capoeira na região do curso médio do rio Jacuí, RS. *Ciência & Natura* 4: 141-150.

Marques MMSM, Barbosa FAR, Callisto M. 1999. Distribution and abundance of Chironomidae (Diptera, Insecta) in an impacted watershed in southeast Brazil. *Revista Brasileira de Biologia* 59(4): 553-561.

Martel N, Rodríguez MA, Bérubé P. 2007. Multi-scale analysis of responses of stream macrobenthos to forestry activities and environmental context. *Freshwater Biology* 52: 85-97.

Marziali L, Giordano D, Armanini MC, Erba S, Toppi E, Buffagni A, Rossaro B. 2009. Responses of Chironomid larvae (Insecta, Diptera) to ecological quality in Mediterranean river mesohabitats (South Italy). *River Research and Applications* 26: 1036-1051.

McKie BG, Pearson RG, Cranston PS. 2005. Does biogeographical history matter? Diversity and distribution of lotic midges (Diptera: Chironomidae) in the Australian Wet Tropics. *Austral Ecology* 30: 1-13.

Morrone JJ. 2006. Biogeographic areas and transition zones of Latin America and the Caribbean islands based on panbiogeographic

and cladistic analyses of the entomofauna. *Annual Review of Entomology* 51: 467-494.

Neri DB, Kotzian CB, Sieglösch AE. 2005. Composição de Heteroptera aquáticos e semi-aquáticos na área de abrangência da U.H.E. Dona Francisca, RS, Brasil: fase de pré-enchimento. *Iheringia, Série Zoologia* 95 (4): 421-428.

Nessimian JL, Sanseverino AS. 1998. Trophic functional characterization of the mountain region of Rio de Janeiro State, Brazil. *Verhandlungen der Internationalen Vereinigung für Theoretische und Angewandte Limnologie* 26: 2115-2119.

Oliver DR. 1971. Life History of the Chironomidae. *Annual Review of Entomology* 16: 211-230.

Panatta A, Stenert C, Freitas SMF, Maltchik L. 2006. Diversity of chironomid larvae in palustrine wetlands of the coastal plain in the south of Brazil. *Limnology* 7: 23-30.

Panatta A, Stenert C, Freitas SMF, Maltchik L. 2007. Diversity and distribution of Chironomid larvae in wetlands in southern Brazil. *Journal of the Kansas Entomological Society* 80(3): 229-242.

Pereira PRB, Garcia Netto LR, Borin CJA. 1989. Contribuição à geografia física do município de Santa Maria: unidades de paisagem. *Geografia Ensino e Pesquisa* 3: 37-68.

Pinder LCV. 1986. Biology of freshwater Chironomidae. *Annual Review of Entomology* 31: 1-23.

Pinder LCV. 1995. The habitats of Chironomid larvae. In: Armitage PD,

Cranston PS, Pinder LCV, Editors. *The Chironomidae. The Biology and Ecology of Non-Biting Midges*. pp. 107-135. Chapman and Hall.

Pinder LCV, Reiss F. 1983. The larvae of Chironominae (Diptera: Chironomidae) of the Holarctic region – keys and diagnoses. In: Wiederholm T, Editor. *Chironomidae of the Holarctic region. Keys and diagnoses*. pp. 293-435. Entomologica Scandinavica Supplement.

Pires MM, Kotzian CB, Spies MR, Neri DB. 2011. *Larval diversity of Odonate (Insecta) genera in a montane region of southern Brazil, and the influence of artificial farm ponds on Odonate community structure*. Monografia de conclusão de curso, Curso de Ciências Biológicas, Universidade Federal de Santa Maria.

Porinchu DF, MacDonald GM. 2003. The use and application of freshwater midges (Chironomidae: Insecta: Diptera) in geographical research. *Progress in Physical Geography* 27: 378-422.

Prado DE. 2000. Seasonally dry forests of tropical South America: from forgotten ecosystems to a new phytogeographic unit. *Edinburgh Journal of Botany* 57(3): 437-461.

Principe RE, Boccolini MF, Corigliano MC. 2008. Structure and spatial-temporal dynamics of Chironomidae fauna (Diptera) in upland and lowland fluvial habitats of the Chocancharava River Basin (Argentina). *Hydrobiology* 93(3): 342-357.

Principe RE, Corigliano MC. 2006. Bentic, drifting and marginal macroinvertebrate assemblages in a low river: temporal and

spatial variations and size structure. *Hydrobiologia* 553: 303-317.

Puntí T, Rieradevall M, Prat N. 2009. Environmental factors, spatial variation, and specific requirements of Chironomidae in Mediterranean reference streams. *Proceedings of the North American Benthological Society* 28 (1): 247-265.

Raunio J. 2008. *The use of Chironomid Pupal Exuvial Technique (CPET) in Freshwater Biomonitoring: Applications for Boreal Rivers and Lakes*. Oulu University.

Resende DLMC, Takeda AM. 2007. Larvas de Chironomidae (Diptera) em três reservatórios do Estado do Paraná. *Revista Brasileira de Zoociências* 9(2): 167-176.

Rios SL, Bailey RC. 2006. Relationship between riparian vegetation and stream benthic communities at three spatial scales. *Hydrobiologia* 553: 153-160.

Rohlf FJ. 2000. NTSYSpc 2.10s. *Numerical Taxonomic and Multivariate Analysis*. Exeter Software.

Romesburg HC. 1984. *Cluster Analysis for Researchers*. Lifetime Learning Publications.

Roque FO, Siqueira T, Bini LM, Ribeiro MC, Tambosi LR, Ciocheti G, Trivinho-Strixino. 2010. Untangling associations between chironomid taxa in Neotropical streams using local and landscape filters. *Freshwater Biology* 37(1): 1-19.

Roque FO, Trivinho-Strixino S, Milan L, Leite JG. 2007. Chironomid species richness in low-order streams in the Brazilian Atlantic Forest: a first approximation through a Bayesian approach. *Journal of the North*

American Benthological Society 26(2): 221-231.

Roque FO, Trivinho-Strixino S, Strixino G, Agostinho RC, Fogo JC. 2003. Benthic macroinvertebrates in streams of the Jaraguá State Park (Southeast of Brazil) considering multiple spatial scales. *Journal of Insect Conservation* 7(2): 63-72.

Rosenberg DM. 1992. Freshwater biomonitoring and Chironomidae. *Aquatic Ecology* 26(2): 101-122.

Rosin GC, Oliveira-Mangarotti DP, Takeda AM, Butakka, CMM. 2009. Consequences of dam construction upstream of the Upper Paraná River Floodplain (Brasil): a temporal analysis of the Chironomidae community over an eight-year period. *Brazilian Journal of Biology* 69(2): 591-608.

Rossaro, B. 1991. Factors that determine Chironomidae species distribution in fresh waters. *Italian Journal of Zoology* 58: 281-286.

Rossaro B, Lencioni V, Boggero A, Marziali L. 2006. Chironomids from Southern Alpine running waters: ecology, biogeography. *Hydrobiologia* 562: 231-246.

Saether OA, Ashe P, Murray DE. 2000. Family Chironomidae. In: Papp L, Darvas B, Editors. *Contributions to a Manual of Palaearctic Diptera (with special reference to the flies of economic importance)*, volume 4 (Appendix A. 6). Science Herald.

Sandin L, Johnson RK. 2004. Local, landscape and regional factors structuring benthic macroinvertebrate assemblages in Swedish streams. *Landscape Ecology* 19: 501-514.

- Sánchez MI, Green AJ, Castellanos, EM. 2006. Spatial and temporal fluctuations in presence and use of chironomid prey by shorebirds in the Odiel salt pans, south-west Spain. *Hydrobiologia* 567: 329–340.
- Sanseverino AM, Nessimian JL. 1998. Habitat preferences of Chironomidae larvae in an upland stream of Atlantic Forest, Rio de Janeiro State, Brazil. *Verhandlungen der Internationalen Vereinigung für Theoretische und Angewandte Limnologie* 26: 2141-2144.
- Sanseverino AM, Nessimian JL. 2001. Habitats de larvas de Chironomidae (Insecta, Diptera) em riachos de Mata Atlântica no Estado do Rio de Janeiro. *Acta Limnologica Brasiliensia* 13(1): 29-38.
- Sanseverino AM, Nessimian JL. 2007. A fauna de Tanytarsini (Insecta: Diptera: Chironomidae) em áreas de terra firme na Amazônia Central. *Anais VIII, Congresso de Ecologia do Brasil* 1-2.
- Sanseverino AM, Nessimian JL, Oliveira ALH. 1998. A fauna de Chironomidae (Diptera) em diferentes biótopos aquáticos na Serra do Subaio (Teresópolis, RJ). In: Nessimian JL, Carvalho AL, Editors. *Ecologia de Insetos Aquáticos. Série Oecologia Brasiliensis* 5: 253-263.
- Santos CM, Henry R. 2001. Composição, distribuição e abundância de Chironomidae (Diptera, Insecta) na Represa de Jurumirim (Rio Paranapanema – SP). *Acta Limnologica Brasiliensia* 13(2): 99-115.
- Santos TG, Vasconcelos TS, Rossa-Feres DC, Haddad CFB. 2009. Anurans of a seasonally dry tropical forest: Morro do Diabo State Park, São Paulo state, Brazil. *Journal of Natural History* 43(15): 973-993.
- Serrano MAS, Severi W, Toledo VJS. 1998. Comunidade de Chironomidade e outros macroinvertebrados em um rio tropical de planície, Rio Bento Gomes, MT. In: Nessimian JL, Carvalho AL, Editors. *Ecologia de Insetos Aquáticos. Série Oecologia Brasiliensis* 5: 265-278.
- Siegloch AE, Froehlich CG, Kotzian CB. 2008. Composition and diversity of Ephemeroptera (Insecta) nymph communities in the middle section of the Jacuí River and some tributaries, southern Brazil. *Iheringia, Série Zoologia* 98(4): 425-432.
- Silva FL, Moreira DC, Ruiz SS, Bochini GL. 2007. Avaliação da importância da unidade de conservação na preservação da diversidade de Chironomidae (Insecta: Diptera) no córrego Vargem Limpa, Bauru, Estado de São Paulo. *Acta Scientiarum Biological Sciences* 29(4): 401-405.
- Silva FL, Ruiz SS, Moreira DC, Bochini GL. 2008. Composição e diversidade de imaturos de Chironomidae (Insecta, Diptera) no Ribeirão dos Peixes, Dois Córregos, SP. *Revista Brasileira de Biociências* 6(4): 341-346.
- Silva FL, Ruiz SS, Moreira DC, Bochini GL. 2008. Composição e diversidade de imaturos de Chironomidae (Insecta, Diptera) no Ribeirão dos Peixes, Dois Córregos, SP. *Revista Brasileira de Biociências* 6(4): 341-346.
- Silva FL, Silveira AL, Talamoni JLB, Ruiz SS. 2009. Temporal variation of Chironomidae larvae (Insecta, Diptera) in the

Batalha River, Midwestern São Paulo State, Brazil. *Ciência et Praxis* 2(3): 7-12.

Simião-Ferreira J, Demarco JP, Mazão GR, Carvalho AR. 2009. Chironomidae assemblage structure in relation to organic enrichment of an aquatic environment. *Neotropical Entomology* 38(4): 464-471.

Siqueira T, Bini LM, Cianciaruso MV, Roque FO, Trivinho-Strixino S. 2009. The role of niche measures in explaining the abundance distribution relationship in tropical lotic chironomids. *Hydrobiologia* 636(1): 163-172.

Siqueira T, Roque FO, Trivinho-Strixino S. 2008. Phenological patterns of Neotropical lotic Chironomids: Is emergence constrained by environmental factors? *Austral Ecology* 33(7): 902-910.

Siqueira T, Trivinho-Strixino S. 2005. Diversidade de Chironomidae (Diptera) em dois córregos de baixa ordem na região central do Estado de São Paulo, através da coleta de exúvias de pupa. *Revista Brasileira de Entomologia* 49(4): 531-534.

Sokal RR, Michener CD. 1958. A statistical method for evaluating systematic relationships. *The University of Kansas Scientific Bulletin* 38: 1409-1438.

Smouse PE, Long JC, Sokal RR. 1986. Multiple regression and correlation extensions of the Mantel test of matrix correspondence. *Systematic Zoology* 35: 627-632.

Sonoda KC, Matthaei CD, Trivinho-Strixino S. 2009. Contrasting land uses affect Chironomidae communities in two Brazilian rivers. *Archiv für Hydrobiologie, Fundamental and Applied Limnology* 174(2): 173-184.

Spies MR, Froehlich CG, Kotzian CB. 2006. Composition and diversity of Trichoptera (Insecta) larvae communities in the middle section of the Jacuí River and some tributaries, State of Rio Grande do Sul, Brazil. *Iheringia, Série Zoologia* 96(4): 389-398.

Spies M, Reiss F. 1996. Catalog and bibliography of Neotropical and Mexican Chironomidae (Insecta, Diptera). *Spixiana Supplement* 22: 61-119.

Statzner B, Higler B. 1986. Stream hydraulics as a major determinant of benthic invertebrate zonation patterns. *Freshwater Biology* 16: 127-139.

Stenert C, Santos EM, Maltchik L. 2004. Levantamento da diversidade de macroinvertebrados em áreas úmidas do Rio Grande do Sul (Brasil). *Acta Biologica Leopoldensia* 26(2): 225-240.

Strahler AN. 1957. Quantitative analysis of watershed geomorphology. *American Geophysical Union Transactions* 33: 913-920.

Suriano MT, Fonseca-Gessner AA. 2004. Chironomidae (Diptera) larvae in streams of Parque Estadual de Campos do Jordão, São Paulo State, Brazil. *Acta Limnologica Brasiliensia* 16(2): 129-136.

Takeda AM, Butakka CMM, Fujita DS, Fugita RH, Bibian JPR. 2005. Larvas de Chironomidae em cascata de reservatórios no Rio Iguaçu (PR). In: Rodrigues L, Thomaz SM, Agostinho AA, Gomes LC, Editors. *Biocenose em Reservatórios: Padrões Espaciais e Temporais*. pp.147-160. Rima.

Takeda AM, Higuti J, Rodrigues LC, Bialecki A, Pilati R, Delariva, RL, Abes SS,

- Campos JRC, Campos JB, Romagnolo MB, Sanches PV. 1997. Distribuição espacial de zoobentos do Ressaco do “Pau Véio” e do canal Cortado. In: *Anais VIII Seminário Regional de Ecologia*. pp. 127-139. Universidade Federal de São Carlos.
- Tokeshi M. 1995. Life cycles and population dynamics. In: Armitage PD, Cranston PS, Pinder LCV, Editors. *The Chironomidae: The Biology and Ecology of Non-Biting Midges*. pp. 226-268. Chapman and Hall.
- Tokeshi M, Pinder LCV. 1985. Microhabitats of stream invertebrates on two submersed macrophytes with contrasting leaf morphology. *Holarctic Ecology* 1(8): 313-319.
- Trivinho-Strixino S. 2011. *Chironomidae (Insecta, Diptera, Nematocera) do Estado de São Paulo, Sudeste do Brasil*. Available on line: <http://www.biotaneotropica.org.br/v11n1a/en/abstract?inventory+bn0351101a2011>
- Trivinho-Strixino S, Strixino G. 1995. *Larvas de Chironomidae (Diptera) do Estado de São Paulo. Guia de Identificação e Diagnose dos Gêneros*. Universidade Federal de São Carlos.
- Trivinho-Strixino S, Strixino G. 1999. Insetos dípteros: quironomídeos. In: Joly CA, Bicudo CEM, Editors. *Biodiversidade do Estado de São Paulo, Brasil: síntese do conhecimento ao final do século XX*, 4: Invertebrados de Água Doce. pp. 141-148. FAPESP
- Trivinho-Strixino S, Strixino G. 2005. Chironomidae (Diptera) do Rio Ribeira (divisa dos Estados de São Paulo e Paraná) numa avaliação ambiental faunística. *Entomologia y Vectores* 12(2): 243-253.
- Valentin JL. 1995. Agrupamento e ordenação. In: Peres-Neto RP, Valentin JL, Fernando F, Editors. *Tópicos em Tratamento de Dados Biológicos. Oecologia Brasiliensis* 2: 27-55.
- Valentin JL. 2000. *Ecologia Numérica: uma Introdução à Análise Multivariada de Dados Ecológicos*. Interciência.
- Vinson MR, Hawkins CP. 1998. Biodiversity of stream insects: Variation at local, basin, and regional scales. *Annual Review of Entomology* 43: 271-293.
- Voelz NJ, McArthur JV. 2000. An exploration of factors influencing lotic species richness. *Biodiversity and Conservation* 9: 1543-1570.
- Walker I. 1998. Population dynamics of Chironomidae (Diptera) in the central Amazonian blackwater river Tarumã-Mirim (Amazonas, Brazil). In: Nessimian JL, Carvalho AL, Editors. *Ecologia de Insetos Aquáticos. Oecologia Brasiliensis* 5: 235-252.
- Wiedenbrug S, Nolte U, Würdig NL. 1997. Macrozoobenthos of a coastal lake in southern Brazil *Achiv für Hydrobiologie* 140(4): 533-548.
- Wiedenbrug S, Trivinho-Strixino S. 2009. *Ubatubaneura*, a new genus of the *Corynoneura* group (Diptera: Chironomidae: Orthoclaadiinae) from the Brazilian Atlantic Forest. *Zootaxa* 1993(41): 41-52.
- Wiederholm T, 1983. Chironomidae of the Holarctic region key and diagnoses. *Entomologica Scandinavica Supplement* 34(1): 125-127.
- Wolda H. 1981. Similarity indices, sample size and diversity. *Oecologia* 50: 296-302.

Woodcock T, Longcore J, McAuley D, Mingo T, Bennatti CR, Stromborg K. 2005. The role of pH in structuring communities of Maine wetland macrophytes and Chironomid larvae (Diptera). *Wetlands* 25(2): 306-316.

Table 1. Characterization of the sampling sites of the Chironomidae larvae assemblages collected between April 2000 and May 2002 in the middle course of the Jacuí River, Rio Grande do Sul, Brasil.

Site	Site Coordinates	Altitude (m)	Location	Order	# of samplings	Width (m)	Depth (m)	Description of sites
1	29° 28' 44" S; 53° 16' 56" W	57	Jacuí River (semi-regulated course)	7	17	200	0.47	Arboreal vegetation, slightly shadowed, rocky bottom, and anthropic influence
2	29° 28' 03" S; 53° 13' 28" W	180	Lajeado do Gringo Stream (non-regulated course)	4	2	9	0.39	Arboreal vegetation in one of the banks, shadowed, presence of <i>Podostemum</i> sp. only in September 2000
3	29° 28' 07" S; 53° 13' 28" W	140	Lajeado do Gringo Stream (non-regulated course)	4	16	5	0.23	Arboreal vegetation in one of the banks, poorly shadowed
4	29° 25' 21" S; 53° 14' 51" W	71	Jacuí River (regulated course)	7	3	70	0.48	Arboreal vegetation in one of the banks, poorly shadowed
5	29° 23' 01" S; 53° 13' 19" W	152	Lajeado da Gringa Stream (non-regulated course)	3	4	13.5	0.38	Arbustive vegetation in one of the banks, not shadowed, presence of <i>Podostemum</i> sp.
6	29° 23' 02" S; 53° 12' 34" W	94	Lajeado da Gringa Stream (non-regulated course)	3	2	6	0.45	Arbustive vegetation in one of the banks, not shadowed, presence of <i>Podostemum</i> sp..
7	29° 22' 57" S; 53° 12' 08" W	107	Lajeado da Gringa Stream (non-regulated course)	3	17	6	0.44	Arbustive vegetation in one of the banks, not shadowed, presence of <i>Podostemum</i> sp.
8	29° 22' 16" S; 53° 13' 41" W	107	Lajeado do Tigre Stream (non-regulated course)	2	3	6	0.28	Arboreal vegetation and shadowing in one of the banks.
9	29° 21' 53" S; 53° 14' 07" W	140	Lajeado do Tigre Stream (non-regulated course)	1	3	3	0.3	Well-preserved riparian vegetation in one of the banks, well shadowed
10	29° 20' 32" S; 53° 09' 57" W	95	Carijinho River (non-regulated course)	4	2	8	0.46	Arboreal vegetation in one of the banks, poorly shadowed.
11	29° 21' 26" S; 53° 09' 11" W	92	Carijinho River (non-regulated course)	4	16	8	0.55	Arbustive vegetation in one of the banks, not shadowed, presence of <i>Podostemum</i> sp.
12	29° 19' 05" S; 53° 14' 20" W	101	Lajeado Estaleiro Stream (non-regulated course)	3	1	6	0.4	Well-preserved riparian vegetation in both banks, well shadowed, presence of <i>Podostemum</i> sp.

Table 2. Ecosystems, hydrographic regions, biomes, and respective locations (see references) used in the comparison between the Chironomidae assemblages with those recorded in the middle course of the Jacuí River, Rio Grande do Sul. Hydrographic regions: Bacia Amazônica (Amazon Basin) – AM, Bacia do Prata (Plate River Basin) – BP, Bacia Costeira do Sudeste (Southeast Coastal Basins) – BCSe and Bacia Costeira Sul (South Coastal Basins) – BCS. Biomes: Amazônia – AM, Cerrado – CR, Pantanal – PN, Mata Atlântica – MT and Pampa – PM. Sampler: Core – 1, Manual grab – 2, Surber – 3, Entomological aquatic net – 4, Modified Petersen grab – 5, Manual net – 6, Eckman-Birge grab – 7, D-frame net – 8, Van Veen grab – 9.

Sites	Locations	State	Coordinates	Average altitude (m)	Ecosystems (lotic/lentic)	Hydrographic regions	Biomes	Collector	# of Samplings				Abbreviations
									1	4	12	> 12	
1	Porto Trombeta (Callisto et al. 1998)	PA	01° 30' 00" S; 56° 20' 00" W	2	lotic	BAM	AM	1	X				BAAM
2	Rio Abraão (Simião-Ferreira et al. 2009)	GO	16° 24' 45" S; 48° 57' 33" W	1130	lotic	BP	CR	3	X				BPCR1
3	Rio Bento Gomes (Serrano et al. 1998)	MT	15° 40' 00" S; 56° 30' 00" W	300	lotic	BP	PN	4	X				BPPAN1
4	Rio Paraguaí (Aburaya et al. 2007)	MT	16° 11' 42" S; 57° 40' 51" W	158	lotic	BP	PN	5		X			BPPAN2
5	Lagoa Guarará, Patos, Rio Ivinhema (Higuti and Takeda 2002)	MS	22° 43' 23" S; 53° 18' 22" W	237	lotic/lentic	BP	MT	5		X			BPMT1
6	Serra do Subaio, Rio Canoas (Sanseverino et al. 1998)	RJ	22° 22' 00" S; 43° 01' 08" W	1100	lotic	BCSe	MT	3, 2, 6	X				BCSeMT1
7	Rio Fazenda (Henriques-Oliveira 2003)	RJ	22° 55' 00" S; 43° 19' 00" W	400	lotic	BCSe	MT	3	X				BCSeMT2
8	Ribeirão dos Peixes (Silva et al. 2008)	RJ	22° 22' 00" S; 48° 22' 00" W	697	lotic/lentic	BCSe	MT	7	X				BCSeMT3
9	Rio Cascatinha (Henriques-Oliveira et al. 1999)	RJ	22° 16' 48" S; 42° 33' 05" W	1110	lotic	BCSe	MT	3	X				BCSeMT4
10	Rio Paquequer (Nessimian and Sanseverino et al. 1999)	RJ	22° 22' 03" S; 42° 59' 57" W	1110	lotic	BCSe	MT	3	X				BCSeMT5
11	11 riachos - Serra do Mar (Sanseverino et al. 1998)	RJ	22° 26' 11" S; 43° 36' 31" W	1100	lotic	BCSe	MT	3, 2, 8	X				BCSeMT6
12	Represa Jurumirim (Santos and Henry 2001)	SP	23° 12' 17" S; 49° 13' 13" W	636	lentic	BP	MT	5		X			BPMT2
13	Córrego Galharada (Suriano and Fonseca-Gessner 2004)	SP	22° 35' 00" S; 45° 27' 00" W	1650	lotic	BP	MT	3	X				BPMT3
14	Reservatório Salto Grande (Dorfeld et al. 2005)	SP	22° 44' 00" S; 47° 20' 00" W	530	lentic	BP	MT	7		X			BPMT4
15	Cabeceira do Rio Iguaçu 1 (Takeda et al. 2005)	SP	25° 26' 38" S; 49° 17' 44" W	940	lotic	BP	MT	5	X				BPMT5
16	Médio - Rio Iguaçu 2 (Takeda et al. 2005)	SP	26° 01' 24" S; 51° 25' 29" W	882	lotic	BP	MT	5	X				BPMT6
17	Médio - Rio Iguaçu 3 (Takeda et al. 2005)	SP	25° 29' 28" S; 51° 32' 35" W	991	lotic	BP	MT	5	X				BPMT7
18	Foz do Rio Iguaçu 4 (Takeda et al. 2005)	SP	25° 31' 44" S; 53° 15' 43" W	508	lotic	BP	MT	5	X				BPMT8
19	Vale do Rio Ribeira (Trivinho-Strixino and Strixino 2005)	SP	24° 31' 21" S; 53° 15' 43" W	502	lotic	BP	MT	8		X			BPMT9
20	Ribeirão Cachim (Freire and Fonseca-Gessner 2001)	SP	21° 57' 35" S; 47° 50' 16" W	839	lotic	BP	CR	3		X			BPCR2
21	Córrego Vargem Limpa (Silva et al. 2007)	SP	22° 19' 18" S; 49° 04' 13" W	530	lotic	BP	CR	7	X				BPCR3
22	Bacia Rio Jacaré-Guaçu, 1 (Corbi and Trivinho-Strixino 2008)	SP	21° 57' 00" S; 48° 20' 00" W	579	lotic	BP	CR	9		X			BPCR4
23	Bacia Rio Jacaré-Guaçu, 2 (Corbi and Trivinho-Strixino 2008)	SP	21° 51' 00" S; 48° 21' 00" W	511	lotic	BP	CR	9		X			BPCR5
24	Bacia Rio Jacaré-Guaçu, 3 (Corbi and Trivinho-Strixino 2008)	SP	22° 00' 00" S; 47° 31' 00" W	734	lotic	BP	CR	9		X			BPCR6
25	Rio Batalha (Silva et al. 2009)	SP	22° 20' 00" S; 49° 00' 00" W	593	lotic	BP	CR	7			X		BPCR7
26	Ressaca do " Pau Véio" (Takeda et al. 1997)	PR	25° 11' 53" S; 48° 58' 15" W	792	lotic	BP	MT	4	X				BPMT10
27	Reservatório Capivari (Resende and Takeda 2007)	PR	22° 36' 00" S; 52° 50' 00" W	254	lentic	BCSe	MT	4		X			BCSeMT7
28	Reservatório Rosana (Resende and Takeda 2007)	PR	25° 25' 10" S; 49° 06' 49" W	884	lentic	BP	MT	4		X			BPMT11
29	Reservatório Irai (Resende and Takeda 2007)	PR	25° 46' 20" S; 53° 16' 03" W	476	lentic	BP	MT	4		X			BPMT12
30	Rio Paraná (Rosin et al. 2009)	PR	23° 17' 27" S; 51° 14' 53" W	615	lotic	BP	MT	5			X		BPMT13
31	Rio Ivinhema (Rosin et al. 2009)	PR	22° 47' 18" S; 53° 19' 16" W	237	lotic/lentic	BP	MT	5			X		BPMT14
32	Jacuí (this study)	RS	29° 28' 03" S; 53° 13' 28" W	137	lotic	BP	MT	2			X		BCSMT
33	Planície Costeira (Panatta et al. 2006)	RS	30° 43' 40" S; 51° 18' 45" W	7	lentic	BCS	PM	9	X				BCSPM

Table 3. Taxonomic composition and abundance of the Chironomidae larvae assemblages at 12 sampling sites (1-12) in the middle course of the Jacuí River basin, Rio Grande do Sul, between April 2000 and May 2002. Genera and species identified by capital letters and with Arabic numerals, respectively, correspond to those described in the dichotomous key of Trivinho-Strixino and Strixino (1995), and genera and species identified by Arabic numbers and capital letters, respectively, correspond to those identified by the authors of the present study.

Subfamily/Taxa	Sampling sites											
	1	2	3	4	5	6	7	8	9	10	11	12
Chironominae												
<i>Aedokritus</i> Roback 1958	3	0	0	0	0	0	18	0	0	0	0	0
<i>Axarus</i> Roback 1980	0	1	0	0	0	0	5	0	0	0	0	0
<i>Beadius</i> sp. 1 Trivinho-Strixino and Strixino 1995	0	0	2	0	0	0	17	0	0	0	5	0
<i>Cladopelma</i> Kieffer 1921	0	0	0	0	0	0	85	0	0	0	3	0
<i>Cf. Pelomus</i>	4	0	0	0	0	0	112	0	0	0	0	0
<i>Dicrotendipes</i> sp. 2 Trivinho-Strixino and Strixino 1995	14	0	0	0	0	0	69	0	0	0	0	0
<i>Dicrotendipes</i> sp. 3 Trivinho-Strixino and Strixino 1995	15	0	12	0	0	2	123	0	0	0	0	0
<i>Cf. Dicrotendipes</i>	0	0	1	0	0	0	0	0	0	0	0	0
<i>Endotribelos</i> Grodhaus 1987	2	0	3	0	0	0	0	0	0	0	9	0
<i>Endotribelos</i> cf. <i>hespellium</i>	0	0	18	0	0	0	3	0	0	0	36	0
<i>Fissimentum</i> Cranston and Nolte	0	0	0	0	0	0	0	0	0	0	15	0
Chironomini Gen. 1	0	0	2	0	0	0	9	0	0	0	0	0
Chironomini Gen. 2	0	0	0	0	0	0	9	0	0	0	0	0
Chironomini Gen. 3	1	0	0	0	0	0	0	0	0	0	0	0
Chironomini Gen. D	0	0	0	0	0	0	0	0	0	0	0	1
<i>Chironomus</i> gr. <i>Riparius</i> Trivinho-Strixino and Strixino 1995	2	0	0	0	0	0	0	0	0	0	0	0
<i>Goeldichironomus pictus</i> Reiss 1974	9	0	0	0	0	0	0	0	0	0	0	0
<i>G. serratus</i> Reiss 1974	0	0	9	0	0	0	0	0	0	0	4	0
<i>G. xiborena</i> Reiss 1974	0	0	0	0	0	0	0	0	0	0	8	0
<i>Harnischia</i> (?) Kieffer 1921	0	0	12	0	0	0	31	0	0	0	0	0
<i>Kiefferulus</i> (?) Goetghebuer 1922	0	0	11	0	1	0	18	0	0	0	6	0
<i>Lauterborniella</i> Thienemann and Bause in Bause 1913	0	0	0	0	0	0	11	0	0	0	0	0
<i>Manoa</i> Fittkau 1963	0	0	6	0	0	0	1	3	0	0	7	0
<i>Nilothauma</i> sp. 1 Trivinho-Strixino and Strixino 1995	0	0	5	0	0	0	0	0	0	0	1	0
<i>Nilothauma</i> sp. 2 (?) Trivinho-Strixino and Strixino 1995	0	0	0	0	0	0	7	0	0	0	0	0
<i>Nimbocera</i> sp. 3 Trivinho-Strixino and Strixino 1995	0	0	4	0	0	0	0	0	0	0	0	0
<i>Oukuriella</i> Epler 1986	0	0	1	1	0	0	0	0	0	0	5	0
<i>Parachironomus</i> sp. 1 Trivinho-Strixino and Strixino 1995	16	0	0	0	0	0	0	0	0	0	0	0
<i>Parachironomus</i> sp. 2 Trivinho-Strixino and Strixino 1995	31	0	0	0	0	0	0	0	0	0	15	0
<i>Parachironomus</i> sp. 3 Trivinho-Strixino and Strixino 1995	0	0	1	0	1	0	0	0	0	0	0	0
<i>Paralauterborniella</i> Lenz 1941	0	0	1	0	0	0	0	0	0	0	0	0
<i>Paratendipes</i> Kieffer 1911	26	0	33	4	0	1	4	1	0	0	71	0
<i>Phaenospsectra</i> Kieffer 1921	3	0	0	0	0	0	5	0	0	0	25	0
<i>Polypedilum</i>	2	0	1	0	0	0	0	0	0	0	2	0
<i>P.(Asheum)</i> Trivinho-Strixino and Strixino 1995	4	0	0	0	0	0	0	0	0	0	0	0

<i>P. fallax</i> Trivinho-Strixino and Strixino 1995	0	0	1	0	0	0	1	0	0	0	23	0
<i>P. (Polypedilum)</i> sp.	0	0	3	0	0	0	0	0	0	0	0	0
<i>P. (Polypedilum)</i> sp. 1 Trivinho-Strixino and Strixino 1995	33	0	200	0	1	0	298	0	1	1	365	0
<i>P. (Polypedilum)</i> sp. 2 Trivinho-Strixino and Strixino 1995	0	0	216	1	5	0	276	3	0	2	296	0
<i>P. (Polypedilum)</i> sp. A	0	0	1	0	0	0	21	0	0	0	23	0
<i>P. (Polypedilum)</i> spp.	0	0	1	0	0	0	0	0	0	0	0	0
<i>P. (Tripodura)</i> Kieffer 1912	6	0	1	0	0	0	1	0	0	0	0	0
<i>Stenochironomus</i> Kieffer 1919	4	0	5	0	0	0	6	0	0	0	37	0
<i>Stenochironomus</i> complex	0	0	1	1	0	0	2	0	0	0	1	0
Cf. <i>Stenochironomus</i>	0	0	1	0	0	0	0	0	0	0	0	0
<i>Xenochironomus</i> complex	0	0	0	0	0	0	0	0	0	0	0	2
<i>Xestochironomus</i> Sublette and Wirth 1972	0	0	1	0	0	0	0	0	0	0	9	0
<i>Xenochironomus</i> complex	0	0	0	0	0	0	0	0	0	0	2	0
<i>Zavreliella</i> sp. 1 Trivinho-Strixino and Strixino 1995	0	0	4	0	0	0	1	0	4	0	0	0
<i>Zavreliella</i> sp. 2 Trivinho-Strixino and Strixino 1995	2	0	0	0	0	0	0	0	0	0	0	0
<i>Pseudochironomini</i>	0	0	1	0	0	0	0	0	0	0	0	0
<i>Pseudochironomini</i> Gen. 1	3	0	0	0	0	0	31	0	0	0	2	0
<i>Pseudochironomini</i> sp. 3	0	0	1	0	0	0	0	0	0	0	0	0
<i>Pseudochironomus</i> ? Malloch 1915	4	0	116	0	34	40	408	0	0	0	205	14
<i>Rheotanytarsus</i> sp.	0	0	1	0	0	0	0	0	0	0	0	0
<i>Rheotanytarsus</i> sp. 1 Trivinho-Strixino and Strixino 1995	169	1	179	0	6	0	1653	2	6	2	252	0
<i>Rheotanytarsus</i> sp. 2 Trivinho-Strixino and Strixino 1995	47	16	76	0	2	0	542	3	0	0	162	0
<i>Caladomyia</i> sp.	0	0	0	0	0	0	8	0	0	0	7	0
<i>C. friederi</i> Trivinho-Strixino and Strixino 2000	0	0	0	0	0	0	13	0	0	0	0	0
<i>C. ortonii</i> Sæwædal 1981	2	0	3	0	1	0	3	0	0	0	2	0
<i>Tanytarsini</i> Gen. A Trivinho-Strixino and Strixino 1995	0	0	4	0	0	0	0	0	0	0	0	0
<i>Tanytarsini</i> Gen. D Trivinho-Strixino and Strixino 1995	0	0	0	0	0	0	0	0	0	0	48	0
<i>Tanytarsini</i> Gen. C Trivinho-Strixino and Strixino 1995	0	0	1	0	0	0	0	0	0	0	5	0
<i>Tanytarsus</i> van der Vulp 1874	1	0	0	0	0	0	89	0	0	0	1	0
<i>T. rhabdomantis</i> Trivinho-Strixino and Strixino 1991	8	0	4	0	0	0	0	0	3	0	0	0
Tanypodinae												
<i>Ablabesmyia</i> (karelia) Trivinho-Strixino and Strixino 1995	7	9	130	1	49	27	213	0	1	0	117	1


<i>Djalmabatista</i> sp.2 Trivinho-Strixino and Strixino 1995	0	0	0	0	0	0	1	0	0	0	1	0
<i>Labrundinia</i> Fittkau, 1962	0	0	0	0	0	0	0	0	0	0	13	0
<i>Larsia</i> Fittkau, 1962	0	0	5	0	0	0	114	0	0	0	4	0
<i>Monopelopia</i> Fittkau, 1962	0	0	0	0	0	0	19	0	0	0	0	0
<i>Pentaneura</i> Philippi, 1865	0	0	4	0	0	0	89	1	0	0	0	0
Cf. <i>Pentaneura</i>	0	0	1	0	0	0	0	0	0	0	0	0
<i>Thienemannimyia</i> (gr.) Fittkau 1957	0	0	0	0	0	0	4	0	0	0	0	0
<i>Thinemannimyia</i> (?) sp. 1 Trivinho-Strixino and Strixino 1995	0	0	3	0	0	0	0	0	0	0	0	0
<i>Thinemannimyia</i> (?) sp. 2 Trivinho-Strixino and Strixino 1995	0	0	1	0	0	0	3	0	0	0	0	0
<i>Zavrelimyia</i> Fittkau 1962	0	0	0	0	0	0	9	0	0	0	1	0
Orthoclaadiinae												
<i>Corynoneura</i> sp. 1 Trivinho-Strixino and Strixino 1995	19	0	0	0	0	0	6	0	0	0	1	0
<i>Corynoneura</i> sp. 2 Trivinho-Strixino and Strixino 1995	135	0	4	0	0	0	0	0	0	0	12	0
<i>Onconeura</i> sp. Trivinho-Strixino and Strixino 1995	0	0	0	0	0	0	1	0	0	0	1	0
<i>Thienemanniella</i> sp. 2 Trivinho-Strixino and Strixino 1995	286	0	4	0	0	0	81	0	0	0	39	0
<i>Thienemanniella</i> (?) sp. 3 Trivinho-Strixino and Strixino 1995	32	0	12	0	0	0	15	0	0	0	0	0
<i>Parametricnemus</i> Gotegebeuer 1932	15	0	7	0	0	1	18	0	1	0	3	0
<i>Cardiocladius</i> (?) Kieffer 1912	2	0	6	6	0	0	10	0	0	0	12	0
<i>Cricotopus</i> sp. 1 Trivinho-Strixino and Strixino 1995	294	0	117	17	16	5	896	3	2	0	38	0
<i>Cricotopus</i> sp. 2 Trivinho-Strixino and Strixino 1995	345	40	510	62	100	23	524	87	0	98	205	92
<i>Cricotopus</i> spp.	0	0	2	0	0	0	0	0	0	0	0	0
<i>Cricotopus</i> sp. A	5	0	2	0	0	0	2	0	0	0	0	0
prox. <i>Cricotopus</i> (?) Trivinho-Strixino and Strixino	23	0	0	0	0	0	0	0	0	0	3	0
Cf. <i>Orthocladus</i>	0	0	1	0	0	0	0	0	0	0	0	0
<i>Orthoclaadiini</i> Gen. 1	1	0	34	0	0	0	0	0	0	0	0	0
<i>Lopescladius</i> Oliveira 1967	6	0	7	0	0	0	8	0	0	0	30	0
<i>Metriocnemus</i> van der Wulp 1874	0	0	0	0	0	0	1	0	0	0	0	0
<i>Nanocladius</i> Kieffer 1913	11	2	0	0	1	0	11	0	0	0	21	1
<i>Paracladius</i> Hirvenoja 1973	11	0	31	0	6	4	10	11	0	0	0	0
<i>Rheocricotopus</i> Thienemann and Harnisch 1932	0	0	1	0	0	0	1	0	0	0	0	0
<i>Thienemanniella</i> (?) sp. 1 Trivinho-Strixino and Strixino 1995	0	0	0	0	0	0	1	0	0	0	1	0
<i>Thienemanniella</i> (?) sp. 2 Trivinho-Strixino and Strixino 1995	2	1	0	2	0	0	0	0	0	0	0	0
<i>Thienemanniella</i> (?) sp. 3 Trivinho-Strixino and Strixino 1995	0	0	0	0	1	0	0	0	0	0	0	0
Number of larvae	1605	70	1825	95	224	103	5917	114	18	103	2154	111
Richness	41	6	58	11	13	8	55	9	7	4	49	6


Table 4. Taxonomic composition and frequency of occurrence of the genera of Chironomidae in the five groups formed in the cluster analysis for the 33 locations compared. (Note. i, ii, iii, iv and v = groups formed in the cluster; Arabic numerals in parentheses = number of locations involved in the formation of the groups; bold numbers = taxa with occurrence frequency $\geq 70\%$; italic numbers = exclusive taxa).


Taxa	Group (# of locations)				
	i(10)	ii(9)	iii(6)	iv(3)	v(2)
<i>Aedokritus</i>	60	78.9	0	0	0
<i>Apedilum</i>	50	22	0	0	0
<i>Axarus</i>	50	11	0	0	100
<i>Beardius</i>	70	44	66.6	66.6	0
<i>Caladomyia</i>	80	55	0	33.3	0
<i>Chironomus</i>	90	78	33.3	100	100
<i>Cladopelma</i>	60	100	0	66.6	0
<i>Cladotanytarsus</i>	0	0	0	0	0
<i>Cryptochironomus</i>	90	88.9	83.3	66.6	100
<i>Cryptotendipes</i>	<i>10</i>	0	0	0	0
<i>Demicryptochironomus</i>	30	0	0	0	50
<i>Dicrotendipes</i>	30	88.9	0	33.3	50
<i>Endotribelos</i>	80	55.5	33.3	33.3	0
<i>Fissiomentum</i>	90	100	16.6	0	0
<i>Glyptotendipes</i>	0	0	0	0	100
<i>Goeldichironomus</i>	70	44	33.3	66.6	50
<i>Harnischia</i>	70	88.9	0	66.6	0
<i>Kiefferulus</i>	10	11	0	0	50
<i>Lauterborniella</i>	40	0	66.6	33.3	0
<i>Manoa</i>	<i>20</i>	0	0	0	0
<i>Micropsectra</i>	0	0	0	0	<i>50</i>
<i>Nilothauma</i>	20	11	33.3	100	0
<i>Nilotanypus</i>	0	0	33.3	0	0
<i>Nimbocera</i>	20	0	83.3	0	100
<i>Oukuriella</i>	20	0	83.3	0	0

<i>Parachironomus</i>	60	22	16.6	66.6	50
<i>Paracladopelma</i>	0	0	0	0	50
<i>Paralauterborniella</i>	50	66.6	0	0	50
<i>Paratanytarsus</i>	0	0	0	0	50
<i>Paratendipes</i>	50	11	50	0	0
<i>Pedionomus</i>	1	0	0	0	0
<i>Pelomus</i>	30	22	83.3	0	0
<i>Phaenospsectra</i>	100	100	100	100	100
<i>Polypedilum</i>	30	11	50	0	0
<i>Pseudochironomus</i>	100	33	100	66.6	0
<i>Rheotanytarsus</i>	20	0	0	0	0
<i>Robackia</i>	30	22	16.6	0	50
<i>Shaeteria</i>	10	0	0	0	0
<i>Stempellina</i>	20	33	16.6	0	0
<i>Stempellinella</i>	0	0	100	33.3	0
<i>Stenochironomus</i>	70	22	66.6	66.6	0
<i>Stictochironomus</i>	10	0	16.6	0	0
<i>Tanytarsus</i>	90	100	83.3	66.6	100
<i>Thienemanniella</i>	0	0	16.6	0	0
<i>Tribelos</i>	20	0	0	33.3	0
<i>Xenochironomus</i>	20	0	0	0	0
<i>Xestochironomus</i>	10	11	50	0	0
<i>Zavreliella</i>	50	0	50	33.3	0
<i>Ablabesmyia</i>	100	88.9	33.3	100	50
<i>Alotanypus</i>	0	0	0	100	50
<i>Brundiniella</i>	10	0	0	0	0

<i>Clinotanypus</i>	70	11	0	33.3	0
<i>Coelotanypus</i>	70	55.5	0	33.3	0
<i>Denopelopia</i>	10	0	0	0	0
<i>Djalmabatista</i>	70	88.9	16.6	66.6	100
<i>Fittkauimyia</i>	20	0	0	33.3	0
<i>Labrundinia</i>	40	11	16.6	100	0
<i>Larsia</i>	60	0	16.6	100	0
<i>Macropelopia</i>	0	0	0	33.3	0
<i>Monopelopia</i>	10	0	0	66.6	0
<i>Pentaneura</i>	40	11	16.6	33.3	0
<i>Procladius</i>	60	44.4	0	33.3	0
<i>Tanypus</i>	20	66	0	66.6	0
<i>Thienemannimyia</i>	30	0	0	33.3	0
<i>Zavreliomyia</i>	20	0	0	0	0
<i>Cardiocladius</i>	20	0	33.3	0	0
<i>Corynoneura</i>	100	22	100	0	0
<i>Cricotopus</i>	90	88.9	66.6	66.6	0
<i>Gymnotriocnemus</i>	30	0	16.6	0	0
<i>Lopescladius</i>	70	44	100	0	0
<i>Mesosmittia</i>	0	0	16.6	0	0
<i>Metriocnemus</i>	10	0	66.6	0	0
<i>Nanocladius</i>	70	11	100	33.3	0
<i>Onconeura</i>	10	0	0	0	0
<i>Orthocladius</i>	0	0	0	0	0
<i>Paracladius</i>	10	0	0	0	0
<i>Parakiefferiella</i>	10	0	0	0	50
<i>Parametricnemus</i>	40	0	100	0	0
<i>Pseudosmittia</i>	0	0	33.3	0	0
<i>Rheocricotopus</i>	0	0	66.6	0	0
<i>Thienemanniella</i>	90	0	100	0	0
N. of taxa	67	39	43	35	21










Figure 4. Similarity (Coefficient of Geographic Resemblance, CGR) in the taxonomic composition of the Chironomidae assemblages from inventories taken in Brazil. r represents the cophenetic correlation coefficient. The abbreviations for the locations inventoried are defined in Table 2. High quality figures are available online.