



## Fluorescent Chromatophore Detected in the Guppy *Poecilia reticulata*

Authors: Nakajima, Masamichi, Shinohara, Eiji, and Fujio, Yoshihisa

Source: Zoological Science, 16(5) : 745-747

Published By: Zoological Society of Japan

URL: <https://doi.org/10.2108/zsj.16.745>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

## [SHORT COMMUNICATION]

## Fluorescent Chromatophore Detected in the Guppy *Poecilia reticulata*

Masamichi Nakajima\*, Eiji Shinohara and Yoshihisa Fujio

Laboratory of Applied Population Genetics, Graduate School of Agricultural Science, Tohoku University,  
Sendai 981-8555, Japan

**ABSTRACT**—Fluorescent chromatophores with bluish white fluorescence were detected in the guppy *Poecilia reticulata*. These chromatophores were observable only by means of a fluorescence microscope, and could not be observed by a standard transmission light microscope or incident light microscope. Resembling other common chromatophores, these fluorescent chromatophores were dendritic cells and similar to other chromatophores in the size, about 50–100  $\mu\text{m}$ .

Of the 17 strains examined, these fluorescent chromatophores were detected only in the R strain, and in all 12 R strain individuals examined, each presented fluorescent chromatophores.

These differences in the occurrence of fluorescent chromatophores among strains suggest that the formation of these chromatophores is the result of genetic polymorphism.

### INTRODUCTION

In mature males of the guppy *Poecilia reticulata*, aquarists have bred a variety of color types and patterns. The modes of inheritance of these body color types and color patterns have been examined (Winge and Ditlevsen, 1948; Ueshima *et al.*, 1998). The body colors of both females and males were determined mainly by the existence of several types of chromatophore, and the genetic polymorphisms of formation of these chromatophores have been described (Ueshima *et al.*, 1998).

Vertebral chromatophores are generally classified into five categories according to Fujii and Oshima (1986). These are the melanophore (black), erythrophore (red), xanthophore (yellow), leucophore (white), and iridophore (reflecting). Recently, the blue chromatophore (cyanophores) was also reported by Goda and Fujii (1995). No chromatophores which have fluorescence have been reported, however, several pigments which have fluorescence have been reported, such as pteridines.

We describe here our preliminary observations regarding fluorescent chromatophores made in the context of our study regarding the inheritance mode of chromatophores.

### MATERIALS AND METHODS

Mature guppies *Poecilia reticulata* which were more than 60 days


old were used as experimental fish. The individuals were collected from 17 strains which were maintained as closed colonies at a density of 300–500 individuals per 60 l aquaria in our laboratory. Each strain was selected by body color type or color pattern, and each trait was fixed in each strain. The examined number of individuals in each strain were more than 6 (3 females and 3 males), as shown in Table 1.

The chromatophores were observed in the scales sampled from specific parts of the body of the guppy (posterior part of head, dorsal fin base, lateral part of trunk and caudal peduncle). These scales were enclosed in the medaka linger solution (Iwamatsu, 1993) for observation. The chromatophore was observed by means of a standard transmission light microscope with differential-interference, and by means of fluorescence microscope (OLYMPUS BX-60). This system used the excitation filter, BP330-385, and the emission filter, BA420.

### RESULTS AND DISCUSSION

Fluorescent chromatophores were detected in the scales sampled in one strain, the R strain, of 17 guppy strains (Table 1). By standard transmission light microscopy, it was not observed (Fig. 1A), however, these bluish white chromatophores were observable by means of fluorescence microscopy (Fig. 1B, 1C). These chromatophores were not visible when standard incidence light was used, however, some granulars were observed by standard transmission light microscopy (Fig. 1A). These chromatophores were composed of dendritic cells and were similar to other chromatophores in size about 50–100  $\mu\text{m}$ . But they differed from previously reported chromatophores, such as melanophores, xanthophores, erythrophores, leucophores, iridophores, and cyanophores (Fujii, 1993; Goda and Fujii, 1995), because of

\* Corresponding author: Tel. +81-22-717-8741;  
FAX. +81-22-717-8743.



**Fig. 1.** Photomicroscopographs showing the fluorescent chromatophore observed in the R strain of the guppy *Poecilia reticulata*. These three photomicroscopographs were taken of the same area under differing optical conditions. **A:** Standard transmission optics with differential-interference microscopy. **B:** Standard transmission optics with differential-interference and fluorescence microscopy. **C:** Fluorescence microscopy.

**Table 1.** The existence of fluorescent chromatophores in each strain of the guppy *Poecilia reticulata*.

Strain	Female	Male
A	—* (n=5)	— (n=6)
B	— (n=3)	— (n=3)
C	— (n=3)	— (n=3)
D	— (n=3)	— (n=3)
D1	— (n=3)	— (n=2)
D2	— (n=3)	— (n=5)
F	— (n=3)	— (n=3)
G	— (n=3)	— (n=3)
G1	— (n=3)	— (n=3)
L	— (n=3)	— (n=3)
O	— (n=3)	— (n=3)
R	+ (n=5)	+ (n=7)
S	— (n=7)	— (n=6)
S3	— (n=3)	— (n=3)
S3HL	— (n=3)	— (n=3)
SC	— (n=3)	— (n=3)
T	— (n=4)	— (n=4)
Y	— (n=4)	— (n=4)

+: Presence —: Absence \*: A few number of fluorescent chromatophores were observed only in the scales picked up from caudal peduncle in one female of the 6 individuals examined.

their fluorescence, the previously described chromatophores having no fluorescence. However, several fluorescent pigments have been reported. Some pteridines, such as biopterin, neopterin, and leucopterin, are known as fluorescent pigments (Matsumoto *et al.*, 1971) and have blue or bluish white fluorescence. In fishes, biopterin has been isolated from goldfish (Hama *et al.*, 1960), *Xiphophorus* fish (Kauffman, 1959) and medaka (Hama *et al.*, 1965). However, the sort of pigments present in the chromatophores detected in this study was not yet identified.

Fluorescent chromatophores were observed only in the R strain of the 17 guppy strains examined (Table 1). The fluorescent chromatophores were not accompanied by chromatophores of other types, because the R strain is an albino strain which has no visible chromatophores. In the A strain, a few fluorescent chromatophores were observed only in scales sampled from the caudal peduncle of one female of the 6 individuals examined. On the other hand, these chromatophores were observed in all the scales of all the individuals examined in the R strain. The genetic control on formation of chromatophores, melanophore, xanthophore and erythrophore, has been observed in the guppy as a body color polymorphism (Ueshima *et al.*, 1998). In regard to the fluorescent chromatophore, the differences of their occurrence among strains suggest that the formation of fluorescent chromatophores in the result of genetic polymorphism.

A cross experiment will be needed for further examination of the genetic control of the fluorescent chromatophore.

## ACKNOWLEDGMENTS

This work was partly supported by a grant-in-aid for scientific research from the Ministry of Education, Science, Culture and Sports in Japan (No. 10660170).

## REFERENCES

- Fujii R, Oshima N (1986) Control of chromatophore movements in teleost fishes. *Zool Sci* 3: 13–47
- Fujii R (1993) Cytophysiology of fish chromatophores. *Int Rev Cytol* 143: 191–255
- Goda M, Fujii R (1995) Blue chromatophores in two species of callionymid fish. *Zool Sci* 12: 811–813
- Hama T, Matsumoto J, Obika M (1960) Localization of pterins in the goldfish-skin melanophores and erythrophores. *Proc Japan Acad* 36: 217–222
- Hama T, Goto T, Tohnoki Y, Hiyama Y (1965) The relation between pterins and chromatophores in the Medaka, *Oryzias latipes*. *Proc Japan Acad* 41: 305–309
- Iwamatsu T (1993) The Biology of the Medaka. Scientist, Tokyo, pp 185–272
- Kauffman T (1959) Untersuchungen über die fluoreszierenden Pigmente in der Haut von Zierfischen. *Z Naturf* 14b: 358–363
- Matsumoto J, Bagnara JT, Taylor JD (1971) Isolation and separation of pteridines from animals. *Exp Physiol Biochem* 4: 289–363
- Ueshima G, Nakajima M, Fujio Y (1998) A study on the inheritance of body color and chromatophores in the guppy *Poecilia reticulata*. *Tohoku J Agr Res* 48: 111–122.

(Received June 10, 1999 / Accepted July 21, 1999)