

Anuran Calcitonins are Diverse in Lower Vertebrates

Authors: Sakamoto, Hidenori, Sasayama, Yuichi, and Ueda, Kunihiko

Source: Zoological Science, 17(1): 97-101

Published By: Zoological Society of Japan

URL: https://doi.org/10.2108/zsj.17.97

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at <u>www.bioone.org/terms-of-use</u>.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Anuran Calcitonins are Diverse in Lower Vertebrates

Hidenori Sakamoto¹, Yuichi Sasayama^{1*} and Kunihiko Ueda²

¹Noto Marine Laboratory, Faculty of Science, Kanazawa University, Uchiura, Ishikawa 927-0553, Japan and ²Department of Biology, Faculty of Science, Kanazawa University, Kakuma, Kanazawa 920-1192, Japan

ABSTRACT—We investigated the nucleotide sequences of cDNA fragments coding calcitonin from ultimobranchial glands in 2 species of urodelans (1 salamander and 1 newt) and 4 species of anurans (1 toad and 3 frogs) by the reverse transcription polymerase chain reaction (RT-PCR) method and rapid amplification of cDNA ends (RACE)-PCR method. The salamander and newt calcitonins were each 97% and 94% similar to the lungfish and caiman calcitonins that we have already reported, in the amino acid sequences. However, anuran calcitonins were not only dissimilar (63–81%) to the lungfish and caiman calcitonins but also diversified (59–91%) even among anurans. The sequence identity of toad calcitonin was always low (59–66%) among anurans.

INTRODUCTION

Calcitonin, a peptide hormone composed of 32 amino acid residues, prevents excessive rises of blood Ca levels (Azria, 1986). As all jawed vertebrates produce calcitonin, this hormone seems to play an important role on Ca homeostasis. The primary structure of calcitonin has so far been determined in some vertebrate groups: mammals, birds, teleosts and stingrays (see Sasayama and Oguro, 1993). Recently, we reported the nucleotide sequences of cDNA fragments coding calcitonin in 4 species of reptiles (Suzuki et al., 1997) and in 4 primitive bony fishes such as lungfish (Suzuki et al., 1999). Reptile calcitonins were extremely similar (88-100% in the amino acid sequences) to both chicken calcitonin and those of the primitive bony fishes. These facts suggest that calcitonin is a very conservative hormone, at least among these animal groups. Similarly, also in amphibians, we clarified the primary structure of bullfrog calcitonin (Yoshida et al., 1997), which was a unique molecule because of a chimera including some amino acid residues common to mammalian and fish calcitonins. However, it is not clear whether or not this characteristic of bullfrog calcitonin is common to all amphibian calcitonins.

In this study, we tried to amplify cDNA fragments coding calcitonin molecule of 2 species of urodelans and 4 species of anurans. Furthermore, we compared the sequences of the amplified nucleotides and of the deduced amino acid sequences to those of the reptiles and fishes (Suzuki *et al.*, 1997; 1999) and among amphibians as well.

* Corresponding author: Tel. +81-768-74-1151; FAX. +81-768-74-1644. E-mail: sasayama@kenroku.kanazawau.ac.jp

MATERIALS AND METHODS

We collected 2 species of urodelans (Tohoku black salamander: *Hynobius nigrescens*; Japanese newt: *Cynops pyrrhogaster*) and 4 species of anurans (Japanese toad: *Bufo japonicus*; Japanese tree frog: *Hyla japonica*; black-spotted pond frog: *Rana nigromaculata*; and forest green tree frog: *Rhacophorus arboreus*) near our laboratory in Uchiura Town, Ishikawa Prefecture. These amphibians were anesthetized by neutralized tricaine methanesulfonate.

Calcitonin-producing ultimobranchial glands were dissected under a binocular microscope and immediately frozen in liquid nitrogen. Total RNA was extracted using an RNA isolation kit (Nippon gene, Toyama), and subjected to a RT-PCR kit (Takara, Kyoto) according to the method of Iwami et al. (1996). Figure 1A shows locations of primers used in the RT-PCR method. As an N-terminal primer (N-1 primer), a mixture of 24 mer was synthesized from the 5'region excluding 1 bp prior to the portion coding calcitonin molecule, on the basis of the consensus of the nucleotide sequences of the human, salmon and chicken calcitonins. On the other hand, three kinds of C-terminal primers were synthesized. The first C-primer (C-1) was composed of 30 mer from the 3'region of the portion coding salmon calcitonin. The second C-primer (C-2) was 30 mer as well but was based on the chicken calcitonin. The third C-primer (C-3) was a mix primer of 26 mer based on the consensus of the portions coding salmon and human calcitonins. The compositions of the reverse-transcription solution, the PCR solution, and the conditions of PCR reaction have been previously described (Suzuki et al., 1997).

Furthermore, 3'-RACE-PCR method was also applied. Oligo(dT) primers were synthesized as Frohman *et al* (1988) suggested. In Fig.1B and C, primers adopted in this method are shown. After the extension with QT primer, in the first PCR, cDNA fragments were amplified by the Q0 primer and N-1 primer used in the RT-PCR method. In the nested PCR, the Q1 primer and each sense primer (gene specific primer, of which nucleotide sequence was obtained in the RT-PCR) of 22 mer in the salamander and 23 mer in the toad were used (Fig. 1B). In the case of newt, we synthesized another N-terminal primer (N-2 primer) for the 3'-RACE-PCR method (Fig. 1C), of which region was at the up-stream of the N-1 primer. The primer was composed of a mixture of 24 mer synthesized on the basis of the





Fig. 1. (A), (B) and (C) Primers' location in the RT-PCR method and 3'-RACE-PCR method. Calcitonin gene is composed of six exons. Exon 4 encodes calcitonin. Figure 1 shows only one part of exon 4. Black portion on the line is a coding region of calcitonin molecule (96 bp). Nucleotide sequences of every primer are also cited. (A) shows the RT-PCR method for all amphibians. (B) and (C) exhibits the 3'-RACE-PCR method for the salamander and the toad, and the newt, respectively.

consensus of the nucleotide sequences of mammal, chicken and salmon calcitonins. In the first PCR for the newt, this N-2 primer and Q0 primer set was used. In the nested PCR, the N-1 primer and Q1 primer set was used.

The conditions for the RACE-PCR are followings. Five μ g of total RNA was reverse-transcribed. The solution was incubated at 55°C for 1 hr. Complementary DNA fragments were subjected to the first PCR. The procedures were done as follows: for 5 min at 97°C, 3 min at 75°C, 3 min at 55°C, 40 min at 72°C, and next 30 cycles of 1 min at

94°C, 1 min at 55°C, 3 min at 72°C, and finally 15 min at 72°C. Furthermore, nested PCR was performed for 2 min at 95°C, 30 cycles of 45 sec at 95°C, 30 sec at 55°C, 1 min at 72°C, and 30 min at 72°C.

The purified cDNA fragments were ligated into pT7 Blue vector (Takara, Kyoto). After screening, nucleotide sequence was determined by a DNA sequencer (Model 373S, Applied Biosystems) using the dideoxynucleotide chain terminal procedure with T7 primer for the vector. At least 4 clones were analyzed in each species.

98

RESULTS

In the Tohoku black salamander and the Japanese toad, neither the C-1 nor the C-2 primer could amplify the cDNA fragments coding calcitonin molecule. Only the C-3 primer was compatible. Therefore, cDNA fragments of 132 bp lacking the information of the last 15 of the 96 bp were amplified. However, those lacking regions were covered by the 3'-RACE-PCR method. In the Japanese newt, only the 3'-RACE-PCR method with the N-2 primer was successful to amplify the full length of the cDNA fragments coding calcitonin molecule. In the 3 species of frogs (Japanese tree frogs, black-spotted pond frogs and forest green frogs), cDNA fragments covering calcitonin molecules were amplified by the RT-PCR method.

Figure 2 shows the nucleotide sequences and the

deduced amino acid sequences of amphibian calcitonins. Bullfrog calcitonin and some other calcitonins are also cited for reference.

In these amphibian calcitonins, the amino acid residues located in the 15th and 19th positions, which are common to those of mammalian calcitonins in bullfrog calcitonin, were not always common even among the amphibians examined. On the other hand, the nucleotide sequence of the salamander calcitonin was 86% similar to that of the lungfish, and 85% to the caiman, 76% to those of the goldfish and sole (Table 1). The sequence identities of the amino acid residues deduced from the nucleotides in these animals were higher than those of the nucleotides, as those values range from 84% to 97% (Table 2). The amino acid sequence of the newt calcitonin coincided completely with that of the salamander. In contrast,

	1	1/	0		20			30)			40			50			6	0		7	0			80			90	,	
LUNGFISH 5	'-TGTTC	CAATI	TGAC	TACI	TGC	GTG	TTG	GGC	AAA	TTG	TCG	CAI	GAA	TTG	CAC	ААА	TTG	CAI	AAT	TAC	сст	CGC	АСТ	GAC	GTC	GGA	GCT	GGA	ACT	сст - 3 '
GOLDFISH	c	GC (-c			c	G		c	c	0	3 G	c			c		-CA	T	A		c	A-T	A		G	T	G	G
SOLE	CA-	GGC (-c	7	r	c			c-c	c		T	A-T			c		CA	T	c		c	A	G		G	G	A	G
TOHOKU BLACK SALAMANDER	c	cc		·	:		c			c-c			3			G			c-							G	c			G
JAPANESE NEWT	c	-gcc	3	·c		r	c		G	c-c		0	3 G	c		G	c	· (3-c-				c			c	c			G
JAPANESE TOAD	CG-	-GCG-C	3			- A-A	cc-	G	-G-	c-c	- AT	0	- 3т	T	A		- 7.1	ACC	TC-					T		G	c	c	G	C
JAPANESE TREE FROG	C					AC-	ст-	т-G		c-c		0					C-2	A-0	170-	T			G		G	c	c	C	G	
BLACK-SPOTTED POND FROG		- GGGCC		.c c			»	C		c													0			0				
FOREST CREEN TREE FROG						C	NC-	1 G		c				» – m			C-7		mc-					N	0					
CATWAN				C e		C	AC-			c			91	X -1			C-2	LAG.					0	A			C			0
CAIMAN	G-					A	C			C												1				G				
								10																				20		
TIBIORTON					6		•	10	•	•		a 1		•		.	• • • •	20	• • • •		D	•	m1	•	**- 1	61	• • •	30	m 1	D
LONGFISH	Cys se	r Asn i	Leu Se	er thr	Cys	vai	Leu	GTĀ	гда	Leu	ser	GTL	i Giu	Leu	HIS	гда	Leu	GTL	Asn	TYT	Pro	Arg	Thr	Asp	vai	σтΫ	ALA	стў	Thr	Pro
GOLDFISH		Ser			-	-	-	-	-	-	-	-	-	-	-	-	-	-	Thr	-	-	-	-	Asn	-	-	-	-	-	-
SOLE	- Th	r Gly			-	7	-	-	-	-	-	-	Asp	Ile	-	-	-	-	Thr	-	-	-	-	Asn	-	-	-	-	-	-
TOHOKU BLACK SALAMANDER		· -			-	-	-	-	-	-	-	-	-	-	-	-	-	-	Thr	-	-	-	-	-	-	-	-	-	-	-
JAPANESE NEWT		· -			-	-	-	-	-	-	-	-	-	-	-	-	-	-	Thr	-	-	-	-	-	-	-	-	-	-	-
JAPANESE TOAD	- Al	.a Arg			-	Ile	Pro	-	Arg	-	Tyr	-	Asp	Phe	Asn	-	Tyr	Thr	Ser	-	-	-	-	-	-	-	-	-	-	-
JAPANESE TREE FROG		Gly			-	Thr	-	Trp	-	-	-	-	Asp	-	-	-	-	Asn	Ser	- '	-	-	-	-	-	-	-	-	-	-
BULLFROG		Gly		· -	-	Ala	-	Met	-	-	-	-	Asp	-	-	Arg	Phe	Asr	Ser	-	-	-	-	Asn	-	-	-	-	-	-
BLACK-SPOTTED POND FROG		· Gly			-	Leu	Thr	Trp	-	-	-	-	Asp	-	Gln	-	-	Asn	Ser	-	-	-	-	-	-	-	-	-	-	-
FOREST GREEN TREE FROG	- Va	ıl -		· -	-	Ala	Thr	Met	-	-	Ile	-	Asp	Ile	-	-	-	Ser	Ser	-	-	-	-	Asn	-	-	-	-	-	-
CAIMAN	- Al	.a Ser			-	_	-	-	-	-	-	-	-	-	-	-	-	-	Thr	-	-	-	-	-	-	-	-	-	_	-

Fig. 2. Nucleotide sequences and amino acid sequences of amphibian calcitonins. Horizontal bars indicate identical bases and amino acid residues to those of the lungfish calcitonin. Data other than amphibians are also cited for reference (Suzuki *et al.* 1997; Yoshida *et al.* 1997; Suzuki *et al.* 1999).

Table 1. Sequence identity of nucleotides of cDNA coding calcitonin molecules shown in Fig. 1.

LUNG	FISH													
91	GOL	DFIS	Н											
81	88	88 SOLE												
97	94	94 84 TOHOKU BLACK SALAMANDER												
97	94	84	4 100 JAPANESE NEWT											
63	59	63	63 63 JAPANESE TOAD											
81	78	78	81	81 81 66 JAPANESE TREE FROG										
72	75	75	72	72 59 84 BULLFROG										
75	72	72	75	75	66	91	78 BLACK-SPOTTED POND FROG							
69	69	75	69	69	63	72	75	72	FOF	REST GREEN TREE FROG				
91	94	84	94	94	66	78	69	72	66	CAIMAN				

H. Sakamoto et al.

2011														
72	GOLDFISH													
72	83 SOLE													
86	76 76 TOHOKU BALCK SALAMANDER													
78	78	76	76 91 JAPANESE NEWT											
66	65	64	72 67 JAPANESE TOAD											
71	74	74	77	77 77 77 JAPANESE TREE FROG										
68	72	71	73	72	73	82	BLACK-SPOTTED POND FROG							
61	68	67	67	69	69	74	80 FOREST GREEN TREE FROG							
89	72	75	85	78	73	71	69	63	CAIMAN					

 Table 2.
 Sequence identity of amino acid residues of calcitonin molecules shown in Fig. 1.

the nucleotide sequences of the 4 species of anurans were 67–77% similar to those of the salamander and newt, 64–74% to those of the goldfish and sole, 63–73% to that of the caiman, 61–71% to that of the lungfish (Table 1). The sequence identities of the deduced amino acid residues in those animals ranged from 59% to 81% (Table 2). Among anurans as well, the sequence identities of the deduced amino acid residues ranged widely from 59% to 91%. The calcitonin of the Japanese toad was clearly distinctive, since the sequence identities of the amino acid residues were only 59-66%. However, the calcitonin of the Japanese tree frog was relatively similar to those of other frogs. The sequence identities of the amino acid residues were 66 - 91%, of which the highest value was obtained with the black-spotted pond frog.

DISCUSSION

The present study did not support the hypothesis that the chimera-like characteristic of bullfrog calcitonin is general among amphibian calcitonins. Therefore, its characteristic is limited so far to the bullfrog calcitonin alone. On the other hand, the salamander and newt calcitonins were very similar to those of the lungfish and the caiman. This fact suggests that urodele calcitonin may have originated from calcitonin of the ancestor common to the primitive bony fishes and that the calcitonin has been transported to reptiles as well. In anurans, however, the sequence identities of the nucleotides coding calcitonin were rather low in comparison to those of other lower vertebrates. Especially, calcitonin of the Japanese toad was distinctive even among amphibians. Recently, Conlon et al. (1998) reported the diversity of the amino acid sequences of pancreatic polypeptide (PP) among amphibians. This peptide is composed of 36 amino acid residues whose molecular size is similar to that of calcitonin. The amino acid sequence of the PP of Rana sylvatica was the same as that of the bullfrog PP. The PP from the green frog (Rana ridibunda) was different only in one amino acid residue from that of the bullfrog. However, the PP of the cane toad (Bufo marinus) was different in 6 amino acid residues from that of the bullfrog PP. As paleontological evidence shows the early diversity of toads from other frogs (Duellman and Trueb, 1985), comparative aspects of calcitonin and PP may contribute our understanding of the phylogenetic location of toads in anurans. In the frog phylogeny, Hylidae (Japanese tree frog) is closer to Bufonidae (Japanese toad) than to Ranidae (black-spotted pond frog) (Duellman and Trueb, 1985). However, the sequence identity of the tree frog calcitonin to the pond frog calcitonin was higher in the amino acid residues than to the toad calcitonin. Therefore, the relationship based on the sequence identity of frog calcitonins did not coincide with the phylogenetic relationship of the frogs.

As above-mentioned, anuran calcitonins were distinctive and diversified from those of lower vertebrates. Amphibians are the first vertebrates that advanced on land. In general, it has been accepted that new environments relax various restrictions to suppress gene changes (Kimura, 1983). Recently, Arai et al. (1998) reported that the alpha subunit of the glycoprotein hormone in the pituitary diversifies more largely in amphibians than in mammals, birds and lungfish. Therefore, results from calcitonin and the study of the subunit of glycoprotein seem to coincide with the theory established by Kimura (1983). Salamanders and newts changed their morphology and physiology moderately from their ancestor on the way of evolution. Anurans, on the other hand, altered drastically. On the path to adaptation, frogs acquired a characteristic physiology known as metamorphosis. In the metamorphic processes, frogs form their leg bones in a manner different from that of other vertebrates. At first, tadpoles develop endolymphatic sacs in their vertebral bones and accumulate Ca in those sacs preceding metamorphosis, since they can not obtain Ca through food on account of the reconstruction of their digestive tracts. Ca deposited in the sacs is utilized to make four leg bones during metamorphosis. Calcitonin accelerates the accumulation of Ca into the sacs (Sasayama and Oguro, 1985). Such function of calcitonin has not been known in other vertebrates. Furthermore, Stiffler et al. (1998) recently reported that, in frogs, calcitonin stimulates Ca influx through the skin into the body. In other vertebrates, such a role of calcitonin has not been known, because this hormone accelerates primarily to decrease blood Ca levels. On the other hand, frogs changed their moving style from crawling to jumping, which deformed their skeletal structure to a large extent. Some of their vertebral bones became united to fix the waist for jumping, and some parts of the skull regressed to lighten the weight. These various characteristics of phenomena related with Ca homeostasis in anurans might have been brought about by the relaxation on the suppression of changes in the calcitonin molecule, or they might have caused the relaxation on the molecule.

ACKNOWLEDGMENTS

We thank Dr. Nobuo Suzuki (Kanazawa University) for his kind suggestions of 3'-RACE-PCR method.

REFERENCES

- Arai Y, Kubokawa K, Ishii S (1998) Cloning of cDNAs for the pituitary glycoprotein hormone alpha subunit precursor molecules in three amphibian species. Gen Comp Endocrinol 112: 46–53
- Azria M (1989) "The Calcitonins." Karger, Basel
- Conlon JM, Platz JE, Chartrel N, Vaudry H, Nielsen PF (1998) Amino acid sequence diversity of pancreatic polypeptide among the amphibia. Gen Comp Endocrinol 112: 146–152
- Duellman WE, Trueb L (1985) "Biology of Amphibians." McGraw-Hill Book Company, New York

- Frohman MA, Dush MK, Martin GR (1988) Rapid pruduction of full length cDNAsfrom rare transcripts: Amplification using a single gene-specific oligonucleotide primer. Proc Nat Acad Sci U.S.A. 85: 8998–9002
- Iwami M, Tanaka A. Hano N, Sakurai S (1996) Bombyxin gene expression in tissues other than brain detected by reverse transcription-polymerase chain reaction (RT-PCR) and in situ hybridization. Experientia 52: 882–887
- Kimura M (1983) "The Neutral Theory of Molecular Evolution." Cambridge University Press, Cambridge
- Sasayama Y, Oguro C (1985) The role of the ultimobranchial glands on Ca balance in bullfrog tadpoles. In "Current Trends in Comparative Endocrinology", Hong Kong University Press, Hong Kong, pp. 837–838.
- Sasayama Y, Oguro C (1993) Ultimobranchial Gland. In "Atlas of Endocrine Organs." Springer-Verlag, Tokyo, pp. 83–92.
- Stiffler DF, Yee JC, Tefft JD (1998) Responses of frog skin, *Rana pipiens*, calcium ion transport to parathyroid hormone, calcitonin, and vitamin D3. Gen Comp Endocrinol 112: 191–199
- Suzuki N, Eguchi C, Hirai T, Sasayama Y (1997) Nucleotide sequences of reptile calcitonins: their high homology to chicken calcitonin. Zool Sci 14: 833–836
- Suzuki N, Ueda K, Sakamoto H, Sasayama Y (1999) Fish calcitonin genes: primitive bony fish genes have been conserved in some lower vertebrates. Gen Comp Endocrinol 113: 369–373
- Yoshida A, Kaiya H, Takei Y, Watanabe TX, Nakajima K, Suzuki N, Sasayama Y (1997) Primary structure and bioactivity of bullfrog calcitonin. Gen Comp Endocrinol 107: 147–152

(Received January 22, 1999 / Accepted August 25, 1999)