



## **Involvement of Calcium, Inositol-1,4,5 Trisphosphate and Diacylglycerol in the Prothoracicotropic Hormone-Stimulated Ecdysteroid Synthesis and Secretion in the Prothoracic Glands of *Bombyx mori***

Authors: Dedos, Skarlatos G., and Fugo, Hajime

Source: Zoological Science, 18(9) : 1245-1251

Published By: Zoological Society of Japan

URL: <https://doi.org/10.2108/zsj.18.1245>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

# Involvement of Calcium, Inositol-1,4,5 Trisphosphate and Diacylglycerol in the Prothoracicotrophic Hormone-Stimulated Ecdysteroid Synthesis and Secretion in the Prothoracic Glands of *Bombyx mori*

Skarlatos G. Dedos<sup>1</sup> and Hajime Fugo<sup>2\*</sup>

<sup>1</sup>*Institute of Biology, National Center for Scientific Research "Demokritos",  
Aghia Paraskevi, Athens 15310, Greece*

<sup>2</sup>*Department of Biological Production, Tokyo University of Agriculture  
and Technology, Fuchu-shi, Tokyo 183, Japan*

**ABSTRACT**—The objective of this study was to determine which intracellular second messenger systems are activated by prothoracicotrophic hormone in the prothoracic glands (PGs) of *Bombyx mori*. Recombinant prothoracicotrophic hormone (rPTTH) could stimulate ecdysteroid synthesis and secretion from day 6 PGs of the 5th instar of *Bombyx mori* within 30 min of *in vitro* incubation. However, rPTTH did not stimulate any increases in the glandular content of inositol 1,4,5-trisphosphate and cAMP during this short incubation period. Extracellular  $\text{Ca}^{2+}$  influenced the basal and rPTTH-stimulated ecdysteroid synthesis and release in a dose-dependent manner. The L-type  $\text{Ca}^{2+}$  channel antagonist, nitrendipine, inhibited the rPTTH-stimulated ecdysteroid synthesis and secretion ( $\text{IC}_{50} \sim 28 \mu\text{M}$ ). The phospholipase C inhibitor, 2-nitro-4-carboxyphenyl-N, N-diphenylcarbamate, inhibited the rPTTH-stimulated ecdysteroid synthesis ( $\text{IC}_{50} \sim 19 \mu\text{M}$ ). The protein kinase C inhibitor, chelerythrine chloride, inhibited the rPTTH-stimulated ecdysteroid synthesis ( $\text{IC}_{50} \sim 14 \mu\text{M}$ ). The protein kinase C activator, phorbol-12-myristate 13-acetate (PMA), could stimulate basal ecdysteroid synthesis and secretion ( $\text{EC}_{50} \sim 1 \mu\text{M}$ ) and its inactive  $\alpha$ -isomer (4  $\alpha$ -PMA) was ineffective. The combined results suggest that the PTTH-stimulated ecdysteroid synthesis and release in the PGs of *Bombyx* is dependent on extracellular  $\text{Ca}^{2+}$  and the bifurcating second messenger signalling cascade of inositol 1,4,5-trisphosphate and diacylglycerol.

**Key words:** prothoracicotrophic hormone, ecdysteroid, inositol-1,4,5 trisphosphate, calcium, prothoracic gland, *Bombyx mori*

## INTRODUCTION

The synthesis and secretion of ecdysteroid hormone from prothoracic glands (PGs) of insects is regulated by a family of cerebral neuropeptides, the prothoracicotrophic hormones (PTTHs) (Gilbert *et al.*, 1996). In the tobacco hornworm, *Manduca sexta*, two size variants of PTTH have been identified: big PTTH and small PTTH (Gilbert *et al.*, 1996). Both forms of *Manduca* PTTH were shown to mediate their signalling cascade via cAMP as a second messenger (Smith, 1993; Watson *et al.*, 1993). The signalling cascade of *Manduca* big PTTH does not involve any increases in glandular inositol phosphate production (Girgenrath and Smith, 1996). Furthermore, phorbol

esters (protein kinase C activators) were not found to stimulate basal and big PTTH-stimulated ecdysone synthesis from *Manduca* PGs, although protein kinase C activity was found in the PGs of this insect (Smith, 1993).

In the silkworm, *Bombyx mori*, only the 30,000-dalton PTTH was found to stimulate ecdysteroid synthesis and secretion from the PGs of this insect (Kiriishi *et al.*, 1992). In the present study, we used the recombinant form of *Bombyx* PTTH (rPTTH; Ishibashi *et al.*, 1994), which was shown to stimulate ecdysteroid synthesis and release from 5th instar PGs (Dedos *et al.*, 1999a). Our objective in this study was to determine which intracellular second messenger systems are activated by *Bombyx* PTTH in the PGs and how such activation is regulated. We present evidences to suggest that activation of ecdysone synthesis and secretion by PTTH is dependent on the presence of extracellular  $\text{Ca}^{2+}$  and on the generation and action of the second messengers, inositol-1,4,5

\* Corresponding author: Tel. +81-42-367-5734;  
FAX. +81-42-367-5734.  
E-mail: fugoh@cc.tuat.ac.jp

trisphosphate (IP<sub>3</sub>) and diacylglycerol which are generated from the bifurcating signalling system of receptor activated phospholipase C.

## MATERIALS AND METHODS

### Animals

All experiments used larvae from the hybrid J106xDAIZO. The larvae were reared on mulberry leaves under a 12:12-L:D photoperiod at 25±1°C and 60% relative humidity. Larvae were staged as we previously described (Dedos *et al.*, 1999a). In this particular hybrid, the 5th instar period lasts about ~208 hr. The onset of wandering behaviour occurs ~144 hr (day 6) after the final larval ecdysis. Female larvae of day 6 were exclusively used in this study.

### Reagents

Recombinant *Bombyx* PTTH (rPTTH) (Ishibashi *et al.*, 1994) was dissolved in either Grace's medium (GIBCO-BRL, Grand Island, NY, USA) or Ca<sup>2+</sup>-free Ringer saline and stored at -20°C until use. Nitrendipine (Calbiochem, La Jolla, CA, USA), chelerythrine chloride (Research Biochemicals International, Natick, MA, USA), 2-nitro-4-carboxyphenyl-*N,N*-diphenylcarbamate (NCDC) (Sigma, St. Louis, MO, USA), phorbol 12-myristate 13-acetate (PMA) and 4 $\alpha$ -phorbol 12-myristate 13-acetate (4 $\alpha$ -PMA) (GIBCO-BRL, Grand Island, NY, USA) were prepared as stock solutions in dimethylsulfoxide (DMSO). Ethylene glycol bis( $\beta$ -aminoethyl ether)-*N,N,N',N'*-tetraacetic acid (EGTA) was purchased from Wako (Osaka, Japan).

### In vitro prothoracic gland assay

Larvae were anaesthetized by submersion in water and PGs were dissected rapidly (~2 min /animal) from each larva in sterile saline (0.85% NaCl). The glands were pre-incubated in Grace's medium for 15–30 min. A paired gland design was used in some experiments. One gland of the pair was incubated in 20  $\mu$ l of medium containing one or more experimental agent(s); the other gland of the pair was incubated in 20  $\mu$ l medium containing solvent. When DMSO was used as solvent, all PGs were incubated in medium containing 1% DMSO. The presence of 1% DMSO does not affect basal or rPTTH-stimulated ecdysteroid synthesis and release. In experiments where the dose response to an experimental agent was investigated, glands from a pooled group were randomly selected for incubation at the indicated doses. In experiments requiring a Ca<sup>2+</sup>-free medium, Ringer's saline (Shirai *et al.*, 1994) was prepared using NaCl (4.5 mM) in place of the standard 4.5 mM CaCl<sub>2</sub>. Hepes buffer (0.01 M, pH 6.8) and either EGTA (0.1 mM) or the indicated amount of Ca<sup>2+</sup> (in the form of CaCl<sub>2</sub>) were added prior to use (Hayes *et al.*, 1995). Incubations were carried out at 25±1°C in high humidity in 96-multiwell plates (Wako, Osaka, Japan). After each designated incubation period, the medium was removed, and an aliquot of the medium was subjected to radioimmunoassay for quantification of ecdysteroid content.

### Quantification of inositol 1,4,5-trisphosphate and cAMP

The content of inositol 1,4,5-trisphosphate (IP<sub>3</sub>) in PGs was quantified by a [<sup>3</sup>H]radioreceptor assay using the kit and protocol available from New England Nuclear Corp (Boston, MA, USA) as we previously described (Dedos *et al.*, 1998). Recovery of IP<sub>3</sub> from the glands was determined to be 90%. Sensitivity of the IP<sub>3</sub> assay system was approximately 0.1 pmol.

The content of cAMP in PGs was quantified by enzyme immunoassay using the kit and protocol available from Cayman Chemical Co. (Ann Arbor, MI, USA), as we previously described (Dedos *et al.*, 1999a).

### Radioimmunoassay

The amount of ecdysteroid in the incubation medium was quantified by radioimmunoassay, as we previously described (Dedos *et*

*al.*, 1999a). Radiolabeled ecdysone, [23,24-<sup>3</sup>H]ecdysone (sp. act. 53 Ci/mmol) was purchased from New England Nuclear Corp. (Boston, MA, USA).

### Statistical analyses

Statistical significance of the results was determined by analysis of variance or Student's t-test. For most experiments, analysis of variance was followed by Tukey multiple comparisons tests. Test results are shown in figure legends. The statistical analyses were done with computer software (GraphPad Prism™ 2.0).

## RESULTS

### Stimulation of ecdysteroid synthesis and secretion by rPTTH and effects of rPTTH on glandular inositol 1,4,5-trisphosphate (IP<sub>3</sub>) and cAMP levels

To determine rPTTH potentiation of ecdysteroid synthesis and release from day 6 PGs, groups of glands were exposed to 1 ng of rPTTH for 30 min. The glands were then transferred to plain Grace's medium and incubated for an additional 90 min (Table 1). The results showed that rPTTH

**Table 1.** Stimulation of ecdysteroid synthesis and release from PGs of *Bombyx mori* by rPTTH


Treatments	Ecdysteroid (ng/gland)	
	30 min incubation	90 min incubation
Basal	1.83±0.16	2.62±0.28
1 ng rPTTH	3.77±0.24	4.57±0.14

Individual glands from day 6 of the 5th instar were incubated in Grace's medium in the presence of 1 ng rPTTH and their contralaterals in the same medium alone (basal) for 30 min. Then glands from each group were transferred and incubated for an additional 90 min period to Grace's medium in the absence of rPTTH. Each group is the mean±SEM of 6 glands. Tests for the difference between control (basal) and experimental (rPTTH) glands (t-tests) revealed that rPTTH stimulated ecdysteroid synthesis and release in both incubation periods. Results of Tukey multiple comparisons for ecdysteroid synthesis and release (pairs of means enclosed by the range of a bracket are not significantly different, P>0.05): [3.77–4.57].


could stimulate ecdysteroid synthesis and secretion from day 6 PGs within the 30 min and even in the subsequent 90 min incubation period (Table 1).

Next we determined whether rPTTH transduces its signalling cascade through activation of a phospholipase C (PLC), thus generating the second messenger inositol 1,4,5-trisphosphate (IP<sub>3</sub>). Groups of day 6 PGs were incubated in the presence of 1 ng rPTTH/gland and their IP<sub>3</sub> contents were determined at various time intervals. As shown in Fig. 1, rPTTH could not stimulate any increase in glandular IP<sub>3</sub> content in time intervals ranging from 30 sec to 2 hr (P=0.95).

In a previous study, we showed that a cerebral prothoracicotrophic factor, which is different from *Bombyx* PTTH, stimulates cAMP accumulation in the PGs of this insect (Dedos *et al.*, 1999a). However, both a cerebral prothoracicotrophic factor and rPTTH did not stimulate cAMP accumulation in day 6 PGs within a 30 min incubation (Dedos *et al.*, 1999a). Since rPTTH stimulated ecdysteroid synthesis and secretion within



**Fig. 1.** Effects of rPTTH on prothoracic gland inositol trisphosphate levels. Each point is the mean  $\pm$  S.E. of 4–5 glands. Data were analysed by one-way analysis of variance ( $P=0.95$ ).




**Fig. 2.** Effects of rPTTH on prothoracic gland cAMP levels. Each point is the mean  $\pm$  S.E. of 6 glands. Data were analyzed by one-way analysis of variance ( $P=0.51$ ).

a 30 min incubation (Table 1), we determined whether rPTTH transduces its signalling cascade through activation of an adenylate cyclase at time periods much shorter than 30 min. The results (Fig. 2) showed that 1 ng rPTTH did not change glandular cAMP content during incubation times ranging from 2 min to 1 hr ( $P=0.51$ ; Fig. 2). The glandular content of another cyclic nucleotide, cGMP, was either not affected by rPTTH in day 6 PGs (data not shown).

#### Dependence of basal and rPTTH-stimulated ecdysteroid synthesis and secretion on extracellular $\text{Ca}^{2+}$

Prothoracic glands from day 6 were incubated for 2 hr in Ringer's saline, in which the indicated concentration of  $\text{CaCl}_2$  was added with or without 1 ng rPTTH/gland (Fig. 3). Both, basal and rPTTH-stimulated ecdysteroid secretions were influenced by extracellular  $\text{Ca}^{2+}$  in a dose-dependent manner. When glands were incubated in  $\text{Ca}^{2+}$ -free Ringer's saline, the rPTTH-stimulated ecdysteroid synthesis and secretion reached levels similar to the basal ecdysteroid secretion in the absence




**Fig. 3.** Effect of external  $\text{Ca}^{2+}$  on basal and rPTTH-stimulated ecdysteroid synthesis and release by day 6 PGs. Each point is the mean  $\pm$  S.E. of 5 to 6 glands. Results of Tukey multiple comparisons (pairs of means enclosed by the range of a bracket are not significantly different,  $P>0.05$ ): Basal; [0.56–0.97], [3.01–3.14], [3.14–3.99]; rPTTH; [0.77–1.73], [8.55–8.63], [10.49–11.47].

of extracellular  $\text{Ca}^{2+}$ . The higher the concentration of  $\text{CaCl}_2$  in saline, the more rPTTH could stimulate ecdysteroid synthesis and release above basal levels (Fig. 3).

#### Effects of nitrendipine on basal and rPTTH-stimulated ecdysteroid synthesis and secretion

Prothoracic glands of day 6 were incubated for 2 hr in the presence of various concentrations of the L-type  $\text{Ca}^{2+}$  channel antagonist, nitrendipine, and 1 ng rPTTH/gland. The results



**Fig. 4.** Effect of nitrendipine on basal and rPTTH-stimulated ecdysteroid synthesis and release by day 6 PGs. Individual glands were incubated for 2 hr with the indicated concentrations of nitrendipine alone (basal) or in the simultaneous presence of 1 ng rPTTH/gland (rPTTH). Each point is the mean  $\pm$  S.E. of 5 to 6 glands. Data for the effects of nitrendipine on basal ecdysteroid synthesis and release were analyzed by one-way analysis of variance ( $P=0.11$ ). Results of Tukey multiple comparisons for the effects of nitrendipine on rPTTH-stimulated ecdysteroid synthesis and release (pairs of means enclosed by the range of a bracket are not significantly different,  $P>0.05$ ): [1.28–2.33], [9.74–11.43].

**Table 2.** Effect of rPTTH on ecdysteroid synthesis and release and IP<sub>3</sub> levels of day 6 larval PGs of *Bombyx mori*

Treatment	Ecdysteroid (ng/gland)		IP <sub>3</sub> (pmol/gland)	
	Basal	Experimental	Basal	Experimental
1 ng rPTTH	2.94±0.23	8.66±0.72	0.31±0.03	0.29±0.02
1 ng rPTTH+50 µM NCDC	2.78±0.22	2.68±0.45	0.27±0.03	0.25±0.01

Glands were incubated for 2 hr in the presence of 1 ng rPTTH/gland or 1 ng rPTTH+50 µM NCDC. After the end of the incubation the IP<sub>3</sub> content of the PGs and the amount of secreted ecdysteroid in the medium were determined. Each value is the mean±SEM of 6 glands. Tests for the difference between experimental and control glands (t-tests) revealed that NCDC inhibited the rPTTH-stimulated ecdysteroid synthesis and release ( $P<0.05$ ). Data of IP<sub>3</sub> determinations were analyzed by one-way analysis of variance ( $P=0.48$ ).

in Figure 4 indicate that nitrendipine competitively inhibited the rPTTH-stimulated ecdysteroid synthesis and release *in vitro* ( $IC_{50}\sim 28\text{ }\mu\text{M}$ ). Nitrendipine did not inhibit the basal secretory activity of the glands ( $P=0.11$ ; Fig. 4). The rPTTH-stimulated ecdysteroid synthesis and release was inhibited in a similar way by the phenylalkylamine derivative, verapamil but not by the benzothiazepine, diltiazem (data not shown).

#### Dependence of rPTTH-stimulated ecdysteroid synthesis and release on the generation of inositol 1,4,5-trisphosphate (IP<sub>3</sub>) in the PGs


Two-nitro-4-carboxyphenyl-*N,N*-diphenylcarbamate (NCDC) is known as a potent inhibitor of PLC (Takei *et al.*, 1991). Glands of day 6 were incubated for 2 hr in the presence of 1 ng rPTTH/gland or 1 ng rPTTH+50 µM NCDC (Table 2). Recombinant PTTH stimulated ecdysteroid synthesis and release but NCDC inhibited the rPTTH-stimulated ecdysteroid

synthesis and release ( $P>0.05$ ; Table 2). The IP<sub>3</sub> content of the PGs remained unchanged for 2 hr (Table 2;  $P=0.48$ ).


In a different approach, PGs of day 6 were incubated for 2 hr in the presence of various concentrations of NCDC and 1 ng rPTTH/gland. The results in Figure 5 indicate that NCDC competitively inhibited the rPTTH-stimulated ecdysteroid synthesis and release ( $IC_{50}\sim 19\text{ }\mu\text{M}$ ), although NCDC did not inhibit the basal secretory activity ( $P=0.11$ ; Fig. 5).

#### Effects of chelerythrine chloride on basal and rPTTH-stimulated ecdysteroid synthesis and release

Prothoracic glands of day 6 were incubated for 2 hr in the presence of various concentrations of the protein kinase C (PKC) inhibitor chelerythrine chloride (Herbert *et al.*, 1990) and 1 ng rPTTH/gland. The results in Fig. 6 indicate that chelerythrine chloride competitively inhibited the steroidogenic effect of rPTTH ( $IC_{50}\sim 14\text{ }\mu\text{M}$ ). Chelerythrine chloride did not



**Fig. 5.** Effect of NCDC on basal and rPTTH-stimulated ecdysteroid synthesis and release by day 6 PGs. Individual glands were incubated for 2 hr with the indicated concentrations of NCDC alone (basal) or in the simultaneous presence of 1 ng rPTTH/gland (rPTTH). Each point is the mean±S.E. of 5 to 6 glands. Data for the effects of NCDC on basal ecdysteroid level were analyzed by one-way analysis of variance ( $P=0.11$ ). Results of Tukey multiple comparisons for the effects of NCDC on rPTTH-stimulated ecdysteroid synthesis and release (pairs of means enclosed by the range of a bracket are not significantly different,  $P>0.05$ ): [0.74–1.95], [6.62–8.77], [8.84–9.15].




**Fig. 6.** Effect of chelerythrine chloride on basal and rPTTH-stimulated ecdysteroid synthesis and release by day 6 PGs. Individual glands were incubated for 2 hr with the indicated concentrations of chelerythrine chloride alone (basal) or in the simultaneous presence of 1 ng rPTTH/gland (rPTTH). Each point is the mean±S.E. of 5 to 6 glands. Data for the effects of chelerythrine chloride on basal ecdysteroid level were analyzed by one-way analysis of variance ( $P=0.41$ ). Results of Tukey multiple comparisons for the effects of chelerythrine chloride on rPTTH-stimulated ecdysteroid synthesis and release (pairs of means enclosed by the range of a bracket are not significantly different,  $P>0.05$ ): [4.81–7.76], [10.49–11.33].

inhibit the basal secretory activity of the glands ( $P=0.41$ ; Fig. 6). Similar results were obtained with another PKC inhibitor, calphostin C (data not shown).

### Effect of phorbol 12-myristate 13-acetate (PMA) and $4\alpha$ -phorbol 12-myristate 13-acetate ( $4\alpha$ -PMA) on basal ecdysteroid synthesis and release

In initial experiments, day 6 PGs were incubated in the presence of varying concentrations (0.1 nM ~ 0.1 mM) of PMA and  $4\alpha$ -PMA for 2 hr. During this incubation period, however, no statistically significant change in ecdysteroid synthesis and release was observed with PMA or  $4\alpha$ -PMA (data not shown). Statistical analysis showed that PMA did not alter basal ecdysteroid synthesis and release ( $P=0.14$ ) and similar results were also observed with  $4\alpha$ -PMA ( $P=0.11$ ; data not shown). Since the possibility remained that the 2 hr incubation period was too short to observe any PMA-mediated effects on basal ecdysteroid synthesis and release, the incubation period was extended to 5 hr. At 5 hr incubation period, PMA stimulated basal ecdysteroid synthesis and release ( $EC_{50}\sim 1\ \mu\text{M}$ ; Fig. 7). Under similar conditions,  $4\alpha$ -PMA did



**Fig. 7.** Effect of phorbol 12-myristate 13-acetate (PMA) and  $4\alpha$ -PMA on basal ecdysteroid synthesis and release by day 6 PGs. Individual glands were incubated for 5 hr with the indicated concentrations of PMA or  $4\alpha$ -PMA. Each point is the mean  $\pm$  S.E. of 5 to 6 glands. Data for the effects of  $4\alpha$ -PMA on basal ecdysteroid level were analyzed by one-way analysis of variance ( $P=0.21$ ). Results of Tukey multiple comparisons for the effects of PMA on basal ecdysteroid synthesis and release (pairs of means enclosed by the range of a bracket are not significantly different,  $P>0.05$ ): [7.31–9.35], [9.05–11.03], [14.50–15.70].

not alter basal ecdysteroid synthesis and release ( $P=0.21$ ; Fig. 7). These results were verified using a paired gland design at 5 hr incubation. Glands incubated in 1  $\mu\text{M}$  PMA in Grace's medium secreted 12.3 ng ecdysone/gland while their contralaterals secreted 7.6 ng/gland ( $n=4$ ), but such an effect was not observed during a 2 hr incubation (data not shown).

## DISCUSSION

The results in the present study demonstrate that the ste-

roidogenic effect of *Bombyx* PTTH in PGs is dependent on extracellular  $\text{Ca}^{2+}$  and the bifurcating second messenger system that generates the two intracellular messengers, inositol 1,4,5-trisphosphate and diacylglycerol.

During a 30 min incubation of day 6 PGs with recombinant PTTH there was a substantial increase of ecdysteroid synthesis and release ( $P<0.05$ ; Table 1). This stimulation by rPTTH was not mediated through the generation of the second messenger  $\text{IP}_3$  since rPTTH did not increase glandular  $\text{IP}_3$  content even after 2 hr incubation (Fig. 1, Table 2). Moreover, the increase in ecdysteroid during 30 min incubation (Table 1) was not preceded or accompanied by any rPTTH-mediated increase in the glandular content of the second messenger cAMP, since rPTTH did not increase cAMP content even after 1 hr incubation (Fig. 2). Gu *et al.* (1996), using as source of PTTH medium in which brain complexes of *Bombyx* were incubated, found that PTTH could stimulate cAMP accumulation in the PGs of this insect. However, we observed that this cAMP stimulating activity in *Bombyx* PGs was mediated by another cerebral prothoracicotrophic factor different from PTTH (Dedos *et al.*, 1999a). Recombinant PTTH stimulated cAMP accumulation only in day 4 and day 5 *Bombyx* PGs, and did not increase the cAMP content after a 30 min incubation of day 6 PGs (Dedos *et al.*, 1999a). In this study, it was further shown that rPTTH did not stimulate cAMP accumulation in PGs even at incubation periods much shorter than the 30 min used in previous experiments (Fig. 2; Dedos *et al.*, 1999a). Previous research using day 6 *Bombyx* PGs showed that  $\text{Ca}^{2+}$  and cAMP signaling pathways can cooperatively, as well as independently, stimulate ecdysteroid synthesis and release from the PGs (Dedos and Fugo, 1999b). All these results suggested that there exists a high degree of complexity and stage-specific variability in the signaling cascades that mediate ecdysteroid synthesis and release by the PGs of *Bombyx*. Therefore, in order to explain the signaling cascade of rPTTH-mediated ecdysteroid synthesis and release, it was important to choose, for the present experiments, a developmental stage (day 6, 5th instar) at which the rPTTH-stimulated ecdysteroid synthesis and release does not involve multiple second messenger cascades.

Both basal and rPTTH-stimulated ecdysteroid secretions were found to be affected by the presence of extracellular  $\text{Ca}^{2+}$  in a dose-dependent manner (Fig. 3). A similar stage-specific dependence of basal ecdysteroid synthesis and release on the presence of extracellular  $\text{Ca}^{2+}$  was reported for the PGs of *Manduca* (Meller *et al.*, 1990). The rPTTH-stimulated ecdysteroid synthesis was reduced to levels similar to basal release in the absence of  $\text{Ca}^{2+}$  from the incubation medium as shown in Fig. 3 and also by Gu *et al.* (1998).

These results suggest that a  $\text{Ca}^{2+}$ -influx pathway into the cytosol of the PG cells may be the primary stimuli in the *Bombyx* PTTH signal transduction cascade. There are several pathways of  $\text{Ca}^{2+}$  influx activation by receptor agonists through the plasma membrane (Fasolato *et al.*, 1994). Among them, the receptor-operated channels together with the voltage-operated channels provide brief and high intensity bursts

of  $\text{Ca}^{2+}$  influx into the cytosol, while the store-operated channels provide a much smaller but sustained influx of  $\text{Ca}^{2+}$  (Berridge, 1997). The existence of receptor-operated  $\text{Ca}^{2+}$  channels was suggested for the PGs of *Manduca* (Girgenrath and Smith, 1996). These authors suggested that the *Manduca* big PTTH receptor mediates  $\text{Ca}^{2+}$ -influx through an L-type  $\text{Ca}^{2+}$  channel based on the ability of the L-type  $\text{Ca}^{2+}$  channel antagonist, nitrendipine, to inhibit *Manduca* big PTTH-stimulated ecdysteroid synthesis and release (Girgenrath and Smith, 1996). This *Manduca* big PTTH-mediated  $\text{Ca}^{2+}$  channel was described to be nitrendipine-sensitive and verapamil-insensitive (Girgenrath and Smith, 1996), while *Manduca* small PTTH-stimulated ecdysteroid synthesis was inhibited by verapamil (Hayes *et al.*, 1995). In this study, we showed that nitrendipine competitively inhibited only the rPTTH-stimulated ecdysteroid secretion, without affecting basal ecdysteroid synthesis and release in *Bombyx* (Fig. 4). We also suggest that binding of *Bombyx* PTTH to its cellular membrane receptor results in the opening of an L-type  $\text{Ca}^{2+}$  channel which promotes  $\text{Ca}^{2+}$  entry into the cytosol of PG cells.

Research on the intracellular  $\text{Ca}^{2+}$  modulation in the PGs of *Galleria mellonella* and *Manduca* (Birkenbeil, 1996; 1998) revealed that there are differences between insect species in the PTTH-mediated mobilization of intracellular  $\text{Ca}^{2+}$  in the PGs. For example, it was shown that the PTTH-mediated increase in intracellular  $\text{Ca}^{2+}$  in the PG cells of *Galleria* was abolished by the removal of extracellular  $\text{Ca}^{2+}$  or in the presence of the  $\text{Ca}^{2+}$  channel antagonists nicardipine and verapamil (Birkenbeil, 1996). Similar experiments with the PGs of *Manduca* showed that the PTTH-mediated increase in intracellular  $\text{Ca}^{2+}$  was not abolished by removal of extracellular  $\text{Ca}^{2+}$  or in the presence of nicardipine and verapamil (Birkenbeil, 1998). Our results suggest that the PTTH-mediated mobilization of intracellular  $\text{Ca}^{2+}$  in the PGs of *Bombyx* is different from those of *Galleria* and *Manduca*.

Although rPTTH did not directly stimulate the generation of  $\text{IP}_3$  (Fig. 1), its ecdysteroidogenic action was dependent on the generation of this second messenger since the rPTTH-stimulated ecdysteroid secretion was competitively inhibited by NCDC (Table 2 and Fig. 5). These results suggest that the  $\text{Ca}^{2+}$  that is mobilized by PTTH into the PG cells is sequestered in  $\text{IP}_3$ -sensitive intracellular  $\text{Ca}^{2+}$  stores and released in the ecdysteroidogenic process by the action of  $\text{IP}_3$ . Thus, we believe that the PTTH signalling cascade in the PG cells involves keeping intracellular  $\text{Ca}^{2+}$  levels elevated during maintained stimulation and speeding the replenishment of intracellular  $\text{Ca}^{2+}$  stores. By doing so, filled internal  $\text{Ca}^{2+}$  stores are maintained, and their  $\text{Ca}^{2+}$  content is ready to be mobilized by  $\text{IP}_3$ . Because the rPTTH-stimulated ecdysteroid synthesis and release was dependent on external  $\text{Ca}^{2+}$  (Fig. 3), it appears that an initial PTTH-mediated influx of  $\text{Ca}^{2+}$  requires the generation of  $\text{IP}_3$  (Table 2), and this explains why the NCDC-mediated inhibition of PLC eliminated the rPTTH-stimulated ecdysteroid synthesis and release (Fig. 5). The NCDC-mediated inhibition of PLC may eliminate increases in glandular  $\text{IP}_3$  content. This, in turn, would eliminate the  $\text{IP}_3$ -medi-

ated  $\text{Ca}^{2+}$  release from intracellular stores which would be the first event in the PTTH-mediated cascade of events that lead to replenishing of intracellular  $\text{Ca}^{2+}$  stores. Since a similar function in the replenishing of intracellular  $\text{Ca}^{2+}$  stores has been proposed for store-operated  $\text{Ca}^{2+}$  channels (Friel, 1996), we believe that if a receptor-operated  $\text{Ca}^{2+}$  channel is regulated by the *Bombyx* PTTH receptor, then this  $\text{Ca}^{2+}$  channel is probably in close proximity to  $\text{IP}_3$  receptors and it is also regulated by the state of filling of the  $\text{IP}_3$ -sensitive intracellular  $\text{Ca}^{2+}$  stores.

Models for conformational coupling mechanisms between  $\text{Ca}^{2+}$  channels in the plasma membrane and cytosolic  $\text{Ca}^{2+}$  stores have been suggested to explain receptor-mediated (Rink, 1990; Tsunoda, 1993) and capacitative  $\text{Ca}^{2+}$  entries (Berridge, 1995; 1997; Parekh, 1997). Capacitative  $\text{Ca}^{2+}$  entry is the process in which  $\text{Ca}^{2+}$  enters the cytosol and replenishes intracellular  $\text{Ca}^{2+}$  stores which are emptied through the action of  $\text{IP}_3$  (Friel, 1996). The action of  $\text{IP}_3$  is essential for capacitative  $\text{Ca}^{2+}$  entry through store-operated  $\text{Ca}^{2+}$  channels (Friel, 1996) in the same way as the action of  $\text{IP}_3$  is essential for the PTTH-stimulated ecdysteroid synthesis and release. Moreover, since basal ecdysteroid synthesis and release decreased in the absence of external  $\text{Ca}^{2+}$  (Fig. 3), it is quite possible that store-operated  $\text{Ca}^{2+}$  channels exist in the PG cells to facilitate agonist-insensitive  $\text{Ca}^{2+}$  entry and replenishment of  $\text{Ca}^{2+}$  stores.

Chelerythrine chloride competitively inhibited the rPTTH-stimulated ecdysteroid synthesis and release (Fig. 6) whereas PMA stimulated basal ecdysteroid synthesis, but only after a 5 hr incubation (Fig. 7). One can assume that the PKC limb of the phosphoinositide signalling pathway is directly involved in the PTTH-evoked  $\text{Ca}^{2+}$  influx and increase of ecdysteroid synthesis and release. Protein kinase C has disparate and cell type specific effects on store-operated  $\text{Ca}^{2+}$  influx and voltage-dependent  $\text{Ca}^{2+}$  channel activity (Shearman *et al.*, 1989). Activation of store-operated  $\text{Ca}^{2+}$  influx by PKC was documented (Tsunoda, 1993; Parekh and Penner, 1997) and this enzyme was found to positively regulate voltage-dependent  $\text{Ca}^{2+}$  channel activity (Shearman *et al.*, 1989). Thus, if a PKC-mediated positive feedback mechanism operates in the PGs, a large rPTTH-stimulated ecdysteroid synthesis and release would not have been expected in the PGs in the presence of PKC inhibitors (Fig. 6). Alternatively, the PKC limb of the phosphoinositide signalling pathway may regulate ecdysteroid synthesis and secretion through an independent pathway. It may also converge with the PTTH-mediated signalling cascade at later stages of the ecdysteroidogenic process for maximal activation of ecdysteroid synthesis and release. Another possibility is that activation of PKC is required for a prolonged and sustained ecdysteroid synthesis and release as suggested for the *Manduca* PGs (Smith, 1993). Such a possibility may explain why a 5 hr incubation was required for PMA to stimulate ecdysteroid synthesis and release by the PGs of *Bombyx* (Fig. 7).

## ACKNOWLEDGEMENTS

We would like to thank Dr. Satoshi Takeda, National Institute of Sericultural and Entomological Science, Tsukuba, Japan, for donating the antiserum for 20-hydroxyecdysone. This study was supported by grants from the Ministry of Education, Science, Sports and Culture of Japan (No. 08276206, 09265204 and 10161205).

## REFERENCES

- Berridge M J (1995) Capacitative calcium entry. *Biochem J* 312: 1–11
- Berridge M J (1997) Elementary and global aspects of calcium signalling. *J Physiol* 499: 291–306
- Birkenbeil H (1996) Involvement of calcium in prothoracicotropic stimulation of ecdysone synthesis in *Galleria mellonella*. *Arch Insect Biochem Physiol* 33: 39–52
- Birkenbeil H (1998) Intracellular calcium in prothoracic glands of *Manduca sexta*. *J Insect Physiol* 44: 279–286
- Dedos S G, Fugo H, Kataoka H (1998) A new cerebral factor stimulates IP<sub>3</sub> levels in the prothoracic glands of *Bombyx mori*. *Insect Biochem Mol Biol* 28: 767–774
- Dedos S G, Fugo H, Nagata S, Takamiya M, Kataoka H (1999a) Differences between recombinant PTTH and crude brain extracts in cAMP-mediated ecdysteroid secretion from the prothoracic glands of the silkworm, *Bombyx mori*. *J Insect Physiol* 45: 415–422
- Dedos S G, Fugo H (1999b) Interactions between Ca<sup>2+</sup> and cAMP in ecdysteroid secretion from the prothoracic glands of *Bombyx mori*. *Mol Cell Endocrinol* 154: 63–70
- Fasolato C, Innocenti B, Pozzan T (1994) Receptor-activated Ca<sup>2+</sup> influx: how many mechanisms for how many channels? *Trends Pharmacol Sci* 15: 77–83
- Friel D D (1996) TRP: Its role in phototransduction and store-operated Ca<sup>2+</sup> entry. *Cell* 85: 617–619
- Gilbert L I, Rybczynski R, Tobe S S (1996) Endocrine cascade in insect metamorphosis. In *Metamorphosis: Postembryonic Reprogramming of Gene Expression in Amphibian and Insect Cells*. Gilbert LI, Tata JR, Atkinson GB (eds): San Diego: Academic Press pp 59–107
- Girgenrath S, Smith W A (1996) Investigation of presumptive mobilization pathways for calcium in the steroidogenic action of big prothoracicotropic hormone. *Insect Biochem Mol Biol* 26: 455–463
- Gu S H, Chow Y S, Lin F J, Wu J L, Ho R J (1996) A deficiency in prothoracicotropic hormone transduction pathway during the early last larval instar of *Bombyx mori*. *Mol Cell Endocrinol* 127:109–116
- Gu S H, Chow Y S, O'Reilly D R (1998) Role of calcium in the stimulation of ecdysteroidogenesis by recombinant prothoracicotropic hormone in the prothoracic glands of the silkworm, *Bombyx mori*. *Insect Biochem Mol Biol* 28: 861–867
- Hayes G C, Muehleisen D P, Bollenbacher W E, Watson R D (1995) Stimulation of ecdysteroidogenesis by small prothoracicotropic hormone: role of calcium. *Mol Cell Endocrinol* 115: 105–112
- Herbert J M, Augereau J M, Gleye J, Maffrand J P (1990) Chelerythrine is a potent and specific inhibitor of protein kinase C. *Biochem Biophys Res Comm* 172: 993–999
- Ishibashi J, Kataoka H, Isogai A, Kawakami A, Saegusa H, Yagi Y, Mizoguchi A, Ishizaki H, Suzuki A (1994) Assignment of disulfide bond location on prothoracicotropic hormone of the silkworm, *Bombyx mori*: A homodimeric peptide. *Biochemistry* 33: 5912–5919
- Kiriishi S, Nagasawa H, Kataoka H, Suzuki A, Sakurai S (1992) Comparison of the *in vivo* and *in vitro* effects of bombyxin and prothoracicotropic hormone on prothoracic glands of the silkworm, *Bombyx mori*. *Zool Sci* 9: 149–155
- Meller V, Sakurai S, Gilbert L I (1990) Developmental regulation of calmodulin-dependent adenylate cyclase activity in an insect endocrine gland. *Cell Regul* 1: 771–780
- Parekh A B, Penner R (1997) Store depletion and calcium influx. *Physiol Rev* 77: 901–930
- Rink T J (1990) Receptor-mediated calcium entry. *FEBS Letters* 268: 381–385
- Shearman M S, Sekiguchi K, Nishizuka Y (1989) Modulation of ion channel activity: A key function of the protein kinase C enzyme family. *Pharmacol Rev* 41: 211–237
- Shirai Y, Iwasaki T, Matsubara F, Aizono Y (1994) The carbachol-induced release of prothoracicotropic hormone from brain-corpus cardiacum-corpus allatum complex of the silkworm, *Bombyx mori*. *J Insect Physiol* 40: 469–473
- Smith W A (1993) Second messengers and the action of prothoracicotropic hormone in *Manduca sexta*. *Am Zool* 33: 330–339
- Takei M, Ueno M, Endo K, Nakagawa H (1991) Effect of NCDC, a protease inhibitor, on histamine release from rat peritoneal mast cells induced by Anti-IgE. *Biochem Biophys Res Comm* 181: 1313–1322
- Tsunoda Y (1993) Receptor-operated Ca<sup>2+</sup> signaling and crosstalk in stimulus secretion coupling. *Biochim Biophys Acta* 1154: 105–156
- Watson R D, Yeh W E, Muehleisen D P, Watson C J, Bollenbacher W E (1993) Stimulation of ecdysteroidogenesis by small prothoracicotropic hormone: Role of cAMP. *Mol Cell Endocrinol* 92: 221–228

(Received May 7, 2001 / Accepted September 7, 2001)