



## **Characterization of the Sperm Receptor for Acrosome Reaction-Inducing Substance of the Starfish, *Asterias Amurensis***

Authors: Kawamura, Mayu, Matsumoto, Midori, and Hoshi, Motonori

Source: Zoological Science, 19(4) : 435-442

Published By: Zoological Society of Japan

URL: <https://doi.org/10.2108/zsj.19.435>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

# Characterization of the Sperm Receptor for Acrosome Reaction-Inducing Substance of the Starfish, *Asterias amurensis*

Mayu Kawamura<sup>1</sup>, Midori Matsumoto<sup>2</sup> and Motonori Hoshi<sup>2\*</sup>

<sup>1</sup>Department of Bioscience, Graduate School of Bioscience and Biotechnology, Tokyo Institute of Technology, 4259 Nagatsuta, Yokohama 226-8501, and

<sup>2</sup>Center for Life Science and Technology, Graduate School of Science and Technology, Keio University, Hiyoshi 3-14-1, Yokohama, 223-8522, Japan

**ABSTRACT**—Acrosome reaction-inducing substance (ARIS) in the jelly coat of starfish eggs is a highly sulfated proteoglycan-like molecule of an apparent molecular size over 10<sup>4</sup>kDa and plays a pivotal role in the induction of acrosome reaction in homologous spermatozoa. It is known in *Asterias amurensis* that ARIS binds to a restricted area of the anterior portion of sperm head, and that a glycan fragment of ARIS, named Fragment 1, consisting of 10 repeats or so of a pentasaccharide unit retains the biological activity of ARIS to an appreciable extent. In this report, we have shown the binding of Fragment 1, a relatively small pure glycan fragment of ARIS, to the putative ARIS receptor on the sperm surface by three independent methods. First, the specific binding of P-ARIS to isolated sperm membranes was monitored in real-time by using a surface plasmon resonance detector, namely a Biacore sensor system. The specific and quantitative binding of Fragment 1 to the intact sperm and to isolated sperm membranes was similarly monitored. Secondly, the binding of <sup>125</sup>I-labeled Fragment 1 to the intact sperm was stoichiometrically measured, for which we had developed a unique procedure for radioiodination of saccharide chains. It is found that Fragment 1 competes with P-ARIS for the binding to ARIS-receptor, suggesting that Fragment 1 is a useful ligand in the search for ARIS receptor protein(s). Thirdly, the putative receptor molecules were specifically labeled by using Fragment 1 as a ligand for photoaffinity crosslink technique. Taking these results into account, we conclude that starfish sperm have the ARIS receptor, which consists most probably of 50 to 60 kDa proteins, of reasonably high affinity (for Fragment 1, K<sub>d</sub>=15 μM, B<sub>max</sub>=8.4×10<sup>4</sup> per cell).

**Key words:** acrosome reaction, ARIS, receptor, photoaffinity crosslink, starfish sperm

## INTRODUCTION

The acrosome reaction of sperm is an essential event for fertilization in most animals (Dan, 1967). It involves the exocytosis of the acrosomal vesicle to expose the devices for penetration through the egg coats and for subsequent fusion with the egg plasma membrane (Longo, 1997). Although it is well established that the egg coat contains signal molecules for the induction of acrosome reaction, our knowledge on the chemical structure of such molecules is much limited except for some echinoderms (for reviews see Miller and Ax, 1990; Litscher and Wassarman, 1993; Mengerink and Vacquier, 2001). In sea urchins, egg jelly

polysaccharides such as the fucose sulfate polymer (FSP) (Vacquier and Moy, 1997; Alves *et al.*, 1997, 1998; Vilela-Silva *et al.*, 1999) plays a key role for the induction of acrosome reaction though some other jelly components are involved in the induction (Yamaguchi *et al.*, 1987; Hirohashi and Vacquier, 2002). A 210-kDa protein named REJ (receptor for egg jelly) is thought to be the receptor for FSP (Moy *et al.*, 1996; Vacquier and Moy, 1997). Although the study of REJ has provided us a new insight into the human polycystic kidney disease (Mengerink *et al.*, 2000), stoichiometric character of REJ as the FSP receptor remains unknown.

In the starfish, *Asterias amurensis*, the actions of three egg jelly components in concert are responsible for the induction of acrosome reaction (Hoshi *et al.*, 1994, 1999). Namely, a highly sulfated proteoglycan-like molecule named acrosome reaction-inducing substance (ARIS) (Uno and Hoshi, 1978; Ikadai and Hoshi, 1981a, b), sulfated steroid

\* Corresponding author: Tel. +81-45-566-1773;  
FAX. +81-45-566-1448.  
E-mail: mhoshi@chem.keio.ac.jp

saponins named Co-ARIS (Nishiyama *et al.*, 1987a, b), and sperm-activating peptides named asterosap (Nishigaki *et al.*, 1996). ARIS alone but no others can induce the acrosome reaction in high calcium or high pH sea water, whereas either Co-ARIS or asterosap is required besides ARIS for the induction in normal sea water (Ikadai and Hoshi, 1981a, b; Matsui *et al.*, 1986). ARIS has an apparent molecular size over  $10^4$ kDa and retains its activity even after extensive digestion with Pronase. The Pronase digest of ARIS (P-ARIS) still has an apparent molecular size of  $10^4$ kDa or so (Ikadai and Hoshi, 1981a).

Although it is reported from our laboratory that a small number of high affinity receptor for ARIS locates in a restricted area of the anterior portion of sperm heads (Ushiyama *et al.*, 1993; Longo *et al.*, 1995), the receptor remains to be isolated and characterized. An extremely large molecular size of ARIS even after Pronase digestion interfered with our trials to characterize and identify the receptor. Poor information on the chemical structure of ARIS also limited our search for the receptor. Thus we tried to disintegrate ARIS into a much smaller fragment with the biological activity. We have recently found that sugar chain fragments liberated from ARIS or P-ARIS by ultra-sonication retains the biological activity of ARIS to an appreciable extent. The fragments are composed of 10 repeats or so of the following pentasaccharide unit and collectively named Fragment 1 (Koyota *et al.*, 1997): [4- $\beta$ -D-Xylp-1-3- $\alpha$ -D-Galp-1-3- $\alpha$ -L-Fucp-4(SO<sub>3</sub><sup>-</sup>)-1-3- $\alpha$ -L-Fucp-4(SO<sub>3</sub><sup>-</sup>)-1-4- $\alpha$ -L-Fucp-1-]. Because we know the structure of Fragment 1 precisely and its size is not too large to handle, Fragment 1 seems a proper ligand for characterization and identification of the ARIS receptor even though its activity was much lower than that of ARIS or P-ARIS.

In this paper, first we show the specific binding of Fragment 1 in real time to the intact sperm and isolated sperm membranes by using a Biacore sensor system, a surface plasmon resonance detector. Then we quantitatively measure the binding of <sup>125</sup>I-Fragment 1 to the intact sperm to obtain kinetic parameters of the receptor. Finally, we use photoaffinity crosslinkers to detect the receptor molecules.

## MATERIALS AND METHODS

### Animals and Gametes

The starfish, *Asterias amurensis*, were collected in the breeding season from Tokyo Bay and Otsuchi Bay on the Pacific coast of Honshu, Japan, and from the coast of Tasmania, Australia. The Tasmanian population is known to be the offspring of animals that were accidentally introduced from Tokyo Bay recently. Spermatozoa of the two populations did not distinguish the egg jelly of domestic animals from that of foreign ones.

Sperm was collected as "dry sperm" by cutting the sperm ducts of the testes. When the acrosome reaction was scored 60% or less with egg jelly, the sperm was not used for experiments. Mature eggs were collected by treating the ovaries with 10  $\mu$ M 1-methyladenine (Sigma Chemicals, St. Louis, MO) and washed with artificial sea water (ASW) consisting of 430mM NaCl, 9mM CaCl<sub>2</sub>, 9mM KCl, 23mM MgCl<sub>2</sub>, 25mM MgSO<sub>4</sub>, 10mM EPPS (N-2-hydrox-

yethylpiperazine-N'-3-propanesulfonic acid; Dojindo Lab., Kumamoto, Japan) buffer, pH 8.2, in MQ water (water purified with a Milli-Q system from Millipore Corp., Bedford, MA). Throughout the present study, MQ water was used unless otherwise specified.

### Preparation of Egg Jelly, P-ARIS and Fragment 1

Egg jelly solution was prepared according to Ikadai and Hoshi (1981a). The egg suspension was gradually acidified to pH 5.5 with 0.1N HCl to dissolve the jelly coat, then returned to pH 8.2 with 0.1N NaOH, and centrifuged at 2,000 g for 5min to remove the eggs. The supernatant was centrifuged at 10,000 g for 30min to remove cell debris and the clear solution of egg jelly was stored at -20°C until use. P-ARIS was prepared according to Matsui *et al.* (1986) with slight modifications; briefly, ethanol precipitation of the egg jelly solution, digestion of the precipitate with Pronase (Sigma Chemicals, St. Louis, MO), gel-filtration of the digest on Sepharose CL-4B (Amersham Pharmacia Biotech AB, Uppsala, Sweden), ion-exchange chromatography of the digest on DEAE Toyopearl 650M (Tosoh Corp., Tokyo), dialysis against MQ water, and lyophilization. Fragment 1 was prepared from purified P-ARIS according to Koyota *et al.* (1997) with slight modifications; briefly dissolution of 30mg P-ARIS in 10ml of MQ water, sonication for 30min by using a Branson Sonifier Cell Disruptor 2000 (Branson Ultrasonics Corp., Danbury, CT), ion-exchange chromatography on DEAE Toyopearl 650M with a linear gradient elution program starting from water to 1.0M NaCl, dialysis of the fraction eluted about 0.9M NaCl against water, and lyophilization.

### Preparation of Biotinyl Fragment 1

Fragment 1 was biotinylated according to Shinohara *et al.* (1995) with slight modifications. A 100 $\mu$ l-aliquot of 10mg (ca. 1 $\mu$ mol)/ml Fragment 1 was mixed with the same volume of 10mM EZ-link biotin-LC-hydrazide (Pierce Chemical Co., Rockford, IL) in 30% acetonitrile, and the mixture was incubated at 90°C for 2hr. The mixture was directly used for the Biacore system as described later.

### <sup>125</sup>I-Labeling of P-ARIS and Fragment 1

<sup>125</sup>I-P-ARIS was prepared according to a modification using iodo-beads (Pierce Chemical Co., Rockford, IL) of the chloramine-T method (Hunter and Greenwood, 1962) as described previously (Ushiyama *et al.*, 1993). Prior to radiiodination, Fragment 1 was coupled with 4-methoxyphenethylamine. Fragment 1 (1.1mg; ca. 0.11 $\mu$ mol) was dissolved in 40 $\mu$ l of 3.1M 4-methoxyphenethylamine (Sigma-Aldrich Japan Corp., Tokyo) in acetic acid and incubated at 90°C for 1hr. Then, 70 $\mu$ l of 22.6M borane dimethylamine complex (Sigma-Aldrich Japan Corp., Tokyo) in 61.5% acetic acid was added and the mixture was stirred at 80°C for 35min. 4-Methoxyphenethylamine-conjugated Fragment 1 was purified with a Bio-Gel P-4 column (1.0 $\times$ 12.7cm; Bio-Rad Labs., Hercules, CA) and lyophilized. 4-Methoxyphenethylamine-conjugated Fragment 1 (1.2mg) was radioiodinated at C3 and C5 of the benzene ring by the iodo-beads/chloramine-T method as described above, and the product was purified with a Bio-Gel P-4 column (1.0 $\times$ 10cm). <sup>125</sup>I-Fragment 1 (11 $\mu$ g as fucose, 1.44 $\times$ 10<sup>7</sup>cpm) thus prepared was stored below -20°C.

### Conjugation of Fragment 1 with Sulfo-SBED

For photoaffinity crosslink of ARIS receptor, we developed a procedure to conjugate the partially oxidized Fragment 1 with an amino-reactive and biotin-containing photoaffinity crosslinker, sulfo-succinimidyl-2-[6-(biotinimido)-2-(p-azidobenzamido)-hexanoamido]ethyl-1,3'-dithiopropionate (Sulfo-SBED; Pierce Chemical Co., Rockford, IL). The reducing terminal of Fragment 1 was first reduced to prevent the "over-oxidation" of sugars from the reducing terminal during modification. Fragment 1 (10mg, ca. 1.0 $\mu$ mol) was dissolved in 200 $\mu$ l of 50mM sodium borate buffer, pH 8.3 containing

5mg (0.13mmol) sodium borohydride. After incubation at room temperature for 4hr, pH was adjusted to 4.0 with glacial acetic acid to destroy excess borohydride. The reaction mixture was left at room temperature for 14hr. The reduced Fragment 1 was purified by chromatography on a Bio-Gel P-4 column (1.0×10cm) equilibrated and eluted with water, and lyophilized. It was then dissolved in 230μl of 40mM imidazole buffer, pH 6.5 and partially oxidized with 2.5mg (12μmol) sodium periodate for 2hr at 0°C. The reaction mixture was directly applied to a Bio-Gel P-4 column (1.0×10cm) equilibrated and eluted with water, and the product was lyophilized. The partially oxidized Fragment 1 was dissolved in 100μl water and 240μl of 2M 1,6-hexanediamine (Wako Pure Chemicals, Osaka) in water was added to convert most if not all aldehyde residues into corresponding Schiff base structures. After incubation for 15min at room temperature, 20mg (0.32mmol) of sodium cyanoborohydride was added to convert the Schiff bases into more stable secondary amines. After incubation for 4hr at room temperature, 3.0mg (79μmol) of sodium borohydride was added, and incubated another 4 hr at room temperature to complete the reduction. The product, namely modified Fragment 1 having primary amine residues, was purified with a Bio-Gel P-4 column (1.0×10cm) as described, and lyophilized.

This product (5.0mg, ca. 0.5μmol) was dissolved in 250μl of 0.1M phosphate-buffered saline (PBS), pH 7.2 and mixed with 20μl of 50mM Sulfo-SBED in dimethylsulfoxide. After incubation for 1hr at room temperature, the mixture was applied to a Bio-Gel P-4 column (1.0×10cm) equilibrated and eluted with water, and the sugar-positive fractions were pooled and lyophilized. Then, the lyophilizate was dissolved in 500μl of 0.1M PBS, pH 7.2 and applied slowly to a monomeric avidin affinity column (2ml; Pierce Chemical Co., Rockford, IL). The column was thoroughly washed with 0.1M PBS, pH 7.2 to remove contaminants. The biotin-labeled, thus most probably the Sulfo-SBED-conjugated, product was eluted out of the column with 2mM D-biotin in 0.1M PBS, pH 7.2. The eluted fractions were pooled, concentrated into 400μl by a centrifuge evaporator, and applied to a Bio-Gel P-4 column (1.0×10cm) equilibrated with water. Then the product was eluted with water to be free of contaminants such as biotin and Sulfo-SBED, and lyophilized. It was estimated for sugar content by resorcinol-sulfuric acid method (Monsigny *et al.*, 1988) and for conjugated Sulfo-SBED moiety by estimation of biotin with 4'-hydroxyazobenzene-2-carboxylic acid (HABA) according to Green (1965).

#### Preparation of Sperm Membrane Fraction


Dry sperm was diluted 5-fold with a cavitation buffer containing 40mM KCl, 1mM MgCl<sub>2</sub>, 10mM piperazine-*N,N'*-bis (2-ethanesulfonic acid) (PIPES; Dojindo Lab., Kumamoto, Japan), pH 6.5. The sperm suspension was subjected to nitrogen cavitation in a Parr nitrogen cavitation bomb (Parr Instrument Co., Moline, IL) for 30min at 1000psi with stirring on ice. The resultant suspension was fractionated by differential centrifugation at 10,000 g for 30min and 100,000 g for 1hr. The microsome fraction was re-suspended in a storage buffer containing 40mM KCl, 1mM MgCl<sub>2</sub>, 20mM MgSO<sub>4</sub> and 10mM PIPES, pH 6.7, and stored at -80°C.

#### Bioassay of Acrosome Reaction


The acrosome reaction was assayed according to Matsui *et al.* (1986) with slight modifications. Dry sperm was diluted 100-fold in ASW containing a test sample like egg jelly, and the mixture was incubated for 5min at room temperature. The spermatozoa were fixed with glutaraldehyde at the final concentration of 0.83% and stained with erythrosine. More than 100 spermatozoa were scored in each experiment under a Nomarski microscope for the extrusion of acrosomal process. For the activity assessment of ARIS, P-ARIS and Fragment 1, high Ca<sup>2+</sup> ASW (430mM NaCl, 50mM CaCl<sub>2</sub>, 9mM KCl, 23mM MgCl<sub>2</sub>, 25mM MgSO<sub>4</sub>, 10mM EPPS buffer, pH 8.2) was used instead of ASW.

#### Binding Assay with Biacore System

The basic principle of the binding assay using a Biacore X (Biacore Inc., Uppsala, Sweden) has been documented previously (Johnsson *et al.*, 1991). P-ARIS was immobilized on the sensor chip surface (C1, research grade) by amino-coupling; briefly, acti-



**Fig. 1.** Acrosome reaction-inducing activity of Fragment 1 in high calcium sea water. The capacity of Fragment 1 (●) to induce the acrosome reaction in 50mM Ca<sup>2+</sup>-sea water was estimated. P-ARIS (□) was used as the positive control. Concentrations of Fragment 1 and P-ARIS are expressed as L-fucose.



**Fig. 2.** Real-time assay of the binding of isolated sperm membrane to immobilized P-ARIS by Biacore system. About 0.2ng of P-ARIS was immobilized on the sensor chip surface. At the flow rate of 20μl/min, 40μl of sperm membrane suspension of the protein concentration as expressed on the corresponding sensogram was introduced to the chip surface. The bottom sensogram represents the competition of the binding by the addition of 0.4 mg P-ARIS to 40μl of the membrane suspension. Values of the ordinate, expressed in resonance units (RU), represent the increments in mass concentration of the sensor surface. The increment of 1000 RU corresponds to the binding of ca. 1ng membrane proteins. The sensograms show relative responses in RU after background subtraction versus time in seconds.

vation of carboxyl groups on the chip surface with *N*-hydroxysuccinimide (Wako Pure Chemicals, Osaka) and *N*-ethyl-*N'*-(3-dimethylaminopropyl) carbodiimide hydrochloride (Tokyo Kasei Kogyo Co., Tokyo), coupling of 0.1ml of 1mg/ml P-ARIS in 10mM sodium acetate buffer pH 4.0, blocking of excess carboxyl groups by 1.0M ethanolamine (Sigma Chemicals, St. Louis, MO). In this way, about 0.2ng of P-ARIS was immobilized. For immobilization of Fragment 1, streptavidin was immobilized on the sensor chip surface (C1, research grade) by amino-coupling. When biotinyl Fragment 1 was introduced to the streptavidin tip, about 70pg of it was immobilized. For the mock-coupled sensor chips, the immobilization procedure was followed except for the addition of the ligand, namely P-ARIS or biotinyl Fragment 1.

Samples were injected over the sensor chip surface at a flow rate of 20 $\mu$ l/min at 20°C. The buffer used for sample dilution was 10mM sodium acetate buffer, pH 4.0, and the running buffer was *N*-2-hydroxyethylpiperazine-*N'*-2-ethanesulfonic acid (HEPES; Dojindo Lab., Kumamoto, Japan) buffered saline (HBS) containing 0.15M NaCl, 3.4mM EDTA (Dojindo Lab., Kumamoto, Japan), 0.005 % Tween 20 and 10mM HEPES, pH 7.4. After every run, the sensor tip was washed with 20 $\mu$ l of 0.05% Nonidet P-40 (Sigma Chemicals, St. Louis, MO). To make sure that the binding is a specific one, it was monitored also in the presence of a large excess of ligand in the running buffer. In each experiment, the sample was simultaneously injected over the mock-coupled sensor chip surface as a blank run. All data were corrected for the background by subtracting the blank run, using BIA evaluation 3.0 software.

#### Binding Assay of $^{125}$ I-P-ARIS and $^{125}$ I-Fragment 1

A 20 $\mu$ l aliquot of sperm suspension in ASW (1.4 $\times 10^7$  cells/ $\mu$ l) was mixed with 30  $\mu$ l of ASW containing  $^{125}$ I-P-ARIS (0.22  $\mu$ g as fucose, 2.5 $\times 10^4$  cpm) or  $^{125}$ I-Fragment 1 (0.33 $\mu$ g as fucose, 4.8 $\times 10^3$ cpm). After incubation for 30min at room temperature, the reaction mixture was layered onto the top of 0.35ml of isotonic, 20% sucrose in buffered saline (138mM NaCl, 9.0mM KCl, 9.3mM CaCl<sub>2</sub>, 23.3mM MgCl<sub>2</sub>, 25.5mM MgSO<sub>4</sub>, 10mM EPPS, pH 8.3 pre-

pared in a 0.4ml test tube (4mm in diameter). Spermatozoa were spun down by centrifugation at 15,000 g for 5min. After removal of the supernatant, the radioactivity bound to the spermatozoa was measured with a gamma counter (Aloka Co. Ltd., Tokyo). For competition assays, the cold ligand of an adequate amount was mixed with the radioiodinated ligand before incubation with sperm.


Kinetic parameters for the binding of  $^{125}$ I-Fragment 1 were estimated by Scatchard analysis (Scatchard, 1949).

#### Photoaffinity Crosslink Experiments

Dry sperm was diluted 5-fold with ASW, and 50 $\mu$ l of the sperm suspension was mixed with 50 $\mu$ l of ASW with or without 1 mg Fragment 1 (ca. 100nmol). After incubation for 10min on ice, 2 $\mu$ g (ca. 2nmol) of Sulfo-SBED-conjugated Fragment 1 in 20 $\mu$ l ASW was added to the mixture and incubated in the dark for another 10min on ice. After the mixture was photoactivated with UV light (365 $\mu$ m) for 15min on ice, the spermatozoa were precipitated by centrifugation at 2,700 g for 1min. The precipitate was washed three times with 1ml ASW, and finally mixed with 50 $\mu$ l of double concentrated sample buffer for SDS-PAGE and boiled for 4min with mixing by a pipette to destroy DNA. Each sample of 5 $\mu$ l was subjected to 10% SDS-PAGE, and protein bands were detected with Western Blot with anti-biotin goat IgG (1mg/ml solution from Pierce Chemical Co., Rockford, IL; 1:1000) or normal goat IgG (0.5mg/ml solution from Santa Cruz Biotechnology Inc., Santa Cruz, CA; 1:500) as the 1st antibody, and anti-goat IgG donkey IgG conjugated with horseradish peroxidase (HRP) as the 2nd. The activity of HRP was detected by an ECL-plus system (Amersham Pharmacia Biotech AB, Uppsala, Sweden).

#### Chemical Analysis

Sugar content was determined by phenol-sulfuric acid method (Dubois *et al.*, 1956) or resorcinol-sulfuric acid method (Monsigny *et al.*, 1988), with L-fucose as a standard. The amount of protein was determined according to Bradford (1976) using bovine serum albumin as a standard.



**Fig. 3.** Real-time assay of the binding of intact sperm (a) and sperm membrane (b) to immobilized Fragment 1 by Biacore system. About 70pg of biotinyl Fragment 1 was immobilized by streptavidin fixed to the sensor chip surface. At the flow rate of 20 $\mu$ l/min, 40 $\mu$ l suspension of sperm (a) or sperm membrane (b) of the concentration as expressed on the corresponding sensogram was introduced to the chip surface.

## RESULTS

### Biological Activity of Fragment 1

The capacity to induce the acrosome reaction in high  $\text{Ca}^{2+}$  (50mM) sea water was estimated with Fragment 1 and P-ARIS. Fig. 1 shows that Fragment 1 alone induces the acrosome reaction significantly in high  $\text{Ca}^{2+}$  sea water though the specific activity is almost two-order lower than P-ARIS.

### Real-Time Monitoring of Ligand Binding to ARIS Receptor with Biacore System


Direct immobilization of about 0.2ng P-ARIS to the sensor tip surface caused the shift of almost 210RU (resonance units), whereas indirect immobilization of about 70pg biotinyl Fragment 1 resulted in the shift of 70RU. These values fit well with those of the standard Biacore assays, suggesting that the system is applicable to the study of ARIS receptor. As shown in Fig. 2, the binding of sperm membrane to immobilized P-ARIS is blocked by an excess of free P-ARIS, confirming the suitability of the Biacore system for the study of ARIS receptor. Fig. 3 shows that Fragment 1 indeed binds to the intact sperm and to the isolated membrane. The binding reaches a plateau within one to two minutes after the injection. Fig. 3 also shows that binding of intact sperm to the immobilized Fragment 1 causes much larger RU shifts than that of sperm membrane, which further supports the methodological validity.

### Stoichiometric Analysis of Ligand Binding to ARIS Receptor


Binding of  $^{125}\text{I}$ -P-ARIS to sperm was inhibited by a large

excess of cold P-ARIS and Fragment 1 as shown in Fig. 4. Here again, Fragment 1 is less potent than P-ARIS in the inhibition of  $^{125}\text{I}$ -P-ARIS binding to sperm. Since the cold ligands inhibit P-ARIS binding greatly but not completely, the binding to sperm should consist of two components at least, specific binding and non-specific one.

Since we cannot figure out the molecular size of P-ARIS, we have to use Fragment 1 for stoichiometric analysis despite its capacity to trigger the acrosome reaction and



**Fig. 4.** Competitive inhibition of  $^{125}\text{I}$ -P-ARIS binding to sperm with Fragment 1 and cold P-ARIS. Sperm ( $2.8 \times 10^9$  cells) was incubated with  $^{125}\text{I}$ -P-ARIS ( $0.22 \mu\text{g}$  as L-fucose,  $2.5 \times 10^4$  cpm) in the presence or absence of Fragment 1 (●) or cold P-ARIS (□) in  $50 \mu\text{l}$  of ASW for 30min. The radioactivity bound to sperm was plotted against the concentration of competitors.



**Fig. 5.** Purification by gel-filtration of 4-methoxyphenethylamine-conjugated Fragment 1 before (a) and after (b) radioiodination. Bio-Gel P-4 columns were used for purification. The size of column was  $1.0 \times 12.7 \text{ cm}$  (a) and  $1.0 \times 10 \text{ cm}$  (b). Fragment 1 represented by sugar content (—●—), 4-methoxyphenethylaminyl residue by UV absorption at 280nm (---△---), and radioactivity (---◇---) were monitored. Sugar content was estimated by resorcinol-sulfuric acid method and expressed by the absorption at 450nm. See Materials and Methods for the details of conjugation and radioiodination.



amine increases. However, the biological activity of Fragment 1 decreases as the oxidation proceeds (Koyota *et al.*, 1997). Taking the results of pilot experiments into account, we chose the procedure summarized in Materials and Methods to make the Sulfo-SBED-conjugate of partially oxidized Fragment 1 (Fig. 7a). Biacore assay showed that this derivative retained enough capacity to bind to the sperm (data not shown).

Fig 7b shows that three sperm proteins of 50-60kDa are cross-linked with the Sulfo-SBED conjugate. The crosslink was specifically blocked by an excess of intact Fragment 1. The band near 100kDa is not sensitive at all to the presence of intact Fragment 1. This band seems to be a sperm protein that has a structure partially at least similar to biotin because similar bands appear whenever avidin is used. The positive bands of 50-60kDa are not distinct, suggesting that they are glycoproteins.

## DISCUSSION

It is well established that saccharide chains of the egg coats play pivotal roles in the induction of acrosome reaction, but their structures remain to be determined except for some echinoderms (for reviews see Miller and Ax, 1990; Litscher and Wassarman, 1993; Mengerink and Vacquier, 2001). In these animals, a very long sulfated linear polysaccharide chain plays a key role in the induction of acrosome reaction and the sperm seem to recognize the highly ordered spatial arrangement of the sulfate moieties (Hoshi *et al.*, 1999). Despite of recent progress in structural analysis of these polysaccharide chains (Koyota *et al.*, 1997; Alves *et al.*, 1997, 1998; Vilela-Silva *et al.*, 1999), our understanding of the sperm receptors for them is not much improved. Indeed, no information is available on the kinetic parameters for FSP binding to REJ in sea urchins. Although the binding of FSP to REJ is reported (Vacquier and Moy, 1997), there is no direct evidence that the binding is really a specific one. Such parameters published for ARIS binding to starfish sperm were calculated by assuming the molecular size of ARIS to be  $1 \times 10^4$  kDa (Ushiyama *et al.*, 1993). This assumption seems not very bad, yet we do not know the real size even now.

In this paper, we have characterized the ARIS receptor by using Fragment 1, which is chemically well characterized, as the ligand. Although the biological activity of Fragment 1 is much lower than ARIS/P-ARIS, our data provide the rationale of using it as the ligand. Firstly, specific and quantitative binding of Fragment 1 to the intact sperm and isolated sperm membranes is shown by two methods of different principles (Figs. 3 and 4). Secondly, it competes quantitatively with P-ARIS for the specific binding to intact sperm (Fig. 4). Thirdly, and most importantly, specific labeling of some sperm proteins of 50-60kDa is achieved by photoaffinity crosslink technique using a Fragment 1 derivative. It is most likely that these proteins are the components of ARIS receptor, or at least those closely related to the receptor.

Analysis of these proteins is now under progress in our laboratory.

The kinetic parameters obtained in this study ( $K_d=15\mu\text{M}$ ,  $B_{\text{max}}=8.4 \times 10^4$  per cell) appear quite different from those that we published previously ( $K_d=2\text{nM}$ ,  $B_{\text{max}}=1 \times 10^2$  per cell) on the assumption that the molecular size of ARIS is  $1 \times 10^4$  kDa. However, the apparent difference may not be irrational if the clustering effect, which is generally observed with protein-carbohydrate interactions (for a review see Lee, 1992), is taken into account; Fragment 1 itself has a repetitive structure (Koyota *et al.*, 1997) and ARIS seems to have many, extremely long saccharide chains ending with 200 repeats or so of Fragment 1 (Hoshi *et al.*, 1994, 1999). In many cases, multivalent binding of low affinity single sites generates high affinity and specificity of protein-carbohydrate interactions. Thus, very high affinity of intact ARIS and P-ARIS may be generated by its extreme polyvalence. Indeed, the activity of ARIS fragments correlates to their size, and the minimal functional size of ARIS is known to be almost the size of Fragment 1 (Ushiyama *et al.*, 1995).

This paper will contribute also to solve some technical problems in glycobiology. Firstly, we developed a procedure to conjugate sulfated sugar chains such as Fragment 1 with the photoaffinity crosslinker, Sulfo-SBED. This method has advantages over those we tried but failed previously; the most important two are the multiplicity of crosslink sites in the derivative and easiness in purification and detection of the derivative due to the presence of biotin. Needless to say, it is essential for any method of ligand modification for specific binding assays that the derivative is free of the original ligand. When we designed the methods described above, this point was carefully considered. Secondly, we developed a simple, reproducible and efficient procedure of radioiodination of sulfated saccharide chains by 4-methoxy-phenetylation of the reducing terminal followed by iodination at C3 and C5 of the benzene ring. Thirdly, we showed that the Biacore system is useful for the study of sperm-egg glycoconjugate interactions.

Identification of ARIS receptor protein(s) using the procedure developed in the present work will expose a new cutting-edge in the study of the molecular mechanism of the induction of acrosome reaction.

## ACKNOWLEDGEMENT

We greatly appreciate Dr. Y. Hatanaka, Toyama Medical and Pharmaceutical University, and Dr. K. Sakurai, Seikagaku Corporation, Tokyo for providing Biotin-*N*-BOC-phenylaminodeazirine. We wish to thank Dr. H. Hashimoto, Tokyo Institute of Technology for his valuable advice on chemical modification of saccharide chains. We thank the directors and staff of Otsuchi Marine Research Center and Misaki Marine Biological Station of University of Tokyo, Ushimado Marine Biological Station of Okayama University, and Marine and Coastal Research Tasmania, Australia, for their help in collecting starfish. This work was supported in part by Grants-in-Aid for Scientific Research on Priority Areas (#10178102, #12045225) from



the Ministry of Education, Science, Culture and Sports, Japan and a Research Grant from Mizutani Foundation for Glycoscience.

## REFERENCES

- Alves AP, Mulloy B, Diniz JA and Mourao PA (1997) Sulfated polysaccharides from the egg jelly layer are species-specific inducers of acrosomal reaction in sperms of sea urchins. *J Biol Chem* 272: 6965–6971
- Alves AP, Mulloy B, Moy GW, Vacquier VD and Mourao PA (1998) Females of the sea urchin *Strongylocentrotus purpuratus* differ in the structures of their egg jelly sulfated fucans. *Glycobiology* 8: 939–946
- Bradford MM (1976) A rapid and sensitive method for the quantization of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem* 72: 248–254
- Dan JC (1967) Acrosome reaction and lysins. In "Fertilization, volume 1: Comparative morphology, Biochemistry and Immunology" Ed by CB Metz and A Monroy, Academic Press, New York, pp 237–293
- Dubois M, Gilles KA, Hamilton JK, Rebers PA and Smith F (1956) Colorimetric method for determination of sugars and related substances. *Anal Chem* 28: 350–356
- Green NM (1965) A spectrophotometric assay for avidin and biotin based on binding of dyes by avidin. *Biochem J* 94: 23c–24c
- Hirohashi N and Vacquier VD (2002) Egg sialoglycans increase intracellular pH and potentiate the acrosome reaction of sea urchin sperm. *J Biol Chem* 277: 8041–8047
- Hoshi M, Nishigaki T, Ushiyama A, Okinaga T, Chiba K and Matsumoto M (1994) Egg-jelly signal molecules for triggering the acrosome reaction in starfish spermatozoa. *Int J Dev Biol* 38: 167–174
- Hoshi M, Kawamura M, Maruyama Y, Yoshida E, Nishigaki T, Ikeda M, Ogiso M, Moriyama H and Matsumoto M (1999) How does the jelly coat of starfish eggs trigger the acrosome reaction in homologous spermatozoa? In "The Male Gamete: From Basic Science to Clinical Applications" Ed by C Gagnon, Cache River Press, Vienna, IL, pp 119–125
- Hunter WH and Greenwood FC (1962) Preparation of iodine-131 labelled human growth hormone of high specific activity. *Nature* 194: 495–496
- Ikada H and Hoshi M (1981a) Biochemical studies on the acrosome reaction of the starfish, *Asterias amurensis*. 1. Factors participating in the acrosome reaction. *Dev Growth Differ* 23: 73–80
- Ikada H and Hoshi M (1981b) Biochemical studies on the acrosome reaction of the starfish, *Asterias amurensis*. 2. Purification and Characterization of acrosome reaction-inducing substance. *Dev Growth Differ* 23: 81–88
- Johnsson B, Lofas S and Lindquist G (1991) Immobilization of proteins to a carboxymethyl-dextran-modified gold surface for bio-specific interaction analysis in surface plasmon resonance sensors. *Anal Biochem* 198: 268–277
- Koyota S, Wimalasiri KMS and Hoshi M (1997) Structure of the main saccharide chain in the Acrosome Reaction-Inducing Substance of the starfish, *Asterias amurensis*. *J Biol Chem* 272: 10372–10376
- Lee YC (1992) Biochemistry of carbohydrate-protein interaction. *FASEB J* 6: 3193–3200
- Litscher ES and Wassarman PM (1993) Carbohydrate-mediated adhesion of eggs and sperm during mammalian fertilization. *Trends Glycosci Glycotech* 5: 369–388
- Longo FJ, Ushiyama A, Chiba K and Hoshi M (1995) Ultrastructural localization of Acrosome Reaction-Inducing Substance (ARIS) on sperm of the starfish *Asterias amurensis*. *Mol Reprod Dev* 41: 91–99
- Longo FJ (1997) Fertilization. 2nd ed, Chapman and Hall, London
- Matsui T, Nishiyama I, Hino A and Hoshi M (1986) Induction of the acrosome reaction in starfish. *Dev Growth Differ* 28: 39–48
- Mengerink KJ, Moy GW and Vacquier VD (2000) suREJ proteins: new signaling molecules in sea urchin spermatozoa. *Zygote* 8 (Suppl 1): S28–S30
- Mengerink KJ and Vacquier VD (2001) Glycobiology of sperm-egg interactions in deuterostomes. *Glycobiology* 11: 37R–43R
- Miller DJ and Ax RL (1990) Carbohydrates and fertilization in animal. *Mol Reprod Dev* 26: 184–189
- Monsigny M, Petit C and Roche AC (1988) Colorimetric determination of neutral sugars by a resorcinol sulfuric acid micromethod. *Anal Chem* 175: 525–530
- Moy GW, Mendoza LM, Schulz JR, Swanson WJ, Glabe CG and Vacquier VD (1996) The sea urchin sperm receptor for egg jelly is a modular protein with extensive homology to the human polycystic kidney disease protein, PKD1. *J Cell Biol* 133: 809–817
- Nishiyama I, Matsui T and Hoshi M (1987a) Purification of Co-ARIS, a cofactor for acrosome reaction-inducing substance, from egg jelly of starfish. *Dev Growth Differ* 29: 161–169
- Nishiyama I, Matsui T, Fujimoto Y, Ikekawa N and Hoshi M (1987b) Correlation between the molecular structure and the biological activity of Co-ARIS, a cofactor for acrosome reaction-inducing substance. *Dev Growth Differ* 29: 171–176
- Nishigaki T, Chiba K, Miki W and Hoshi M (1996) Structure and function of asterosaps, sperm-activating peptides from the jelly coat of starfish eggs. *Zygote* 4: 237–245
- Scatchard G (1949) The attractions of proteins for small molecules and ions. *Ann NY Acad Sci* 51: 660–672
- Shinohara Y, Sota H, Kim F, Shimizu M, Gotoh M, Tosu M and Hasegawa Y (1995) Use of a biosensor based on surface plasmon resonance and biotinyl glycans for analysis of sugar binding specificities of lectins. *J Biochem* 117: 1076–1082
- Uno Y and Hoshi M (1978) Separation of the sperm agglutinin and the acrosome reaction-inducing substance in egg jelly of starfish. *Science* 200: 58–59
- Ushiyama A, Araki T, Chiba K and Hoshi M (1993) Specific binding of acrosome reaction-inducing substance to the head of starfish spermatozoa. *Zygote* 1: 121–127
- Ushiyama A, Shima A and Hoshi M (1995) Estimation by radiation inactivation of the minimum functional size of acrosome reaction-inducing substance (ARIS) in the starfish, *Asterias amurensis*. *Zygote* 3: 351–355
- Vacquier VD and Moy GW (1997) The fucose sulfate polymer of egg jelly binds to sperm REJ and is the inducer of the sea urchin sperm acrosome reaction. *Dev Biol* 192: 125–135
- Vilela-Silva AC, Alves AP, Valente AP, Vacquier VD and Mourao PA (1999) Structure of the sulfated alpha-L-fucan from the egg jelly coat of the sea urchin *Strongylocentrotus franciscanus*: patterns of preferential 2-O- and 4-O-sulfation determine sperm cell recognition. *Glycobiology* 9: 927–933
- Yamaguchi M, Niwa T, Kurita M and Suzuki N (1987) The participation of speract in the acrosome reaction of *Hemicentrotus pulcherrimus*. *Dev Growth Differ* 30: 159–167

(Received January 31, 2002 / Accepted February 25, 2002)