BioOne COMPLETE

The Drumming of Isoperla pallida Aubert and Guadalgenus franzi (Aubert) (Plecoptera, Perlodidae), and Review and Evolutionary Considerations of Southern Iberian Peninsula Perlodidae

Authors: de Figueroa, J. M. Tierno, Luzón-Ortega, J. M., and Stewart, K. W.

Source: Zoological Science, 19(8) : 871-875

Published By: Zoological Society of Japan

URL: https://doi.org/10.2108/zsj.19.871

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Downloaded From: https://complete.bioone.org/journals/Zoological-Science on 28 Mar 2024
Terms of Use: https://complete.bioone.org/terms-of-use

ZOOLOGICAL SCIENCE **19**: 871–875 (2002) © 2002 Zoological Society of Japan

The Drumming of *Isoperla pallida* Aubert and *Guadalgenus franzi* (Aubert) (Plecoptera, Perlodidae), and Review and Evolutionary Considerations of Southern Iberian Peninsula Perlodidae

J. M. Tierno de Figueroa[1]*, J. M. Luzón-Ortega[1] and K. W. Stewart[2]

[1]*Departamento de Biología Animaly Ecología, Facultad de Ciencias, Universidad de Granada, 18071 Granada, Spain*
[2]*Department of Biological Sciences, University of North Texas, Denton, TX 76203, USA*

ABSTRACT—The drumming calls of two perlodid species from Southern Spain (Sierra Morena, Jaén) are analyzed and described. Nineteen calls from one male of *Guadalgenus franzi* were recorded, showing a simple call pattern composed of two beats with an average duration between them of 184.32 msec (range: 165–227). This is the simplest call pattern yet discovered in the Plecoptera. The female of this species also presented a simple call consisting of only one beat. The male of *Isoperla pallida* showed an ancestral call pattern, more simple than the other Iberian *Isoperla* species (*I. grammatica, I. nevada* or *I. curtata*, all of them with multiphasic drumming calls or *I. bipartita* with a notably different model consisting of rubbing). Calls from six different males (the female did not drum) consisted of an average of 65.94 beats (range: 25–83) with a duration average of 1462.69 msec (range: 548–1835), regularly repeated with intervals with a duration average of 22.27 msec (range: 17–30). With this study, the vibrational calls of all the Perlodid species from the Southern Iberian Peninsula, except *Perlodes microcephalus* and *Besdolus bicolor*, are known. Moreover, it presents the first record and description of the call of the monospecific genus *Guadalgenus*. An evolutionary approach to the vibrational communication diversification in the perlodids of the Southern Iberian Peninsula is presented, showing a complexity graduation from the more simple drumming calls (shown by *Hemimelaena flaviventris* and *G. franzi)* to the rubbing call of *I. bipartita* and intermediate drumming calls of other *Isoperla* species, consistent with the drumming evolution paradigm of Stewart (2001) .

Key Words: Vibrational communication, drumming, stonefly, Perlodidae, Plecoptera

INTRODUCTION

Intersexual communication using low-frequency, substrate-borne vibrations is a little explored mode of communication in arthropods (Stewart, 1997, 2001). Intersexual vibrational communication appears to have evolved its most complex form in the Arctoperlarian Plecoptera as a mechanism for mate encounter and pair forming (Stewart *et al*., 1995; Stewart, 1997). Stoneflies produce vibrational signals by percussion (drumming), rubbing (or scraping), tremulation (abdominal or body jerking), or combinations of these methods (Stewart, 1997). Vibrational calls are genetically programmed and can be considered as a fixed-action, species-specific behavioral pattern (Szczytko and Stewart, 1979a, 1979b), useful for inferring phylogenies within the order (Stewart and Zeigler, 1984). Particularly interesting, is the study of the effects of evolutionary pressures in the diversification of vibrational calls among species (inside the same genus and between related genera) from the same family sharing the same geographical distribution area.

Stonefly drumming has been known from the XIX century (Newport, 1851), and was first quantified some decades ago by Rupprecht (1968). Since then, the vibrational calls of about 160 world species have been studied, principally in North America (Stewart, 1994) and, to a lesser extent, in Europe and Japan.

Nine species of Perlodidae exist in the Southern Iberian Peninsula: *Besdolus bicolor* (Navás, 1909), *Guadalgenus franzi* (Aubert, 1963), *Perlodes microcephalus* (Pictet, 1833), *Hemimelaena flaviventris* (Pictet, 1841), *Isoperla bipartita* Aubert, 1962, *I. curtata* Navás, 1924, *I. grammatica* (Poda, 1761), *I. nevada* Aubert, 1952 and *I. pallida* Aubert, 1963 (Sánchez-Ortega and Tierno, 1996).

* Corresponding author: Tel. 0034-958242372;
FAX. 0034-958243238.
E-mail: acuatica@ugr.es

Downloaded From: https://complete.bioone.org/journals/Zoological-Science on 28 Mar 2024
Terms of Use: https://complete.bioone.org/terms-of-use

G. franzi and *H. flaviventris* represent monospecific stonefly genera, the first one with a Iberic distribution and the second one with a Iberic-North African distribution (Sánchez-Ortega and Alba-Tercedor, 1987). The drumming calls of the *G. franzi* are unknown. *H. flaviventris* drumming calls were described by Tierno de Figueroa et al. (1998). *P. microcephalus* is widely distributed European species, but is a relict one in the South of Iberian Peninsula, where it is only present in scarce number in the Sierra Nevada (Sánchez-Ortega and Alba-Tercedor, 1987), and *B. bicolor* is an Iberian endemic species, scarcely distributed by Central Spain, with a punctual cite in the South (Zwick and Weinzierl, 1995). The drumming calls of these two species have not been previously recorded. *Isoperla* is an Holarctic Plecoptera genus with a high number of species. Four Iberic endemic species are present in the Southern Iberian peninsula: *I. bipartita*, *I. curtata*, *I. nevada* and *I. pallida* (Sánchez-Ortega and Alba-Tercedor, 1987). The drumming calls of the first three of them have been studied (Tierno de Figueroa and Luzón-Ortega, 2002; Tierno de Figueroa et al., 2000; Tierno de Figueroa and Sánchez-Ortega, 1999). Another *Isoperla* species, *I. grammatica*, with a wide European distribution is also present (Sánchez-Ortega and Alba-Tercedor, 1987). The drumming call of this species has been studied in other parts of its distribution area: Central Europe by Rupprecht (1968, 1969, 1983, 1984) and Northern Spain by Membiela and Vidal (1998).

The flight period of all the perlodid species from Southern Iberian Peninsula is spring-summer, with maximum peaks from May to July, with slight variations in relation to the altitude and latitude (Sánchez-Ortega and Alba-Tercedor, 1987). Only, in the case of *I. nevada* and *I. grammatica* in Sierra Nevada (S Spain) a temporal segregation (and also a spatial one) has been detected with a slight seasonal difference between their capture maximums, later in *I. nevada* (Tierno *et al.*, 1994).

In the present paper, we describe for the first time the drumming calls of *G. franzi* and *I. pallida* and review the vibrational communication of the seven Perlodidae species from the Southern Iberian Peninsula (all of them with hammer or ventral lobe, see Rupprecht, 1976, and with a size oscillating 7–12 mm for *Isoperla* genus, 10–14 mm for *H. flaviventris* and 12–18 mm for *G. franzi*, according with the species descriptions), whose calls are known in light of the drumming evolution paradigm proposed by Stewart (2001), in which Arctoperlarian stoneflies represent the epitome of evolution of vibrational communication. Their ancestral signals were monophasic volleys of evenly spaced drumbeats. Derived signals to achieve species-specificity and possibly enable sexual selection or some measure of reproductive fitness as involved modification of the ancestral form toward complex signals through: changes in the rhythmic patterning of calls, patterns of male-female duetting, and/or changes in the method of signal production such as rubbing or tremulation.

MATERIALS AND METHODS

The specimens of *Guadalgenus franzi* and *Isoperla pallida* were collected in the Martín Pérez seasonal stream from the Sierra Morena (Jaén, Spain), on May, first, 2001. The specimens were transported individually to the laboratory in bottles kept in an ice chest. The drumming signals were recorded over the following four days between 15:30 to 18:00 local time at a temperature between 23 and 26°C, and under natural light entering the room through a window.

Male calls of the two species were recorded by placing individuals in a 2.5×2.5×2.5 cm. bottomless plastic box resting directly on a 7.5 cm diameter, 8 ohm, 0.8 W speaker connected to a Sony CFS-W350L model cassette-recorder. We placed the female close to the male in a separate bottomless plastic box similar to the one containing the male, permitting the pair to communicate with each other with vibrations. Previously to the record, females of both species never were observed drumming when they were in the bottles, such it occurs with the males. Recorded calls were displayed and measured using the IBM-compatible computer program Creative Wave Studio for Windows 1993.

RESULTS

Only a male and a female of *Guadalgenus franzi* were studied. A total of 19 drumming calls from the male (Fig. 1) was obtained, showing a simple call pattern (Stewart, 1997, 2001). The vibrational call was composed of only two beats with an average duration between them of 184.32 msec (range: 165–227; SD=17.11; N=19). This call is simpler than the one described for *Hemimelaena flaviventris* (Tierno de Figueroa *et al.*, 1998), and the simplest yet discovered in Plecoptera.





Fig. 1. *Guadalgenus franzi* male and female drumming call.

The female of this species also presented a simple call consisting of only one beat (N=8). Thus, the female call is also simpler than the one presented by *H. flaviventris,* composed of 1 to 3 beats (Tierno de Figueroa *et al.*, 1998), and similar to the one-beat answers of the North American species *Perlinella drymo* (Newman, 1839) (Zeigler and Stewart, 1977).

It was not observed overlapping between male and female calls. Intervals between male call and female answer were very variable and oscillated between 215 msec. and more than 4000 msec. Moreover, in one case the female repeated the answer twice (with intervals of 376 and 775

Downloaded From: https://complete.bioone.org/journals/Zoological-Science on 28 Mar 2024
Terms of Use: https://complete.bioone.org/terms-of-use



Isoperla pallida male call

Fig. 2. *Isoperla pallida* male drumming call.

msec.).

Isoperla pallida (Fig. 2) also showed an ancestral call pattern (Stewart, 1997) with simple modifications of percussion increasing the beat number (Stewart, 2001), extremely long (only surpassed by the *I. nevada* call) and simpler than other studied Iberic *Isoperla* species. Thus, *I. grammatica*, *I. nevada* and *I. curtata*, present multiphasic drumming calls (Membiela and Vidal, 1998; Tierno de Figueroa and Sánchez-Ortega, 1999; Tierno de Figueroa *et al*., 2000) and *I. bipartita,* exhibit a notably different vibrational communication pattern consisting in rubbing (Tierno de Figueroa and Luzón-Ortega, 2002). The male drumming call of *I. pallida* consisted of 65.94 beats (range: 25-83; SD=15.5; N=16) with a duration of 1462.69 msec (range: 548-1835; SD=352.56; N=16) regularly repeated with intervals with a mean value of 22.27 msec (range: 17-30; SD=1.74; N=974), obtained from six different males (from more than 10 studied specimens).

Although five females of *I. pallida* were also located close to the males in separate plastic box, they never drummed.

DISCUSSION

It has been pointed out that intersexual vibrational calls are species-specific (Szczytko and Stewart, 1979a, 1979b) and could play an important role as an isolating reproductive mechanism (Tierno de Figueroa and Sánchez-Ortega, 1999). This idea could be supported by the fact that from the nine perlodid species present in the Southern Iberian Peninsula, almost all of them can share habitat. Thus, *I. curtata* has been collected by us from the same locality and with the same flight period as *H. flavinentris, I. pallida* and *G. franzi.* The same thing occurs with *I. curtata* and *I. bipartita* in other streams and potentially all of them can share habitat with *I. grammatica* (Sánchez-Ortega and Alba-Tercedor, 1987). *I. grammatica* and *I. nevada* seem to have a spatial and temporal segregation between them (Tierno *et al*., 1994). *P. microcephalus* and *B. bicolor* present a punctual distribution in Southern Iberian Peninsula (Sánchez-Ortega and Alba-Tercedor, 1987; Zwick and Weinzierl, 1995).

Rupprecht (1969, 1983) noted that sympatric species of *Isoperla* showed more different call models than the allopatric ones. This idea could be supported by the call analysis of studied species in Southern Iberian Peninsula (Table 1). In fact, we have detected a gradual complexity in the vibrational calls of all these perlodids. *H. flaviventris* and *G. franzi* are monospecific genera and show call patterns deviating only by number of beats and intervals from ancestral patterns (Stewart, 2001) (Fig. 3). In other genera, such as *Isoperla*, where some species coexist, the call must be subject to selective pressures related with its role as an isolating reproductive mechanism.

Fig. 3 shows a theoretical evolution of Southern Iberian male perlodid species: increasing or decreasing beat number, changes of intervals or rhythm inside a sequence, division of calls into phases, and derived signal production by rubbing. This model is consistent with the evolution paradigm of Stewart (2001). These data (Table 1; Figs. 1–3) corroborate the previously proposed hypothesis that in male Arctoperlarian stoneflies, sexual selection acts to favor derivation from the ancestral monophasy to calls with altered (more or fewer) beats, distinct interbeat intervals (often phased and/or grouped) and/or changed method of vibration production such as with *I. bipartita* rubs (Table 1; Fig. 3). In many cases, these changes lead to calls that are more complex than ancestral ones.

Stewart *et al*. (1995) indicated that seven species of stoneflies in different families and genera have derived what might be considered “evolutionarily converged” calls into 3-beat signals. They hypothesized that this trend toward deriving such 3-beat calls (particularly in the families Peltoperlidae, Perlodidae and Perlidae) might represent a minimal amount of information (two interbeat time intervals) necessary for female recognition of potential calling conspecific mates. If correct, 3-beat calls would represent an evolutionary end-point of derived reduction of beats, with their corresponding interval alterations, of ancestral signals that could achieve communicative isolation. They pointed out that the intervals of the seven species known to have 3-beat calls were species-specific, without overlap, and that none of them was known to co-occur in time and place. Our discovery here of the 2-beat call of *G. franzi* now demonstrates that only a single time interval is required, at least for females of some species, for female recognition. And, if time intervals are important recognition parameters in percussion-produced signals, as indicated by computer-simulation models (Stewart and Maketon, 1990), then this 2-beat call of *G. franzi* would represent the simplest possible percussion-produced call that could be effectively derived from monophasic ancestral signals in a drumming mode of communication.

One-beat answers of females, such as here in *G. franzi*, are possible because such a simple female answer conveys information to the male only that she is responding with location to her recognition that he is a conspecific caller.

Downloaded From: https://complete.bioone.org/journals/Zoological-Science on 28 Mar 2024
Terms of Use: https://complete.bioone.org/terms-of-use

Table 1. Vibrational call description of seven Southern Iberian Perlodid species. Seq= sequence; Ph= phase; N= number; T= time; Int= interval. * For *I. grammatica*, we have considered the Membiela & Vidal (1998) (*) description based in Spanish *I. grammatica* specimens, but Ruprecht (1968, 1984) (**) pointed a highly variable drumming call for this species in Central Europe.

	males								females		
Species	**Call production**	**N. seq/ call**	**N. ph/ seq**	**T. seq & ph (msec) Mean (range)**	**N. beats or rubs/seq & ph Mean (range)**	**Beat Int. reduction**	**Seq Int. Mean (range)**	**Final beat**	**Call production**	**N. beats Mean (range)**	**Beat Int Mean (range)**
G. franzi	Drumming	1	1	182.32 (165–227)	2	–	–	No	Drumming	1	–
H. flaviventris	Drumming	1	1	204.64 (176–261)	3	Yes	–	No	Drumming	1.41 (1–3)	100.91 (68–227)
I. bipartita	Rubbing	6.73 (3–10)	1	257.74 (24–594)	3.11 (1–4)	Yes	180.75 (142–290)	No	Drumming Rubbing	1 beat + 1 rub	479.09 (258–544)
I. curtata	Drumming	1	2	792.9 (228–1312) Phase I: 649.9 (149–1140) Interphase: 17 (13–24) Phase II: 118.2 (37–185)	17.3 (8–27) Phase I: 9.5 (5–17) Phase II: 8.1 (3–12)	Phase I: yes Phase II: No	–	No	Tremulation?	–	–
I. grammatica	Drumming	6.70 (5–8)* 14 (2–15)**	2	123 (25–334)* Phase I: 65 (25–122)* Phase II: 58 (0–212)*	7.48 (2–11)* Phase I: 2* Phase II: 5.48 (0–9)*	Phase I: yes* Phase II: No*	173.6 (68–469)*	No*	No detected	No detected	No detected
I. nevada	Drumming	2.73 (1–5)	2	1595–4028 **Seq I** Phase I: 3446 Interphase: 19 Phase II: 563 **Seq II-V** Phase I: 853–1089 Interphase: 23–36 Phase II: 719–827	**Seq I** Phase I: 43.3 Phase II: 54.9 **Seq II-V** Phase I: 13–22 Phase II: 50–62	No	134–236	Yes (1–12)	No detected	No detected	No detected
I. pallida	Drumming	1	1	1462.69 (548–1835)	65.94 (25–83)	–	–	No	No detected	No detected	No detected




Fig. 3. Possible evolution of vibrational call complexity in Southern Iberian Perlodids.

Downloaded From: https://complete.bioone.org/journals/Zoological-Science on 28 Mar 2024
Terms of Use: https://complete.bioone.org/terms-of-use

She is not required to give more complex information on her species identity, and in fact such derivation toward simpler answer is less energy demanding, thus saving resources for egg-yolking and oviposition activity. The duet descriptions (Table 1) show for regional species a simplification of female answer (*H. flaviventris* 1-3 beats; *G. franzi* 1-beat) or change in method of answering (*I. bipartita* 1 beat + 1 rub; *I. curtata* possible tremulation). In other species, no answer has been successfully detected or recorded.

In Central Europe, where *Hemimelaena* and *Guadalgenus* genera are not present, there are some *Isoperla* species (such as *I. rivulorum* in Rupprecht, 1969, or *I. difformis* in Rupprecht, 1981) whose males show a similar call pattern to them, supporting the importance of the reproductive isolating mechanism among sympatric species. Likewise, in the pointed Central Europe species, female answers have been detected that show relatively similar patterns to those of *Hemimelaena* and *Guadalgenus.*

AKNOWLEDGEMENTS

The authors want to thank J. Pérez Contreras and M. García Sánchez for their collaboration in the field sampling.

REFERENCES

Membiela P, Vidal M (1998) Las llamadas sexuales de *Isoperla grammatica* (Poda, 1761) (Plecoptera: Perlodidae). Boln Asoc esp Ent 22(1–2): 15–21

Newport G (1851) On the anatomy and affinities *Pteronarcys regalis*, Newm.: with a postscript containing descriptions of some American Perlidae, together with notes on their habits. Trans Linn Soc 20: 425–451

Rupprecht R (1968) Das Trommeln der Plecopteren. Z Vergl Physiol 59: 38–71

Rupprecht R (1969) Zur Artspezifität der Trommelsignale der Plecopteren (Insecta). Oikos 20: 26–33

Rupprecht R (1976) The structure (ultra) and function of the ventral lobe and the hammer of Plecoptera. Proc Biol Soc Wash 88: 415

Rupprecht R (1981) A new system of communication within Plecoptera and a signal with a new significance. Proceedings of the Seventh International Symposium on Plecoptera. Nara Women's University, 19–22, August, 1980

Rupprecht R (1983) Kommunicationssignale von Arten der Gattung *Isoperla* (Plecoptera) und deren Einung zur Abgrenzung von Arten. Verh. Dtsch Zool Ges 1983: 198

Rupprecht R (1984) *Isoperla grammatica* Poda 1761. Beschreibung eines Neotypus (Plecoptera). Annls Limnol 20(1–2): 81–90

Sánchez-Ortega A, Alba-Tercedor J (1987) Lista faunística y bibliográfica de los Plecópteros de la Península Ibérica. Asociación española de Limnología. Lista de la flora y fauna de las aguas continentales de la Península Ibérica 4: 1–133

Sánchez-Ortega A, Tierno JM (1996) Current situation of stonefly fauna (Insecta: Plecoptera) in the Iberian Peninsula and Balearic Islands. Mitt Schweiz Ent Ges 69: 77–94

Stewart KW (1994) Theoretical considerations of mate finding and other adult behaviours of Plecoptera. Aquat Insects 16: 95–104

Stewart KW (1997) Vibrational Communication in Insects. Epitome in the Language of Stoneflies? Am Entomol, summer 1997: 81–91

Stewart KW (2001) Vibrational communication (drumming) and mate-searching behavior of stoneflies (Plecoptera); evolutionary considerations. In "Trends in Research in Ephemeroptera and Plecoptera" Ed by E Domínguez, Kluwer Academic/Plenum Publishers, pp 217–225

Stewart KW, Abbott JC, Bottorff RL (1995) The drumming signals of two stonefly species *Cosumnoperla hypocrena* (Perlodidae) and *Paraperla wilsoni* (Chloroperlidae); a newly discovered duet pattern in Plecoptera. Ent News 106(1): 13–18

Stewart KW, Maketon M (1990) Intraspecific variation and information content of drumming in three Plecoptera species. In "Mayflies and Stoneflies" Ed by IC Campbell, Kluwer Academic Publishers, pp 259–268

Stewart KW, Zeigler DD (1984) The use of larval morphology and drumming in Plecoptera systematics, and further studies of drumming behavior. Ann Limnol 20(1-2): 105-114

Szczytko SW, Stewart KW (1979) Drumming behavior of four western Nearctic *Isoperla* (Plecoptera) species. Ann Ent Soc Am 72(6): 781–786

Szczytko SW, Stewart KW (1979) Stonefly drumming as a model classroom study of aquatic insect behavior. In "Innovative teaching in aquatic entomology" Ed by VH Resh, DM Rosemberg, Can Spec Publ Fish Aquat Sci 43: 1–118, pp 31–37

Tierno de Figueroa JM, Luzón-Ortega JM (2002) The drumming call of *Isoperla bipartita* Aubert, 1962 (Plecoptera, Perlodidae). Aquat Insects 24: 87–90

Tierno de Figueroa JM, Luzón-Ortega JM, Sánchez-Ortega A (1998) Imaginal biology of *Hemimelaena flaviventris* (Pictet, 1841) (Plecoptera, Perlodidae). Ann Zool fennici 35: 225–230

Tierno de Figueroa JM, Luzón-Ortega JM, Sánchez-Ortega A (2000) Calling, mating and oviposition in *Isoperla curtata* (Plecoptera, Perlodidae). Eur J Entomol 97: 171–175

Tierno de Figueroa JM, Sánchez-Ortega A (1999) The male drumming call of *Isoperla nevada* Aubert, 1952 (Plecoptera, Perlodidae). Aquat Insects 21: 33–38

Tierno JM, Sánchez-Ortega A, Picazo-Muñoz J (1994) Sobre la distribución espacial y temporal de *Isoperla nevada* Aubert, 1952 (Insecta,Plecoptera). Zool baetica 5: 107–112

Zeigler DD, Stewart KW (1977) Drumming behavior of eleven nearctic stonefly (Plecoptera) species. Ann Ent Soc Am 70(4): 495–505

Zwick P, Weinzierl A (1995) Reinstatement and revision of genus *Besdolus* (Plecoptera: Perlodidae). Ent Scand 26(1): 1–16

(Received January 29, 2002 / Accepted May 22, 2002)

Downloaded From: https://complete.bioone.org/journals/Zoological-Science on 28 Mar 2024
Terms of Use: https://complete.bioone.org/terms-of-use