

Transfer of *Hymenoxys argentea* var. *Thoreauensis* to *Tetraneuris* (Asteraceae: Helenieae: Tetraneuridinae)

Authors: Bierner, Mark W., and Turner, Billie L.

Source: Lundellia, 2010(13) : 1-2

Published By: The Plant Resources Center, The University of Texas at Austin

URL: <https://doi.org/10.25224/1097-993X-13.1.1>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

TRANSFER OF *HYMENOXYS ARGENTEA* VAR. *THOREAUENSIS* TO *TETRANEURIS* (ASTERACEAE: HELENIEAE: TETRANEURIDINAE)

Mark W. Bierner^{1,2} and Billie L. Turner²

¹School of Natural Resources and the Environment and School of Plant Sciences, University of Arizona, Tucson, Arizona 85721

²Plant Resources Center, 1 University Station F0404, The University of Texas, Austin, Texas 78712
bierner@ag.arizona.edu

Abstract: Based on evidence supporting the recognition of *Tetraneuris* as a genus separate from *Hymenoxys*, *Hymenoxys argentea* var. *thoreauensis* is basionym of a new combination, *Tetraneuris argentea* var. *thoreauensis*.

Keywords: Asteraceae, Helenieae, Tetraneuridinae, *Hymenoxys*, *Tetraneuris*.

Hymenoxys argentea (A. Gray) K. F. Parker var. *thoreauensis* N. D. Atwood, S. L. Welsh & A. Clifford was applied by Atwood and Welsh (2007) to discoid plants growing north of Thoreau, New Mexico, "... with other mound-forming plants on a remaining lens of the windswept Todilto Limestone with limy sands and limestone gravel on the point of the Mesa above the limestone cliffs."

Atwood and Welsh (2007) stated, "We maintain use of *Hymenoxys* over the segregate *Tetraneuris* out of personal choice and belief in a stable nomenclature." It is their prerogative to use *Hymenoxys* rather than *Tetraneuris*. How this helps maintain a stable nomenclature is unclear to us, given that the most recent and most comprehensive treatments of the group to which these plants are referable recognize *Tetraneuris* as a genus separate from *Hymenoxys* (Bierner and Turner, 2003; Panero, 2007).

Some relatively recent workers have recognized *Tetraneuris* as an infrageneric grouping within *Hymenoxys* (e.g., Barkley, 1986; Cronquist, 1994; Karis and Ryding, 1994; Welsh, 1987). Other recent workers have recognized *Tetraneuris* at the generic level (e.g., Baldwin and Wessa, 2000; Baldwin et al., 2002; Bierner, 1994; Bierner and Jansen, 1998; Bierner and Turner, 2003; Spring et al., 1994). Morphological and chemical differences between taxa referable to *Tetraneuris* and taxa referable to *Hymenoxys* have been discussed at length (e.g.,

Bierner, 1978, 2001; Bierner et al., 1992; Díaz et al., 1992; Ferracini et al., 1994).

Atwood and Welsh (2007) cited Bierner and Turner's (2006) treatment for *Flora of North America North of Mexico*, but they did not cite Bierner and Turner's (2003) taxonomic treatment, which presented multiple lines of evidence for the recognition of *Tetraneuris*, nor did they cite Panero (2007) or Baldwin et al. (2002). The latter not only supports recognition of *Tetraneuris*, it corroborates Bierner (1994) and Bierner and Jansen (1998), supporting recognition of *Tetraneuris*.

TAXONOMY

Tetraneuris argentea (A. Gray) Greene var. ***thoreauensis*** (N. D. Atwood, S. L. Welsh & A. Clifford) Bierner & B. L. Turner, comb. nov.

BASIONYM: *Hymenoxys argentea* (A. Gray) K. F. Parker var. *thoreauensis* N. D. Atwood, S. L. Welsh & A. Clifford, *Rhodora* 109: 395–414, 2007. TYPE: **UNITED STATES. NEW MEXICO: McKinley Co.:** North of Thoreau, 27 May 2003, A. Clifford 03-385 (HOLOTYPE: BRY; ISOTYPES: ASU! MO—as database image!).

ACKNOWLEDGMENT

We thank John Strother for his help.

LITERATURE CITED

- Atwood, N. D. and S. L. Welsh. 2007. New taxa of *Camissonia* (Onagraceae); *Erigeron*, *Hymenoxys*, and *Tetradymia* (Compositae); *Lepidium* and *Physaria* (Cruciferae) from Arizona, New Mexico, and Utah. *Rhodora* 109: 395–414.
- Baldwin, B. G. and B. L. Wessa. 2000. Phylogenetic placement of *Plucha* and new subtribes in *Helenieae* sensu stricto (Compositae). *Syst. Bot.* 25: 522–538.
- , ———, and J. L. Panero. 2002. Nuclear rDNA evidence for major lineages of Helenioid Heliantheae (Compositae). *Syst. Bot.* 27: 161–198.
- Barkley, T. M. 1986. *Hymenoxys*, Pp. 962–964 in *Flora of the Great Plains*, eds. T. M. Barkley, R. E. Brooks, and E. K. Schofield. Lawrence, Kansas: University of Kansas Press.
- Biernier, M. W. 1978. Flavonoids of *Plummera*: taxonomic implications at the generic level within the Gaillardiiinae. *Biochem. Syst. Ecol.* 6: 293–295.
- . 1994. Submersion of *Dugaldia* and *Plummera* in *Hymenoxys* (Asteraceae: Heliantheae: Gaillardiiinae). *Sida* 16: 1–8.
- . 2001. Taxonomy of *Hymenoxys* subgenus *Picradenia* and a conspectus of the subgenera of *Hymenoxys* (Asteraceae: Helenieae: Tetraneurinae). *Lundellia* 4: 37–63.
- and R. K. Jansen. 1998. Systematic implications of DNA restriction site variation in *Hymenoxys* and *Tetraneuris* (Asteraceae, Helenieae, Gaillardiiinae). *Lundellia* 1: 17–26.
- and B. L. Turner. 2003. Taxonomy of *Tetraneuris* (Asteraceae: Helenieae: Tetraneurinae). *Lundellia* 6: 44–96.
- and ———. 2006. *Tetraneuris*. Pp. 447–453 in *Flora of North America North of Mexico. Volume 21. Asteraceae, part 3*, eds. Flora of North America Editorial Committee. New York: Oxford University Press.
- , J. G. Díaz, B. Barba, and W. Herz. 1992. *Tetraneuris linearifolia* var. *arenicola* (Asteraceae: Heliantheae): a new variety from south Texas. *Sida* 15: 231–239.
- Cronquist, A. 1994. *Hymenoxys*. Pp. 78–86 in *Intermountain Flora. Vascular Plants of the Intermountain West, U.S.A. Volume 5. Asterales*, A. Cronquist, eds. A. H. Holmgren, N. H. Holmgren, J. L. Reveal, P. K. Holmgren. Bronx, New York: The New York Botanical Garden.
- Díaz, J. G., B. Barba, W. Herz, and M. W. Bierner. 1992. Sesquiterpene lactones and monoterpene glucopyranosides of *Tetraneuris linearifolia* var. *arenicola*. *Phytochemistry* 31: 3471–3477.
- Ferracini, V. L., F. Gao, M. W. Bierner, and T. J. Mabry. 1994. Sesquiterpene lactones and one flavonoid from *Hymenoxys scaposa* (DC.) K. Parker var. *linearis* (Nutt.) K. Parker. *Biochem. Syst. and Ecol.* 22: 111–112.
- Karis, P. O. and O. Ryding. 1994. Tribe Helenieae. Pp. 521–558 in *Asteraceae: Cladistics & Classification*, ed. K. Bremer. Portland, Oregon: Timber Press.
- Panero, J. L. 2007. Subtribe Tetraneuriinae. Pp. 404–405 in *The Families of Vascular Plants*, ed. K. Kubitzki, *Volume VIII. Flowering Plants. Eudicots: Asterales*, eds. J. W. Kadereit, C. Jeffrey. Berlin, Heidelberg: Springer-Verlag.
- Spring, O., B. Zitterell-Haid, M. W. Bierner, and T. J. Mabry. 1994. Chemistry of glandular trichomes in *Hymenoxys* and related genera. *Biochem. Syst. Ecol.* 22: 171–195.
- Welsh, S. L. 1987. Compositae. *Great Basin Naturalist Mem.* 9: 131–240.