

All quiet in the East: molecular analysis Retrieves *Chionomys layi* as part of *C. nivalis*

Authors: Mahmoudi, Ahmad, Kryštufek, Boris, Aliabadian, Mansour, Moghaddam, Faezeh Yazdani, and Darvish, Jamshid

Source: Folia Zoologica, 66(1) : 67-71

Published By: Institute of Vertebrate Biology, Czech Academy of Sciences

URL: <https://doi.org/10.25225/fozo.v66.i1.a9.2017>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

All quiet in the East: molecular analysis retrieves *Chionomys layi* as part of *C. nivalis*

Ahmad MAHMOUDI¹, Boris KRYŠTUFEK², Mansour ALIABADIAN^{1,3}, Faezeh Yazdani MOGHADDAM^{1,3} and Jamshid DARVISH^{1,4*}

¹ Department of Biology, Faculty of Sciences, Ferdowsi University of Mashhad, Mashhad, Iran; e-mail: a.mahmoudi.bio@gmail.com

² Slovenian Museum of Natural History, Prešernova 20, SI-1000 Ljubljana, Slovenia

³ Zoological Innovations Research Department, Institute of Applied Zoology, Faculty of Sciences, Ferdowsi University of Mashhad, Mashhad, Iran

⁴ Rodentology Research Department, Institute of Applied Zoology, Faculty of Sciences, Ferdowsi University of Mashhad, Mashhad, Iran; e-mail: darvishj2001@yahoo.com

Received 13 November 2016; Accepted 5 March 2017

Abstract. Snow voles from Zard Kuh in Zagros Mts., Western Iran, were identified as distinct from the European snow vole *Chionomys nivalis* and described as a species new to science under the name *C. layi* Zykov, 2004. This description was missed in subsequent years by majority of authors dealing with the genus *Chionomys*, and taxonomic identity of the new species remains uncertain. In this study we conducted the first testing ever performed on taxonomic status of *C. layi*. We sequenced cytochrome *b* (1140 bp) and nuclear growth hormone receptor *ghr* (870 bp) in topotypes of *layi* and used them, together with all available haplotypes for snow voles, in phylogenetic reconstructions. Topology of phylogenetic trees accorded published results and topotypes of *layi* nested within reference samples of *C. nivalis*. They hold a supported sister position against sequences from the geographically nearby Caucasus and Alborz. Kimura-2 parameter metrics retrieved relatively low divergence (3.2 ± 0.5) between samples from the Zagros and the Alborz + Caucasus. We conclude that *layi* is synonymous with *C. nivalis*. Our results also show that the ancient refugium for the European snow vole in the Kopet Dagh was detached from the Hyrcanian refugium in the mesic mountains to the south of the Caspian Sea.

Key words: Hyrcanian refugium, Iran, Middle East, molecular taxonomy, phylogeography, Zagros Mts.

Introduction

The snow vole genus *Chionomys* consists of three species (Kryštufek 1999, Musser & Carleton 2005) and this taxonomic view remained remarkably stable ever since proposed by Ognev (1950) and Ellerman & Morrison-Scott (1951). While phylogenetic reconstructions utilizing molecular markers confirmed such a taxonomic arrangement (Yannic et al. 2012), they also challenged the taxonomic scope of the European snow vole *C. nivalis* on the eastern edge of its range. Bannikova et al. (2013) reported a highly divergent phylogenetic lineage from Aladağ in the Taurus Mts. of Turkey, and taxonomically linked it with the name *spitzenbergerae*. Arslan et al. (2016) showed however that cytochrome *b* (*cytb*) sequences of topotypes of *spitzenbergerae* (the type locality is Maden Köy, Ulukışla, Niğde, Turkey) cluster with *C. nivalis* haplotypes from the Taurus Mts. of south Turkey. Another candidate taxonomic name which could be linked with the snow voles from Aladağ is *C.*

layi. The name was proposed for snow voles from Zard Kuh in Zagros Mts., Western Iran, which were originally classified as *C. nivalis* (Lay 1967) but were subsequently recognized as a species on its own right on the basis of morphological characteristics (Zykov 2004). The name *layi* however was missed in the influential compilation of the superfamily Muroidea by Musser & Carleton (2005) and therefore attracted very little attention. We provide in this study molecular evidence which removes ambiguities associated with the taxonomic status of *layi* by firmly nesting it in *C. nivalis*. Simultaneously we made available new sequences from north Iran which better define the geographic scope of phylogeographic lineages of *C. nivalis* in the Middle East.

Material and Methods

Samples

This study consists of seven snow voles collected in three regions of Iran (Table 1 and Fig. 1), two of

* Corresponding Author




Fig. 1. Records of snow voles *Chionomys* in Iran: 1 – Central Alborz Mts., 2 – Binaloud Mts., 3 – Zagros Mts. See Table 1 for further locality information. Range (shaded grey) is modified from Kryštufek & Amori (2008).

which were toptotypical with *C. layi*. All specimens are preserved as standard museum skins and skulls and deposited at the Museum of Vertebrate Zoology, Berkeley, U.S.A. (MVZ) and the Zoological Museum of Ferdowsi University of Mashhad, Iran (ZMFUM, Table 1). Taxonomic identification was facilitated by examination of traditional taxonomic characteristics (colour, external and cranial measurements, and enamel molar pattern) provided by Kryštufek & Vohralík (2005) and Zykov (2004). Cranial measurements were scored by Vernier calliper to the nearest 0.1 mm and the external measurements were deduced from specimen tags. *Cytb* sequences of our voles were combined with 94 sequences downloaded from GenBank

(supplementary information, Table S1) and belonging to all three currently recognized species of snow voles (Jaarola et al. 2004, Pfunder et al. 2004, Galewski et al. 2006, Bužan & Kryštufek 2008, Castiglia et al. 2009, Bannikova et al. 2010, 2013, Fink et al. 2010, Yannic et al. 2012, Barbosa et al. 2013, Arslan et al. 2016). Phylogenetic trees were rooted using three arvicolines which are closely related to *Chionomys* (Zou et al. 2008, Bannikova et al. 2009): *Blanfordimys afghanus* (GenBank Accession Numbers EF599108-9), *B. juldaschi* (EF599112-3), and *Microtus fortis* (EU126809). To exclude possibility of introgression of alien mtDNA via hybridisation (Harrison & Larson 2014), we sequenced the nuclear gene for growth hormone receptor (*ghr*) in two topotypes of *C. layi* and two individuals from Binaloud Mts. All available *ghr* sequences from GenBank were used as supplementary data (*C. nivalis*: AM392378, JN244718-9, JX440339, JX440340, KP057338; *C. gud*: JN244708-14; *C. roberti*: JN244715-17; Galewski et al. 2006, Bannikova et al. 2013, Petrova et al. 2016). *Arvicola amphibius* (ZMFUM_4397), *Blanfordimys afghanus* (ZMFUM_1426), *Microtus pennsylvanicus* (AF540633, Adkins et al. 2003) and *Ellobius talpinus* (PMS_223) were used as putative close relations to rooting phylogenetic trees.

Molecular analyses

Nucleotide sequences were edited manually in CodonCode Aligner software (CodonCode Corp.), and aligned with the Clustal W (Thompson et al. 1997) algorithm using BioEdit 7.0.5 (Hall 1999). The best fit model of DNA substitution was chosen using jModeltest 0.1.1 (Posada 2008), based on the Akaike Information

Table 1. Details of sample localities (mapped in Fig. 1) for snow voles from Iran. Geographic origin of the samples and GenBank accession numbers are provided. Museum vouchers are deposited at the Museum of Vertebrate Zoology, Berkeley, U.S.A. (indicated by asterisk *) and the Zoological Museum of Ferdowsi University of Mashhad, Iran (the rest).

No. on map	Sub species	Region/Locality	Coordinates	Specimen voucher	Accession no.	
					<i>cytb</i>	<i>ghr</i>
1	<i>C. n. trialeticus</i>	Northern Iran, Central Alborz Mts, Tar Lake	35°47'59.99" N 52°04'0.012" E	MVZ 198808*	KY761973	
2	<i>C. n. dementievi</i>	North-eastern Iran, Binaloud Mts.	36°15'42.5" N 59°07'13.5" E	ZMFUM 5215	KY761972	KY761978
				ZMFUM 5216	KY761971	KY761979
				ZMFUM 5217	KY761970	
				ZMFUM 5218	KY761969	
3	<i>C. n. layi</i>	Western Iran, Zard Kuh –Zagros Mts.	32°28'57.58" N 50°05'49.92" E	ZMFUM 5155	KY761974	KY761976
				ZMFUM KR02	KY761975	KY761977




Fig. 2. (A) Bayesian inference tree reconstructed from cytochrome *b* sequences of the genus *Chionomys* and rooted on *Microtus fortis*, *Blanfordimys afghanus* and *B. juldaschi*. (B) BI tree generated from the *ghr* sequences of the genus *Chionomys*. The numbers on the branches correspond to posterior probability values (BPP > 0.90) and bootstrap supports (BP > 70 %). The triangles represent species and lineages of snow voles which are based entirely on published sequences. For haplotype designations see Table 1 and supplementary information (Table S1).

Table 2. K2P genetic distances between the geographical lineages of the European snow vole *Chionomys nivalis*. Within lineage distances are shown in diagonal.

Lineage	1	2	3	4	5	6	7	8
1. Kopet-Dagh, Binaloud	0.5 ± 0.1							
2. Alborz, Caucasus	6.1 ± 0.9	1.7 ± 0.3						
3. Zagros	6.0 ± 0.9	3.2 ± 0.5	0.4 ± 0.2					
4. Turkey	5.7 ± 0.8	2.5 ± 0.4	3.1 ± 0.5	0.6 ± 0.2				
5. Alps, Apennines, Tatra	5.7 ± 0.7	3.9 ± 0.5	4.0 ± 0.6	3.5 ± 0.5	1.6 ± 0.2			
6. Spain	5.9 ± 0.8	4.2 ± 0.6	4.5 ± 0.7	3.6 ± 0.5	3.2 ± 0.5	1.3 ± 0.3		
7. Israel, Syria	6.1 ± 0.9	2.9 ± 0.5	2.9 ± 0.5	1.8 ± 0.4	3.3 ± 0.6	3.8 ± 0.6	0.6 ± 0.2	
8. Balkans, Carpathians	5.8 ± 0.9	4.2 ± 0.6	4.2 ± 0.5	3.5 ± 0.5	2.9 ± 0.4	3.5 ± 0.5	3.8 ± 0.6	1.5 ± 0.2

Criterion (AIC). The General Time-Reversible (GTR) model with proportion of invariant sites ($I = 0.5502$) and gamma distribution of rates across sites ($G = 1.1005$) best fitted *cytb* dataset. The best fit model for *ghr* dataset was HKY + G ($G = 0.2193$). Pairwise genetic distances, both between species and within them, were estimated with Kimura-2 parameter model (K2P) using Mega v6 (Tamura et al. 2013).

Bayesian inference (BI) and Maximum Likelihood (ML) phylogenetic trees were constructed using selected substitution model. BI analysis was conducted in MrBayes 3.1.2 (Huelsenbeck & Ronquist 2001, Ronquist & Huelsenbeck 2003), using two simultaneous analyses with four Markov Chain Monte Carlo (MCMC) models which started from random trees and were run for 4 million iterations. The trees were sampled every 1000th generation after removing the first 10 % of the trees as the burn-in stage. ML tree estimation was carried out using PAUP 4.0b10 (Swofford 2003) under 100 bootstrap pseudo-replicates with a ML heuristic tree search using 10 random additional sequence replicates. Nodal robustness for BI and ML analyses were assessed using Bayesian Posterior Probability (BPP) and Bootstrap values (BP) for BI and ML analyses, respectively. The values BPP > 0.95 and BP > 80 were assumed as good support for BI and ML analyses, respectively.

Results

Taxonomic identity

Comparison with published character states classified our specimens as snow voles (*Chionomys*). Any of them showed similarities with the Caucasian endemics *C. gud* and *C. roberti* but matched *C. nivalis-layi*. Zykov (2004) distinguished *C. layi* from *C. nivalis* by longer ears (20 mm), longer diastema (above 9.5 mm) and wider bullae (breadth over 7 mm). Morphological comparison with the data provided by Zykov (2004) showed broad similarity between the topotypes and the type series of

C. layi. Corresponding measurements in our topotypes of *layi* were (parenthesized is range for *C. nivalis* in Zikov 2004): length of ear 19 and 20 mm (12–18 mm), length of diastema 9.0 and 9.7 mm (7.5–9.3 mm), and breadth of bulla 7.0 and 7.4 mm (6.0–7.0 mm).

Molecular analyses

Altogether, five new *cytb* haplotypes were found in our material generating 74 different snow vole *cytb* haplotypes. Within the 1140 bp long sequences considered here, 193 polymorphic sites were found with a total of 267 mutations, 170 of which were parsimony informative. Our analysis of *ghr* retrieved three new sequences (one from Zard Kuh and two from Binaloud Mts.) generating 20 different *ghr* sequences. Within 870 bp long sequences, 25 and 20 sites were found to be polymorphic and parsimony informative, respectively. BI and ML trees of mitochondrial data showed similar topologies, therefore only the BI tree is shown (Fig. 2A). Branching pattern in both trees was congruent with published results (Bannikova et al. 2013, Arslan et al. 2016) therefore three species of snow voles emerged also in our reconstruction: *C. nivalis*, *C. gud*, and *C. roberti*. Sister position of *C. nivalis* against the clade of *C. gud* + *C. roberti* benefited strong BP support (100 %). Within *C. nivalis* the enigmatic sequence from Aladağ (JN244707, Bannikova et al. 2013) hold a supported (BP 95 %) basal position. The remaining haplotypes clustered into two strongly supported (BPP 1.00, BP 99 %) major lineages which contained also our new sequences from Iran. Four samples from Binaloud Mts. hold a supported (BPP 1.00, BP 99 %) sister position against the reference samples of *C. n. dementievi* from adjacent Kopet Dagħ in Turkmenistan (Turkmenian lineage). The remaining haplotypes from Europe and Asia were in a single cluster (Euro-Asiatic lineage). Firstly, two strongly supported (BPP 1.00, BP 89 %) sublineages emerged in this lineage, the European and the Asiatic. Further substructuring within the latter retrieved two supported (BPP 0.98) clusters, the Caucasian and the Anatolian. The latter contained published samples from Turkey and the Levant coast in Syria and Israel. Our samples from Alborz (= Elburz) and Zagros clustered with the Caucasian haplotypes. This cluster however further substructured (BPP 0.98) and the two sequences from the Zagros were in a sister position against samples from Alborz + Caucasus. Phylogenetic analysis of *ghr* gene showed identical tree topology (Fig. 2B). Putting aside the enigmatic haplotype from Aladağ, Turkey, K2P distances between European snow vole lineages varied between 1.8 ± 0.4 (Syria + Israel vs. Anatolia) and 5.7 ± 0.7 – 6.1 ± 0.9 (Turkmenian vs.

the remaining lineages); median K2P was 3.8 %. Within-lineage distances were mainly below 1 %; higher values (1.3 ± 0.3 – 1.7 ± 0.3) evidently point on further substructuring, particularly so in the European lineages, and within the Caucasian lineage. Samples from Zard Kuh were separated from a cluster Alborz + Caucasus by K2P distance of 3.2 ± 0.5 (Table 2).

Discussion

In this study we conducted the first testing ever performed on taxonomic status of *Chionomys layi* which was described more than a decade ago (Zikov 2004) and attracted very little attention ever since. The two haplotypes obtained from the type locality in Zard Kuh (the Zagros Mts., Western Iran) nested with reference samples of *C. nivalis* and hold a supported sister position against sequences from the geographically nearby Caucasus and Alborz. K2P metrics in our results retrieved relatively low divergence from the remaining *C. nivalis* samples. We therefore synonymize *C. layi* Zikov, 2004, with *C. nivalis* (Martins, 1842). Since the divergence between *layi* and the sequences from the Caucasus and Alborz overlaps the distances among the major lineages occupying Europe, Levant coast, Anatolia, and Western Iran, *layi* may suit as a subspecific name for European snow voles occupying the Zagros Mts. Samples from Northern Iran clustered in accordance with their geographical position, either with the Caucasian lineage (Alborz) or with the Turkmenian lineage (Binaloud). Evidently, the ancient refugium for the European snow vole in the Kopet Dagħ and the Binaloud Mts. was detached from the Hyrcanian refugium in the mesic mountains to the south of the Caspian Sea. On the other hand, snow voles from the Alborz are phylogeographically remarkably similar to their conspecifics from the Caucasus. Therefore, although the Hyrcanian refugium harbours several small mammal endemics, most notably *Microtus schelkovnikovi*, *Apodemus hyrcanicus*, and *Glis glis persicus* (Naderi et al. 2013) it also shows historical links with the Caucasian refugium. Further sampling is unavoidable to identify the actual biodiversity pattern and retrieve the underlying process in this important refugial area.

Acknowledgements

We would like to thank Chris J. Conroy for providing one sequence (Voucher number; MVZ 198808). The first author is deeply thankful to Mojtaba Karami and Mehdi Dezhman who helped to collect *C. n. layi* at the type locality. This study was funded by the grants from Ferdowsi University of Mashhad (Project numbers: 2/37624) to A. M.

Literature

- Adkins R.M., Walton A.H. & Honeycutt R.L. 2003: Higher-level systematics of rodents and divergence time estimates based on two congruent nuclear genes. *Mol. Phylogenet. Evol.* 26: 409–420.
- Arslan A., Arslan E., Mahmoudi A. et al. 2016: Taxonomic identity of *Chionomys nivalis spitzenbergerae* (Mammalia: Rodentia). *Zool. Middle East* 63: 1–7.
- Bannikova A.A., Lebedev V.S. & Golenishchev F.N. 2009: Taxonomic position of Afghan vole (Subgenus *Blanfordimys*) by the sequence of the mitochondrial *cytb* gene. *Russ. J. Genet.* 45: 91–97.
- Bannikova A.A., Lebedev V.S., Lissovsky A.A. et al. 2010: Molecular phylogeny and evolution of the Asian lineage of vole genus *Microtus* (Rodentia: Arvicolinae) inferred from mitochondrial cytochrome *b* sequence. *Biol. J. Linn. Soc.* 99: 595–613.
- Bannikova A.A., Sighazeva A.M., Malikov V.G. et al. 2013: Genetic diversity of *Chionomys* genus (Mammalia, Arvicolinae) and comparative phylogeography of snow voles. *Russ. J. Genet.* 49: 561–575.
- Barbosa S., Pauperio J., Searle J.B. & Alves P.C. 2013: Genetic identification of Iberian rodent species using both mitochondrial and nuclear loci: application to noninvasive sampling. *Mol. Ecol. Res.* 13: 43–56.
- Bužan E.V. & Kryštufek B. 2008: Phylogenetic position of *Chionomys gud* assessed from a complete cytochrome *b* gene. *Folia Zool.* 57: 274–282.
- Castiglia R., Annesi F., Kryštufek B. et al. 2009: The evolutionary history of a mammal species with a highly fragmented range: the phylogeography of the European snow vole. *J. Zool. Lond.* 279: 243–250.
- Ellerman J.R. & Morrison-Scott T.C.S. 1951: Checklist of Palaearctic and Indian mammals, 1758–1946. *Trustees of the British Museum, London*.
- Fink S., Fischer M.C., Excoffier L. & Heckel G. 2010: Genomic scans support repetitive continental colonization events during the rapid radiation of voles (Rodentia: *Microtus*): the utility of AFLPs versus mitochondrial and nuclear sequence markers. *Syst. Biol.* 59: 548–572.
- Galewski T., Tilak M.K., Sanchez S. et al. 2006: The evolutionary radiation of Arvicolinae rodents (voles and lemmings): relative contribution of nuclear and mitochondrial DNA phylogenies. *BMC Evol. Biol.* 6: 80.
- Hall T.A. 1999: BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symp. Ser.* 41: 95–98.
- Harrison R.G. & Larson E.L. 2014: Hybridization, introgression, and the nature of species boundaries. *J. Hered.* 105 (Suppl. 1): 795–809.
- Huelsenbeck J.P. & Ronquist F. 2001: MrBayes: Bayesian inference of phylogeny. *Bioinformatics* 17: 754–755.
- Jaarola M., Martinková N., Gündüz I. et al. 2004: Molecular phylogeny of the speciose vole genus *Microtus* (Arvicolinae, Rodentia) inferred from mitochondrial DNA sequences. *Mol. Phylogenet. Evol.* 33: 647–663.
- Kryštufek B. 1999: Snow voles, genus *Chionomys*, of Turkey. *Mammalia* 63: 323–339.
- Kryštufek B. & Amori G. 2008: *Chionomys nivalis*. The IUCN Red List of Threatened Species. www.iucnredlist.org
- Kryštufek B. & Vohralík V. 2005: Mammals of Turkey and Cyprus. *Annales, Koper*.
- Lay D.M. 1967: A study of the mammals of Iran: resulting from the street expedition of 1962–63 (vol. 54). *Field Museum of Natural History*.
- Musser G.G. & Carleton M.D. 2005: Superfamily Muridae. In: Wilson D.E. & Reeder D.A.M. (eds.), *Mammal species of the world. A taxonomic and geographic reference*, vol. 2nd, 3rd ed. *John Hopkins University Press, Baltimore*: 894–1531.
- Naderi G., Kaboli M., Koren T. et al. 2013: Mitochondrial evidence uncovers a refugium for the fat dormouse (*Glis glis* Linnaeus, 1766) in Hyrcanian forests of northern Iran. *Mamm. Biol.* 79: 202–207.
- Ognev S.I. 1950: Mammals of the USSR and adjacent countries: Rodents (continued). (Mammals of eastern Europe and northern Asia). *Akademiya Nauk SSSR* 7. (in Russian)
- Petrova T.V., Tesakov A.S., Kowalskaya Y.M. & Abramson N.I. 2016: Cryptic speciation in the narrow-headed vole *Lasiopodomys (Stenocranius) gregalis* (Rodentia: Cricetidae). *Zool. Scr.* 45: 618–629.
- Pfunder M., Holzgang O. & Frey J.E. 2004: Development of microarray-based diagnostics of voles and shrews for use in biodiversity monitoring studies, and evaluation of mitochondrial cytochrome oxidase I vs. cytochrome *b* as genetic markers. *Mol. Ecol.* 13: 1277–1286.
- Posada D. 2008: jModelTest: phylogenetic model averaging. *Mol. Biol. Evol.* 25: 1253–1256.
- Ronquist F. & Huelsenbeck J.P. 2003: MrBayes 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* 19: 1572–1574.
- Swofford D.L. 2003: PAUP: Phylogenetic analysis using parsimony (and other methods), version 4b10. *Sinauer Associates, Sunderland, MA*.
- Tamura K., Stecher G., Peterson D. et al. 2013: MEGA6: molecular evolutionary genetics analysis, version 6.0. *Mol. Biol. Evol.* 30: 2725–2729.
- Thompson J.D., Gibson T.J., Plewniak F. et al. 1997: The CLUSTAL – windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Res.* 25: 4876–4882.
- Yannic G., Burri R., Malikov V.G. & Vogel P. 2012: Systematics of snow voles (*Chionomys*, Arvicolinae) revisited. *Mol. Phylogenet. Evol.* 62: 806–815.
- Zou Y., Wang J.B., Gaowa H.S. et al. 2008: Isolation and genetic characterization of Hantaviruses carried by *Microtus* voles in China. *J. Med. Virol.* 80: 860–868.
- Zykov A.E. 2004: Review of snow voles (Rodentia, Arvicolinae, *Chionomys*) from Iran with description of new species from Zagros Mountains (Central Iran). *Pratsi Zoologichnogo Muzeju KNU imeni Tarasa Shevchenka* 2: 116–127. (in Russian with English summary)

Supplementary online material

Table S1. Information on cytochrome *b* sequences of three species of snow voles (*Chionomys*) retrieved from GenBank: geographic origin of the samples and GenBank accession no. are provided. Subspecies names are reported only for *C. nivalis* (http://www.ivb.cz/folia_zoologica/supplementarymaterials/mahmoudi_a_et_al_table_s1.docx).