



## **ITS sequence data confirm the presence of *Schistidium marginale* in Scandinavia and indicate connections to the Caucasus**

Authors: Kiebacher, Thomas, and Blom, Hans H.

Source: Lindbergia, 2024(1)

Published By: Dutch Bryological and Lichenological Society and Nordic Bryological Society

URL: <https://doi.org/10.25227/linbg.025153>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

## Bryological note

# ITS sequence data confirm the presence of *Schistidium marginale* in Scandinavia and indicate connections to the Caucasus

Thomas Kiebacher<sup>1</sup> and Hans H. Blom<sup>2</sup>

<sup>1</sup>Department of Botany, State Museum of Natural History Stuttgart, Stuttgart, Germany

<sup>2</sup>Norwegian Institute of Bioeconomy Research, Bergen, Norway

Correspondence: Thomas Kiebacher ([thomas.kiebacher@smns-bw.de](mailto:thomas.kiebacher@smns-bw.de))

Lindbergia

2024: e025153

doi: [10.25227/linbg.025153](https://doi.org/10.25227/linbg.025153)

Subject Editor: Kristian Hassel

Editor-in-Chief: Nils Cronberg

Accepted 19 December 2023

We document the presence of *Schistidium marginale* (Grimmiaceae) in Trollheimen, central Norway. The sequence of the nuclear ITS region aligns identically with an accession from the Caucasus, distinct from Alpine accessions. This may suggest post-glacial colonization originating from south-eastern rather than southern populations. The new locality exhibited only a few tufts and shared habitat similarities with Alpine locations, albeit with a potentially more base-rich substrate.

Keywords: biodiversity, biogeography, grimmiaceae, norway, saxicolous

Although being only recently described (Blom et al. 2016), *Schistidium marginale* H.H. Blom, Bedn.-Ochyra & Ochyra is one of the widespread and regionally common species of the genus. In Europe, it is locally frequent in the Alps (Austria, France, Italy, Switzerland) and the Caucasus mountains but also known from Albania, Macedonia, NW Russia (Murmansk Province) and Spain (Blom 1997, Köckinger et al. 2008, Ignatov et al. 2017, Hodgetts and Lockhart 2020). In Asia, it is reported from east and south Siberian mountains: Altai, Kuznetzky Alatau, Putorana Plateau and the Transbaikalian region (Ignatov et al. 2017). Notwithstanding the intensive work on *Schistidium* carried out by HHB in Scandinavia (Blom 1996, 1998), *S. marginale* has so far not been recorded there. Only recently HHB discovered small plants that showed characteristics of *S. marginale* at Mt Høghøa in the Trollheimen mountains, Trøndelag county (Fig. 1):

Norway, Trøndelag, Oppdal, Høghøa, SE slope, ca 1120 m a.s.l., 22 July 2020, leg. H.H. Blom (TRH).

At this locality *S. marginale* was only found in 2–3 small tufts at one site where it grew on low inclined (40°) phyllite ledges on a steep south-faced slope. The bed-rock is base-rich siliceous mica-shist with garnets ([https://geo.ngu.no/kart/berggrunn\\_mobil](https://geo.ngu.no/kart/berggrunn_mobil)) and *S. marginale* grew together with *Schistidium dupretii* and a yet undescribed *Schistidium* species.

© 2024 The Authors. This is an Open Access article



We aim to further explore these issues in our future studies, which will require an in depth morphological and genetic analysis of an extended set of well-developed specimens sampled across the area of occurrence of *S. marginale*. Here, we primarily aim to update our knowledge on the moss flora of the Scandinavian mountains and to rise the attention to this inconspicuous species. *Schistidium marginale* is a representative of the *Confertum* group (Blom 1996, Ignatova et al. 2010) as indicated by its small to medium size, the spinulose hair point, the partially bistratose lamina and the slightly sinuous lamina cells. It is distinctive in the prominent costa that is often subangular or even dorsally furrowed in transverse section and the very thick leaf margins, that are often four-stratous in the upper part of the leaves.

In the Alps, *S. marginale* typically colonises rather poor and acidic rocks at warm sites in the subalpine and alpine zones. This pattern aligns with the Norwegian locality, with the exception of potentially more base-rich conditions. Hence, the search for further localities in the Scandinavian mountains should focus on such habitats, especially S-facing cliffs in the alpine zone.

**Acknowledgements** – The Norwegian specimen of *S. marginale* was collected in the course of the *Alpine bryophyte* project (project no. KNR: 22-19) which is part of the Norwegian Taxonomy Initiative financed by the Norwegian Biodiversity Information Center, Trondheim.

## Data availability statement

There are no additional data for this paper.

## References

- Blom, H. H. 1996. A revision of the *Schistidium apocarpum* complex in Norway and Sweden. – Bryophyt. Bibl. 49: 1–333.
- Blom, H. H. 1997. An account on the genus *Schistidium* in two areas of the Swiss Alps. – Meylania 12: 27–29.
- Blom, H. H. 1998. 96. *Schistidium* Bruch & Schimp. in B.S.G., Bryol. Eur., 1845 nom. cons. – In: Nyholm, E. (ed.), Illustrated flora of Nordic mosses. Fasc. 4. Nordic Bryological Society, pp. 287–330.
- Blom, H. H., Bednarek-Ochyra, H. and Ochyra, R. 2016. Studies on *Schistidium* (Grimmiaceae, Bryophyta) in Europe, with particular reference to the Alps: I. A description of *S. marginale* sp. nov. – Phytotaxa 247: 210–218.
- Hodgetts, N. and Lockhart, N. 2020. Checklist and country status of European bryophytes – Update 2020. – Irish Wildl. Manuals 123: 1–214.
- Ignatova, E. A., Blom, H. H., Goryunov, D. V. and Milyutina, I. A. 2010. On the genus *Schistidium* (Grimmiaceae, Musci) in Russia. – Arctoa 19: 195–233.
- Ignatov, M. S., Ignatova, E. A., Fedosov, V. E., Ivanova, E. I., Blom, H. H., Mucoz, J., Bednarek-Ochyra, H., Afonina, O. M., Kurbatova, L. E., Czernyadjeva, I. V. and Cherdantseva, V. Ya. 2017. Moss flora of Russia. 42. Oedipodiales-Grimmiales. – KMK Scientific Press Ltd.
- Kiebach, T. 2020. *Schistidium succulentum* Ignatova & H. H. Blom (Grimmiaceae) new to Europe. – J. Bryol. 42: 277–280.
- Kiebach, T., Köckinger, H. and Blom, H. H. 2021. *Schistidium foraminis-martini* sp. nov. (Grimmiaceae), a high mountain calcicole from the European Alps molecularly related to *S. agassizii*. – Bryophyte Divers. Evol. 44: 1–11.
- Köckinger, H., Suanjak, M., Schriebl, A. and Schröck, C. 2008. Die Moose Kärntens. – Verlag des Naturwissenschaftlichen Vereins für Kärnten.
- Kyrkjeeide, M. O., Stenøien, H. K., Flatberg, K. I. and Hassel, K. 2014. Glacial refugia and post-glacial colonization patterns in European bryophytes. – Lindbergia 37: 47–59.
- Milyutina, I. A., Goryunov, D. V., Ignatov, M. S., Ignatova, E. A. and Troitsky, A. V. 2010. The phylogeny of *Schistidium* (Bryophyta, Grimmiaceae) based on the primary and secondary structure of nuclear rDNA internal transcribed spacers. – Mol. Biol. 44: 883–897.