

Evaluating Potential Factors Affecting Puma Puma concolor Abundance in the Mexican Chihuahuan Desert

Authors: Laundré, John W., Salazar, Joel Loredó, Hernández, Lucina, and López, Daniel Nuñez

Source: Wildlife Biology, 15(2) : 207-212

Published By: Nordic Board for Wildlife Research

URL: <https://doi.org/10.2981/07-077>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Evaluating potential factors affecting puma *Puma concolor* abundance in the Mexican Chihuahuan Desert

John W. Laundré, Joel Loredó Salazar, Lucina Hernández & Daniel Nuñez López

The distribution and abundance of pumas *Puma concolor* within mountain ranges of similar size in the Mexican Chihuahuan desert is known to vary. In 2001-2002, we tested 11 variables pertaining to habitat composition, prey abundance and anthropogenic factors to identify which ones might explain the difference in puma abundance between two mountain ranges (El Cuervo and Sierra Rica) of similar size. We found that shrub density (32.2 ± 1.9 (SE) vs 30.0 ± 1.7 shrubs/km²) and diversity (2.1 ± 0.1 vs 1.9 ± 0.1) did not differ between the two ranges. However, El Cuervo had significantly lower density of mule deer *Odocoileus hemionus* (158.3 ± 62.6 /km² vs 703.3 ± 296.1 /km²) and collared peccary *Tayassu tajacu* (5.0 ± 2.8 /km² vs 146.7 ± 70.1 /km²) faecal groups than Sierra Rica. Conversely, anthropogenic factors such as road density (52.4 km/100 km² vs 43.9 km/100km²), town density (25 towns/100 km² vs 6 towns/100 km²) and human density (6 individuals/100 km² vs 0.08 individuals/100 km²), were higher for El Cuervo than for Sierra Rica. We hypothesized that anthropogenic factors were the most important in explaining the difference in abundance of pumas between the two ranges. We propose that the higher number of people and accessibility to El Cuervo results in a high incidence of illegal hunting which suppresses prey and puma populations. We discuss the consequences of our results to the conservation of pumas in the Mexican Chihuahuan desert.

Key words: abundance, anthropogenic factors, Chihuahuan desert, Mexico, mountain range, puma, *Puma concolor*

John W. Laundré* & Lucina Hernández*, Instituto de Ecología, A.C., Centro Regional Durango, Km 5 Carr. a Mazatlan, C.P. 34100, Durango, Durango, Mexico. - e-mail addresses: launjohn@hotmail.com (John Laundré), lhernan1@oswego.edu (Lucina Hernández)

Joel Loredó Salazar, Instituto de Ecología, A.C. Km 2.5 Antigua Carr. A Coatepec, Xalapa, Veracruz, Mexico. - e-mail: joel.loredo@inecol.edu.mx

Daniel Nuñez López, Instituto de Ecología, A.C., Centro Regional Chihuahua, Km 33.3 Carr. Chihuahua a Ojinaga, C.P. 32900 Aldama, Chihuahua, Mexico. - e-mail: daniel.nunez@inecol.edu.mx

*Present address: Department of Biological Sciences, 133 Piez Hall, SUNY Oswego, Oswego, New York 13126, USA

Corresponding author: John W. Laundré

Received 15 October 2007, accepted 19 January 2008

Associate editor: Urs Breitenmoser

The puma was once the most widely distributed large mammalian carnivore in North America (Currier 1983). Although its range in the United States of America declined dramatically over the last two centuries, current populations, including the U.S. portion of the Chihuahuan desert, are considered viable (Harveson et al. 1999, Pittman et al. 1999, Logan & et al. 2001). However, in the Mexican

portion of the Chihuahuan Desert little is known of its current status and abundance.

The puma is the only remaining large mammalian carnivore in the Mexican Chihuahuan desert (Anderson 1972). Large carnivores are often of conservation concern because they are recognized as important components in some ecosystem functions (Boyce & Anderson 1999). Consequently, identifying

factors that affect the abundance of pumas in the Chihuahuan Desert can help in conserving pumas in this ecosystem.

To identify factors influencing puma abundance, we compared abiotic and biotic factors between two mountain ranges of similar physiognomy but known to have different relative abundances of pumas (Hernández & Laundré 2003). We assumed that factors that did not differ between the two areas could be eliminated as possible impacts on puma abundance. Factors that did differ between the areas were considered possible influences on puma abundance.

Study sites

Our two study areas were located in the northern part of the Mexican Chihuahuan Desert (Loredo-Salazar 2003; Fig. 1). The mountain range 'El Cuervo' (28°57'N-29°22'N latitude and 105°44'W-106°18'W longitude) is located 80 km northeast of the city of Chihuahua and covers approximately 619 km². The other mountain range studied was 'Sierra Rica' (28°38'N-29°29'N and 103°16'W-

104°32'W; 717 km²) located within the Area de Protección de Flora y Fauna Cañon de Santa Elena near the Mexican-United States border.

Predominant shrub species in both study areas were creosote bush *Larrea tridentata*, mesquite *Prosopis glandulosa* and ocotillo *Fouquieria splendens* (Loredo-Salazar 2003). The climate in both areas is typical of the hot desert with average maximum daily temperatures of 38-44°C and average rainfall of approximately 300 mm.

Methods

The assessment of puma abundance in El Cuervo was based on five preliminary visits which we made in the fall-winter of 1999-2000 to parts of the mountain range most likely to contain puma sign (i.e. dry river beds, rocky ledges and small reservoirs), and on interviews with local ranchers concerning possible sightings of pumas. During these visits we found no puma sign and local ranchers commented that pumas were known to occur in the area but were extremely rare. Based on these findings, and on the assumption that puma abundance is related to the amount of sign found in an area (Smallwood 1997), we classified this mountain range as an area of low puma abundance (Hernández & Laundré 2003). During six visits in the fall-winter of 1999-2000 to likely locations in Sierra Rica we found fresh puma tracks on every occasion. Local ranchers reported that pumas were relatively common; they often saw their tracks and occasionally saw individuals, and they reported that pumas were a problem regarding predation on their livestock (Bueno-Cabrera et al. 2005). Thus, we classified this mountain range as an area of high puma abundance (Hernández & Laundré 2003).

In each area we measured the abundance of two native prey species, the mule deer *Odocoileus hemionus* and collared peccary *Tayassu tajacu*, and two domestic prey, cattle *Bos taurus* and equines (horses *Equus feris* and donkeys *E. africanus*). We estimated relative prey abundance by counting the number of faecal groups found in 500 × 20 m belt transects (Gallina et al. 1991, Alvarez-Cardenas et al. 1999). As the mountain ranges were relatively large and many areas inaccessible, we randomly placed the transect lines perpendicular to and starting 20 m from existing roads that lead into the ranges. We determined the minimum number of transects needed (Krebs 1999) to be 30 in each area.




Figure 1. Location of the study areas El Cuervo and Sierra Rica in the state of Chihuahua in northern Mexico.

We divided the number of faecal groups found per species in each transect by the area of the transect ($500\text{ m} \times 20\text{ m}$) to estimate the density ($\#$ faecal groups/ km^2).

The structure of the desert plant community can affect the distribution of mule deer (Alvarez-Cardenas et al. 1999, Sánchez-Rojas & Gallina 2000), which could then affect puma abundance. Therefore, we estimated shrub density, cover, height and species diversity using the point quarter method and formulae from Brower et al. (1990), at 50-m intervals along the 30 transects used for studying faecal groups in both areas.

Human activity can influence the behaviour and distribution of pumas (Van Dyke et al. 1986, Murphy et al. 1999), thus we measured the density ($\#$ /100 km^2) of towns, humans and roads in and around the study areas. To calculate these densities we first circumscribed a 20-km buffer around each mountain range. We chose a 20-km buffer as a reasonable distance that local people might travel to visit the mountain ranges. We estimated the number of towns, the human population and kilometres of roads within the buffers from Geographic Information System (GIS) layers available from the local office of the Instituto Nacional de Estadística Geografía e Informática (INEGI). Based on the areas of the 20-km buffers, we then calculated the various densities.

The statistical design for all comparisons was the parametric group *t* or the nonparametric equivalent, Mann-Whitney Test (Zar 1999). All statistical tests were conducted using the Statistica (Statsoft 1999) program, and all GIS analyses were conducted using ARC/info[®] software. All means are \pm standard error.

Results

During our field work in 2001-2002, we found evidence of two native prey species, mule deer and the collared peccary, in our two study areas. For mule deer, the density of faecal groups was significantly higher in Sierra Rica ($703.3 \pm 296.1/\text{km}^2$) than in El Cuervo ($158.3 \pm 62.6/\text{km}^2$; $Z = 2.86$, $P < 0.01$; Fig. 2A). For collared peccary, the density of faecal droppings in Sierra Rica ($146.7 \pm 70.1/\text{km}^2$) was significantly higher than in El Cuervo ($5 \pm 2.8/\text{km}^2$; $Z = 2.43$, $P < 0.05$; see Fig. 2A). For domestic animals, we found no difference in the density of cattle faecal droppings between the two areas (Sierra Rica:




Figure 2. Density of faecal groups for deer, peccary and equines (A) and of towns ($\#$ /100 km^2), inhabitants ($\#$ /100 km^2) and roads ($\text{km}/100\text{ km}^2$) within a 20 km radius (B) of the study areas Sierra Rica (■) and El Cuervo (□). The data for cattle are not presented in (A) because there was no difference in faecal density between the two areas.

$3,716.7 \pm 615.8/\text{km}^2$ vs El Cuervo: $2,586.7 \pm 634.6/\text{km}^2$). However, we did find significantly more faecal droppings of equines (horses and donkeys) in Sierra Rica ($661.7 \pm 156/\text{km}^2$) than in El Cuervo ($3.3 \pm 2.3/\text{km}^2$; $Z = 5.95$, $P < 0.001$; see Fig. 2A).

We found no difference in shrub density ($32.2 \pm 1.9/\text{km}^2$ vs $30.0 \pm 1.7/\text{km}^2$), cover ($7.2 \pm 0.3\text{ m}^2$ vs $6.6 \pm 0.2\text{ m}^2$) or species diversity (2.1 ± 0.1 vs 1.9 ± 0.1) between Sierra Rica and El Cuervo, respectively. However, shrub height in Sierra Rica ($1.3 \pm 0.02\text{ m}$) was significantly higher than in El Cuervo ($1.2 \pm 0.02\text{ m}$; $t = 2.96$, $P < 0.01$). This difference (0.1 m), however, was likely not biologically significant.

The density of towns within the 20-km buffer of El Cuervo was > 4 times higher than in Sierra Rica ($25/100\text{ km}^2$ vs $6/100\text{ km}^2$; Fig. 2B). Human density around El Cuervo was 75 times higher than around Sierra Rica (6 individuals/100 km^2 vs 0.08 individuals/100 km^2 ; see Fig. 2B). El Cuervo had a high-

er road density than Sierra Rica (52.4 km/100 km² vs 43.9 km/100 km²; see Fig. 2B).

Discussion

In this study we attempted to determine some of the factors that might contribute to the difference in puma abundance between two mountain ranges in the Chihuahuan desert of Northern Mexico. Differences in shrub density or cover could affect relative puma abundance, either indirectly by impacting prey abundance or directly by reducing the amount of successful hunting habitat that pumas need (Laundré & Hernández 2003). However, we found no difference in the composition and structure of the vegetal communities in the two areas. Thus, at least for our two study sites, habitat differences were unlikely to be a contributing factor.

The abundance of prey will influence the abundance of the predator (Laundré et al. 2007), and we did find differences in the abundance of mule deer, collared peccary and equines between the two areas, with higher relative abundance of each in Sierra Rica. Because the higher abundance of these species coincided with higher relative abundance of pumas, we considered the differences in prey abundance to be a possible factor contributing to the differences between the two areas. Of these, the number of native prey was probably more important because predation by pumas on equines in the Chihuahuan Desert of Mexico is relatively low (Bueno-Cabrera et al. 2005). The higher abundance of equines in Santa Elena is perhaps a result of more communal ejido lands than are found in El Cuervo; residents of ejidos commonly maintain more equines than private ranchers (Bueno-Cabrera et al. 2005, J.W. Laundré, pers. obs.).

The differences in deer and peccary abundance but not in habitat composition and structure between the two areas may seem contradictory. However, this contradiction can be resolved if we consider that the level of human presence is higher in El Cuervo than in Sierra Rica. In the United States and Canada, puma abundance can be relatively high close to major metropolitan areas (Beier 1995). In these countries, the hunting of pumas and their prey (deer and elk *Cervus elaphus*) is strictly regulated, and there is a high adherence to these regulations by the general public. In contrast, in Latin America, illegal and subsistence hunting of wildlife by humans is a major problem for most wildlife species (Robin-

son & Redford 1991, Alvard et al. 1997, Merriam 1997, Chiarello 1999, Escamilla et al. 2000). This is also the case in Mexico (Leopold 1959, Ezcurra & Gallina 1981 Galindo-Leal & Weber 1998).

Because much of the prey taken by illegal and subsistence hunters overlaps with prey species taken by pumas (Redford, 1992 Carrillo et al. 2000, Escamilla et al. 2000), human hunters may reduce the prey populations needed to support viable populations of pumas. We propose that the higher number of people near the El Cuervo mountain range contributes to a higher level of illegal hunting of both pumas and their principle prey, mule deer and the peccary. Consequently, we further hypothesize that this higher level of illegal harvest may be a factor in the differences in relative abundance of pumas and their prey between the two areas.

This hypothesis has consequences for puma conservation in the Mexican Chihuahuan Desert. Firstly, isolated mountain ranges would be more important to puma conservation than ranges close to large metropolitan areas. Secondly, conservation efforts aimed at pumas, even in high human impact areas, should concentrate on intensive environmental education efforts. If this effort succeeds in raising the environmental consciousness and respect for wildlife in general, and for deer and pumas specifically, we predict that deer and, subsequently, puma numbers in these areas would rebound.

In summary, habitat factors are not the likely reason for differences in puma abundance in mountain ranges of the Mexican Chihuahuan Desert. The reason for these differences is most likely the intensity of use by humans which, we propose, has a negative impact on prey and puma abundance, primarily as a result of illegal hunting. Although our study was limited to a comparison of two mountain ranges, it is the first of its kind in the Chihuahuan Desert of Mexico, where even the most basic information on puma ecology and abundance is lacking. As such, our findings provide a valuable first step in the evaluation of possible factors influencing puma abundance in the greater portion of the largest North American desert.

Acknowledgements - our study was funded by grants from the Lincoln Park Zoo Neotropical Fund, Earthwatch Institute and Cadbury Schweppes Corporation. We thank the staff of the Área de Protección de Fauna y Flora del Cañon de Santa Elena for their cooperation and help in the field. We also thank Juan Pablo Esparza for his help in the field and Carlos A. Muñoz Robles for his help with the

GIS analyses. Lastly, we thank Harley Shaw for revising an earlier version of this manuscript.

References

- Alvard, M.S., Robinson, J.G., Redford, K.H. & Kaplan, H. 1997: The sustainability of subsistence hunting in the Neotropics. - *Conservation Biology* 11: 977-982.
- Alvarez-Cardenas, S., Gallina, S., Galina-Tessaro, P. & Dominguez-Cadena, R. 1999: Habitat availability for the mule deer (*Cervidae*) population in a relictual oak-pine forest in Baja California Sur, Mexico. - *Tropical Zoology* 12: 67-78.
- Anderson, S. 1972: Mammals of Chihuahua Taxonomy and distribution. - *Bulletin of American Museum of Natural History* 148: 149-410.
- Beier, P. 1995: Dispersal of juvenile cougars in fragmented habitat. - *Journal of Wildlife Management* 59: 228-237.
- Boyce, M.S. & Anderson, E.M. 1999: Evaluating the role of carnivores in the Greater Yellowstone Ecosystem. - In: Clark, T.W., Curlee, A.P., Minta, S.C. & Kareiva, P.M. (Eds); *Carnivores in ecosystems: The Yellowstone experience*. Yale University Press, New Haven CT, USA, pp. 265-283.
- Brower, J.E., Zar, J.H. & von Ende, C.N. 1990: Field and Laboratory Methods for General Ecology. 3rd edition. - Wm. C. Brown. Dubuque IA, USA, 237 pp.
- Bueno-Cabrera, A., Hernández-García, L., Landré, J., Contreras-Hernández, A. & Shaw, H. 2005: Cougar impact on livestock ranches in the Santa Elena Canyon, Chihuahua, Mexico. - In: Beausoleil, R.A. & Martorello, D.A. (Eds); *Proceedings of the Eighth Mountain Lion Workshop*. Olympia, Washington, USA, pp. 141-149.
- Carrillo, E., Wong, G. & Cuarón, A.D. 2000: Monitoring mammal populations in Costa Rican protected areas under different hunting restrictions. - *Conservation Biology* 14: 1580-1591.
- Chiarello, A.G. 1999: Effects of fragmentation of the Atlantic forest on mammal communities in southeastern Brazil. - *Biological Conservation* 89: 71-82.
- Currier, M.J.P. 1983: *Felis concolor*. - *The American Society of Mammalogists. Mammalian Species No.* 200: 1-7.
- Escamilla, A., Sanvicente, M., Sosa, M. & Galindo-Leal, C. 2000: Habitat mosaic, wildlife availability, and hunting in the tropical forest of Calakmul, Mexico. - *Conservation Biology* 14: 1592-1601.
- Ezcurra, E. & Gallina, S. 1981: Biology and population dynamics of white-tailed deer in northwestern Mexico. - In: Ffolliott, P.F. & Gallina, S. (Eds); *Deer biology, habitat requirements, and management in Western North America*. Instituto de Ecología, A.C. Mexico, D.F., Mexico, pp. 78-108.
- Galindo-Leal, C. & Weber, M. 1998: El venado de la Sierra Madre Occidental: Ecología, manejo y conservación. - EDICUSA-CONABIO, Mexico, D.F., Mexico, 272 pp. (In Spanish).
- Gallina, S., Galina-Tessaro, P. & Alvarez-Cárdenas, S. 1991: Mule deer density and pattern distribution in the pine-oak forest at the Sierra de la Laguna in Baja California Sur, Mexico. - *Ethology, Ecology & Evolution* 3: 27-33.
- Harveson, L.A., Route, B., Armstrong, F., Silvy, N.J. & Tewes, M.E. 1999: Trends in populations of mountain lion in Carlsbad Caverns and Guadalupe Mountains National Parks. - *The Southwestern Naturalist* 44: 490-494.
- Hernández, L. & Landré, J.W. 2003: Status of the puma in the Mexican Chihuahuan Desert. - In: Harveson, L.A., Harveson, P.M. & Adams, R.W. (Eds); *Proceedings of the 6th Mountain Lion Workshop* Austin, Texas, USA, p. 60.
- Krebs, C.J. 1999: *Ecological Methodology*. - Addison Welsey Longman Publishers, Menlow Park, California, USA, 620 pp.
- Landré, J.W. & Hernández, L. 2003: Winter hunting habitat of pumas *Puma concolor* in northwestern Utah and southern Idaho, USA. - *Wildlife Biology* 9(2): 123-129.
- Landré, J.W., Hernández, L. & Clark, S.G. 2007: Numerical and demographic responses of pumas to changes in prey abundance: Testing current predictions. - *Journal of Wildlife Management* 71: 345-355.
- Leopold, A.S. 1959. *Wildlife of Mexico; the game birds and mammals*. - University of California Press, USA, 568 pp.
- Logan, K.A. & Sweanor, L.L. 2001: *Desert Puma Evolutionary Ecology and Conservation of an Enduring Carnivore*. - Island Press, Washington, USA, 464 pp.
- Loredo Salazar, J. 2003: Evaluación de los componentes de hábitat que afectan la abundancia de pumas en el desierto Chihuahuense a través de un análisis comparativo. - M.Sc. thesis, Instituto de Ecología, A.C., Xalapa, Ver. Mexico, 98 pp. (In Spanish).
- Merriam, J.C. 1997: Community wildlife management by Mayangna Indians in the Bosawas Reserve, Nicaragua. - M.Sc. thesis, Idaho State University, Pocatello, 155 pp.
- Murphy, K.M., Ross, P.I. & Hornocker, M.G. 1999: The ecology of anthropogenic influences on cougars. - In: Clark, T.W., Curlee, A.P., Minta, S.C. & Kareiva, P.M. (Eds); *Carnivores in ecosystems: The Yellowstone experience*. Yale University Press, New Haven Connecticut, USA, pp. 77-101.
- Pittman, M.T., Guzman, G.J. & McKinney, B.P. 1999: Ecology of the mountain lion in Big Bend Ranch State Park in Trans-Pecos Texas. - Unpublished Final Report. Wildlife Division Project # 86. Texas Parks & Wildlife, Austin, TX, USA, 53 pp.
- Redford, K.H. 1992: The empty forest. - *Bioscience* 42: 412-422.

- Robinson, J.G. & Redford, K.H. 1991: Neotropical wildlife use and conservation. - University of Chicago, Chicago, USA, 520 pp.
- Sánchez-Rojas, G. & Gallina, S. 2000: Mule deer (*Odocoileus hemionus*) density in a landscape element of the Chihuahuan Desert, Mexico. - *Journal of Arid Environments* 44: 357-368.
- Smallwood, K.S. 1997: Interpreting puma (*Puma concolor*) population estimates for theory and management. - *Environmental Conservation* 24: 283-289.
- Statsoft 1999: Statistica 1999. Edition Quick References. - Stat Soft, Inc. Tulsa, Oklahoma.
- Van Dyke, F.G., Brocke, R.H., Shaw, H.G., Ackerman, B.B., Hemker, T.P. & Lindzey, F.G. 1986: Reactions of mountain lions to logging and human activity. - *Journal of Wildlife Management* 50: 95-102.
- Zar, J.H. 1999: Biostatistical Analysis. 4th edition. - Prentice-Hall, Inc. New Jersey, USA, 718 pp.