

BIAŁOWIEŻA FOREST BIRD SURVEYS: THE END OF AN ERA

Source: Acta Ornithologica, 57(1) : 1-18

Published By: Museum and Institute of Zoology, Polish Academy of Sciences

URL: <https://doi.org/10.3161/00016454AO2022.57.1.001>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Special issue: A tribute to Ludwik Tomiałoć, Wiesław Walankiewicz and Tomasz Wesołowski

Foreword

BIAŁOWIEŻA FOREST BIRD SURVEYS: THE END OF AN ERA

Among ecologists worldwide, Białowieża Primeval Forest is now one of the best-recognized windows into our evolutionary past. This is not only due to the extensive patch of a primeval forest that – against all odds – remained largely governed by natural processes, but also due to researchers that studied the unique ecosystem and published results of their studies, bringing novel findings to an international audience. In the ornithological bubble, research in Białowieża Forest was always inextricably linked with three names: Ludwik Tomiałoć, Tomasz Wesołowski, and Wiesław Walankiewicz, who studied local birds for almost 50 years since 1975. Sadly, they all passed away recently, within a single year. In recognition of

their unique contribution to ornithological science, we devoted this issue of the *Acta Ornithologica* to the papers linked with forest birds, preceded by three essays about our missed colleagues and their bird studies in Białowieża, written by their friends. We very much hope the legacy of their research will be continued for another 50 years to come at the very least. Truly effective protection of the whole Białowieża Forest (currently just one-third of the Polish part is strictly protected) using their research results would be the best tribute to their lives.

The Editors

HOW THE LONG-TERM BIRD MONITORING IN THE BIAŁOWIEŻA NATIONAL PARK WAS LAUNCHED

Wanda WESOŁOWSKA

Department of Biodiversity and Evolutionary Taxonomy, University of Wrocław, Przybyszewskiego 65, 51–148, Wrocław, POLAND

Ludwik Tomiałoć (1939–2020) was the original initiator of research on birds in the Białowieża Forest. For a long time, Ludwik was interested in the adaptation of birds to human transformed environments and conducted research on urban populations of the WoodPigeon *Columba palumbus*. While gathering information from scattered reports for the first critical summary of information about Polish birds after the Second World War (Tomiałoć 1972), he noticed that the eastern parts of the country were relatively little known, with many poorly explored areas. Eastern Poland has

not undergone such a profound anthropogenic transformation as the remaining part of the country and many only little changed natural areas have been preserved there.

During his stay in England (1973), Ludwik noticed how different the behaviour of birds in forests transformed by man is from their behaviour in better conserved forests in Poland. It was probably then, when he decided that filling the gap in the knowledge on avian communities (i.e. recognition of both composition and densities) in undisturbed forests and studying their

DOI 10.3161/00016454AO2022.57.1.001

life-histories there would be of importance. He chose the Białowieża National Park (BNP) as the site of his research. Similarly as in mid 1970's, the Białowieża Forest is the last remaining fragment of primeval lowland mixed forests that once covered our continent. Ludwik invited two younger colleagues who were just starting their scientific careers to participate in this research: Wiesław Walankiewicz (1949–2020) and Tomasz Wesołowski (1950–2021). Following a year of reconnaissance, monitoring of the breeding birds of the Białowieża Forest began in the spring of 1975. With great effort, in difficult field conditions and in unfavorable early spring weather, ten large plots (24–33 ha) were delimited. They were located in various types of forest — coniferous, ash-alder and lime-hornbeam. The names of these plots followed the circumstances of their designation or location. These common names have survived to this day, although of course in scientific publications they were later identified only with their initials. With time, the number of plots was reduced, but they were still large and represented different types of forest. An improved version of the mapping technique for censusing breeding birds has been applied for 48 years now. Tomiałojć modified Enemar's cartographic method to increase the effectiveness of a single visit to the plot. During the breeding season, ten field visits lasting several hours were carried out on each plot, including one visit at dusk. Additionally, bird nests were actively searched for.

The research was originally planned to last for five years. Their results, however, turned out to be so revealing and bringing so much new understanding of the functioning of primeval forest ecosystems that it was decided to continue them. By 2022, 48 years of uninterrupted censuses of birds in the BNP were completed (since 2020, the team continues monitoring with the mapping technique on just four plots, but other methods — i.e. point counts — are applied on all the seven plots). Of course, the composition of the team of ornithologists conducting the research has changed over time. Ludwik Tomiałojć withdrew from participating in the field work at some point (but he collaborated in data analysis until the end). Wiesław Walankiewicz compared the results obtained with the mapping method with the results from transects, a method that requires much less effort and time. Unfortunately, this valuable work has never been published. Although Wiesław no longer contributed to the monitoring, he continued to conduct other

research at BNP. Over time, the team has grown to include Dorota Czeszczewik, Marta Maziarz, Cezary Mitrus, Patryk Rowiński, Grzegorz Hebda and Grzegorz Neubauer. After Ludwik Tomiałojć's withdrawal, Tomasz Wesołowski became the leader of this group and his research on bird assemblages in the BNP continued for 47 years. If it had not been for his enormous commitment and charisma, the monitoring would have ceased but it continued even during the most difficult times. Over the years, countless students and volunteers, several dozen people in total, also participated in the field work. They all left with unforgettable experiences and learned much about both birds and the natural forest. The results of the censuses was published every five years and the article summarizing the 45-years of research appears in this issue. In addition to monitoring, each team member conducted his own research on individual bird species or on biological processes in the forest.

Years of research in the BNP have shown that primeval temperate forests are similar to their tropical counterparts. Both are characterized by an extraordinary species richness (but individual species are found in low densities) and by predation as one of the most important factors influencing these ecosystems. The results of the research in the Białowieża Forest have become famous throughout the world. As a consequence, over the years naturalists, especially ornithologists, from many countries around the world made pilgrimages to this unique forest. Bird monitoring in the BNP is one of the longest continued ornithological research projects in the world, and certainly the only one conducted on such a large scale.

During this period, over 100 articles devoted to the birds of the BNP have been published. Their co-authors were often ornithologists from other countries. A list of these works is included at the end of the Foreword. Ornithologists working in the Białowieża Forest were not only interested in birds. There were also other research topics on ecological relationships among organisms that make up and living in the forest, including phenology of foliage development, fruiting cycles of forest trees, fluctuations in the number of leaf-eating caterpillars, changes in the number of small rodents, bird parasites or the fate of nesting holes occupied by birds. This research has also been published in multiple articles (the most important of which are listed below). Some of these studies take the form of annual monitoring and provide data that is unique on a global scale.

All members of the Białowieża trio, who started their lifelong adventure there nearly half a century ago, also devoted much time and energy to defending the Białowieża Forests against intensive forest management and logging. Ornithologists fought to preserve this most valuable ecosystem in Poland and tried to ensure that the entire forest was under protection in the national park. They were deeply involved in these activities, ignoring the many difficulties that they

encountered as a result. Unfortunately, this unique Forest, included in the UNESCO World Heritage List, is still more widely appreciated in the rest of the world than in Poland. Recently, it was the EU institutions that had to take steps to prevent logging in the Białowieża Forest.

It is hard to believe, that all three initiators of the BNP bird censuses left us within a single year. This volume is devoted to their memory.

All photos from from archives of Wanda Wesołowska, authors unknown.



The team that began monitoring birds in the BNP, 3 April 1977. Left to right: Tomasz Wesołowski, Ludwik Tomiałoć, Wiesław Walankiewicz.



Ludwik Tomiałoć designing the Łęgoles plot, 1975.



Wiesław Walankiewicz next to a large oak tree, 1970s.



Ludwik Tomiałoć while working in the woods. 1970s, Łęgoles plot.



Tomasz Wesołowski on the trunk of an overturned ash, 2012.



Tomasz Wesołowski in front of the root shield of an overturned tree, 2012.



Ian Newton (left) and Tomasz Wesołowski (right) in the BNP, 2013.



Marta Maziarz at the camera trap during the Wood Warbler study, 2015.



Census team on the 40th anniversary of the monitoring, 2015. Left to right: Cezary Mitrus, Marta Cholewa, Grzegorz Hebda, Tomasz Wesołowski (holding the card), Patryk Rowiński, Marta Maziarz and Dorota Czeszczewik (crouching).



Grzegorz Neubauer with the ladder, 2016. Fot. T. Wesołowski.

BIRDS AND BEYOND — COMMEMORATING PIONEERS OF ORNITHOLOGICAL RESEARCH IN BIAŁOWIEŻA FOREST

Grzegorz MIKUSIŃSKI

Department of Ecology, Grimsö Wildlife Research Station, and School for Forest Management, Swedish University of Agricultural Sciences (SLU), SWEDEN

We are usually preoccupied with our own problems, challenges, and visions. For active scientists, these might be research projects, graduate students, securing and managing grants, publishing papers, or pursuit of an academic career. But sometimes, events in life break our self-absorption and cause us to reflect on the past, present, and future with a new eye. Occasionally, such events

come in series like the passing away of the three eminent researchers who studied birds for decades in the Białowieża Forest. All three of them, Prof. Ludwik Tomiałojć, Prof. Wiesław Walankiewicz and Prof. Tomasz Wesołowski have abruptly left us between June 2020 and July 2021, creating painful hole in Polish ornithology and in our hearts.

Ludwik, Wiesław and Tomasz influenced my life-long adventure with forest birds in diverse ways. These three scientists were of utmost importance for initiating and conducting ornithological studies in Europe's most famous forest, and I was lucky to meet all three of them on many occasions mostly through a shared research interest in forest birds and our mutual concern for the Białowieża Forest. However, my perception and knowledge of their deeds as people and scientists is probably not directly comparable since my relationships with them differed. I visited Białowieża Forest for the first time in May 1987, and at this time I already knew Tomasz quite well and even met Ludwik a few times. I believe that I met Wiesław seven years later.

What had linked three of them most? Of course, Białowieża Forest and its birds. When we review their classic (and very long) paper published in 1984 in *Acta Ornithologica* (Tomiałojć et al. 1984), we readily understand that starting systematic ornithological research in this renowned forest must have been a journey of discovery for them. The paper summed up the first results of their excellently designed bird censuses (1975–1979) in the best-preserved part of the forest. Perhaps, the most important finding was that birds in Białowieża Forest have low breeding bird densities that were tentatively explained by heavy predatory pressure. This finding, being in stark contrast to what we knew from some other European forests and woodlands, affected the direction of future studies on birds of this primeval forest. The lives of some of the species also differed in other ways, especially when compared with western Europe — notably the selection of habitat and nest sites by some species. As Tomasz nicely explained to me later, birds of Białowieża Forest were apparently unaware of the contemporary scientific knowledge of how they should “behave”. All three of them continued ecological studies of birds in Białowieża Forest for decades (see the list of their publications included at the end of the Foreword).

Tomasz and Wiesław studied mostly cavity-nesting species (tits, the nuthatch, flycatchers, woodpeckers), whereas Ludwik pursued open-nesting species (hawfinches, thrushes). Their fieldwork required innumerable nest inspections across a large range of heights — a challenge that is particularly striking in the highly diversified and unusually tall (as for the European lowlands) vegetation of Białowieża Forest. And here comes a remarkable feature shared by these three pioneers

that made me admire them a lot — all of them climbed Białowieża trees into their 60s. This dedication to science had its costs — both Tomasz and Wiesław endured injuries, some more serious than others, falling from trees. The accident in 2001 that very seriously injured Tomasz and almost led to his death (he was working alone that day) changed his perception of life. Since then, he said that he appreciated any good day without having long-term plans. Fortunately, Ludwik, who potentially faced the gravest risk due to the height of the nest of hawfinches located on average at 18 m, with some over 30 m above the ground (Tomiałojć 2005), never fell out of a tree.

Only two common peer-reviewed publications were co-authored by all three: Ludwik, Wiesław and Tomasz (Tomiałojć et al. 1977, 1984). Nevertheless, the latter paper, as mentioned above, became a classic with the highest bibliometric achievement for Ludwik and Wiesław, and, until recently, also Tomasz (almost 300 citations in Google Scholar). Ludwik and Tomasz who were employed at Wrocław University, produced some ten additional joint papers, and Tomasz and Wiesław (employed by the university located in Siedlce) co-authored three. Not surprisingly, the majority of their scientific production as senior scientists, were published within their respective research groups or as single authors.

Ludwik, born in 1939, was older than Tomasz and Wiesław and already in the middle of 1970-ties well-known to anyone in Poland interested in ornithology. Personally, I believe that the main reason was his book “Birds of Poland — the checklist and distribution” published in 1972, the first modern account of the Polish avifauna. I remember that after purchasing it as a teenager sometime around 1975, this book boosted my interest in birds enormously and sharpened my “birder’s imagination and dreams.” A most surprising achievement considering the complete lack of any illustrations except of one little map! For years, the phrase “we have to look it up in Tomiałojć” was a typical part of discussions with my fellow birders when it came to the distribution of birds. I never forget my first, very brief, personal meeting with Ludwik one early morning of March 1978 in Gdansk. It was like meeting a deity. His personal charisma, great ability to share his enthusiasm for research, and his work ethic was a mixture rarely spotted in academia. During my PhD-studies at Grimsö in the mid-1990s, he visited me in Sweden for a month or so to work undisturbed on one of his blackbird’s papers. I remember that he was

almost glued to his desk with complete focus on his work.

I really appreciated his interest in macro-ecological questions reaching far beyond Białowieża Forest. He published a major work that looked at the distributional patterns of the European avifauna as shaped by natural and anthropogenic factors (Tomiałojć 2000). He has also published several papers concerning comparisons of birds living urban and non-urban environments and was particularly interested in studies on Wood-pigeons *Columba palumbus*. Ludwik's research, conservation and wider environmental interests have left a deep mark. I am convinced that he, through stimulating the development of birding, also largely contributed to the advance of Polish "citizen science" and the advance in conservation interests in society. It has been demonstrated elsewhere that birdwatchers are much more likely than other people to engage in conservation behaviours (Cooper et al. 2015).

I had the pleasure and honour of working, travelling, and socialising with Wiesław on numerous occasions. As I recall, it started in 1994, when he guided me and a couple of friends through limited access reserves of Białowieża National Park. Kindness, great knowledge of birds and nature, and sincere attention to people are characteristics of him that I will remember the most. He was also continuously full of potential research questions and hypotheses, even if many of them never led to any formal research and publication. One of the limiting factors in his research was the massive amount of teaching duties at his university. He was especially proud of his 2002 *Acta Ornithologica* paper that identified nest predation as a limiting factor for the population size of Collared Flycatcher *Ficedula albicollis* (Walankiewicz 2002). This paper and an earlier companion paper (Walankiewicz 1991) were milestone contributions that identified the important role of predation on some birds in primeval forest conditions.

Wiesław was a very talkative person always having a story for almost anything. In addition to his impressive biological knowledge, he possessed almost encyclopaedic familiarity of history, anthropology, human foods, physical geography and more. He was kind in so many ways. I never forget Wiesław feeding Dorota Czeszczewik and I with fantastic avocados that he found in the jungle on our joint adventure in Colombia in 2017. Moreover, he almost never turned down any call for help in field work or other activities. I really

miss the guy, and I also miss Wiesław and Dorota being an almost inseparable team of workhorses in their research for over 25 years. Finally, I would like to add that he was fierce conservation advocate. On numerous occasions, Wiesław engaged himself in discussions and other actions concerning the future of Białowieża Forest. This included, in more recent years, several publications that he co-authored that go much beyond biological questions (e.g. Czeszczewik et al. 2019 or Niedziałkowski et al. 2019).

I met Tomasz in the late 1970s, when I started to interact frequently with the group of prominent bird researchers at Department of Avian Ecology in Wrocław. Tomasz, with his extreme rigor in "doing science" captivated my attention and respect from the very beginning. As a free-lance ornithologist with ambition of becoming scientist, I benefited from his generosity and help. He provided me with excellent advice in my early studies on Icterine Warbler *Hippolais icterina* and thanks to his support I got an opportunity to start my internships in Sweden in the late 1980s that eventually transformed into PhD-studies on woodpeckers. There is no doubt that Tomasz was highly respected among his colleagues in Wrocław; I remember when prof. Dyrz introduced him once for a largely "internal" audience at one of the departmental seminars as department's scientific "forefront".

Tomasz had such a deeply rooted knowledge of his field that often it was senseless to argue with him. Throughout the friendship that I enjoyed for over 30 years, I always felt the "little guy" when it came to discussing the intricacies of forest bird ecology. Interestingly, his focus on Białowieża Forest, was almost total. I asked him in 2020 why he basically only worked there and not in other interesting forest environments. His answer was simple — after 3 months of intensive fieldwork in Białowieża Forest he had no desire to work somewhere else. Tomasz was upset about the recent trend for ornithologists to publish as many papers as possible in high-ranked journals using advanced technologies and complex statistical methods but frequently based on weak data that rarely posed or answered biologically relevant questions. His advice for young students of forest ornithology was to try to think like the object of your study — a good recipe to finding new and exciting questions and hypotheses. He was particularly proud of his research on biology of species that nest in natural cavities, in contrast to knowledge derived from nest-box studies that

he largely criticized. He and his colleagues delivered several dozen papers unravelling the biology of species in primeval, naturally dynamic conditions observed in Białowieża Forest with virtually all of them based on field-data collected during many years. A few years ago, Tomasz mentioned to me that he still has some 65 papers to write. Unfortunately, 50 of them will never be written. We should also remember Tomasz' involvement in nature conservation, particularly the fight for protection of Białowieża Forest. He told me once that his involvement in conservation did not produce easily measurable outcomes and took a lot of time; if he hadn't gotten involved he probably would have produced twice as many scientific papers. However, all of us familiar with the protection of Białowieża Forest remembers the importance of the quantitative description of old stands suggested by Tomasz (later called "Wesołowski's definition") in saving logging hundreds, if not thousands of hectares of stands valuable for biodiversity.

All three pioneers of ornithological research in Białowieża Forest were initially more interested in birds of non-forest habitats. Tomasz, for example, mentioned to me that his hope was to work on the Biebrza Marches instead, and the fact that he ended up in Białowieża Forest was accidental — the project led by Ludwik simply started earlier. Without exaggeration, we may say that ornithological knowledge of forest birds in Europe would

be much more limited without their contributions. Remarkably, both Tomasz and Wiesław admitted that their interest in birds started with knowledge acquired from Zenon Lewartowski, a living legend of Polish non-professional ornithology. Zenon still roams Białowieża Forest observing birds or guiding people interested in our famous Puszcza, as we call this forest.

Ludwik, Wiesław and Tomasz put a lot of energy in fighting for the protection of Białowieża Forest and other Polish natural areas. While their scientific contributions are well-preserved in their publications, I desperately wish that their conservation engagement and occasional personal sacrifices will not be for nothing. The hope must be that the entire Białowieża Forest will become a national park and the recent decades of no progress in nature conservation in Poland will end with the creation of additional national parks. This is what our three pioneers would wholeheartedly wish for.

ACKNOWLEDGEMENTS

I wish to express my gratitude to Todd Engstrom and Rob Fuller for their comments on earlier draft of the manuscript and suggestions concerning linguistics. Thanks to Dorota Czeszczewik for the discussions and information. Finally, I thank Piotr Tryjanowski for his idea of interviewing Tomasz and Wiesław in 2020.

ALMOST HALF A CENTURY OF ORNITHOLOGY IN BIAŁOWIEŻA FOREST

Dorota CZESZCZEWIK

Siedlce University of Natural Sciences and Humanities, Siedlce, POLAND

While I was a student in Poland during the 1990s, I dreamed of working as a researcher in a forested national park. Moreover, it was a dream not only for me, but probably for the majority of young ornithologists to meet the famous ornithological figures, who were known to us only from the

professional literature. The publications of Tomiałojć (1990) and Tomiałojć et al. (1984) were already classics of Polish ornithology. I was lucky — immediately after graduation I got the opportunity to do research in the Białowieża Forest (BF). In addition, I had the pleasure to work with

Wiesław Walankiewicz, Tomasz Wesołowski and Ludwik Tomiałoć. Tomasz and Wiesław have known each other from their biology studies at the University of Poznań, and Ludwik was their senior colleague. Wiesław was now the head of the Department of Zoology at Siedlce University, where I got a job after graduation, while Tomasz was working at the University of Wrocław and became the supervisor of my doctorate a few years later. However, the official relationships were of minor importance while working together in the field as part of the BF ornithological team — everyone was equal and worked equally hard. I learned a lot from them all, and certainly many more people can say the same.

When I joined the BF ornithological team in 1994, my older colleagues and Cezary Mitrus (the youngest member of the team at this time) were celebrating the 20-year anniversary of ornithological studies in the Białowieża National Park (BNP). Tomasz usually encouraged new ornithologists that were able to recognize birds' calls and songs to join the team, but few people liked to get up long before dawn, day after day, during the long field season each spring. However, I enthusiastically accepted such an invitation as an honour, of course. Then, a few more young ornithologists joined the team: Patryk Rowiński, Marta Maziarz, Grzegorz Hebda and Grzegorz Neubauer joined the mapping team in the following years. More recently, Julia Barczyk and Marta Cholewa (after a 4-year break) joined the team as well. One of the core research activities was the monitoring of breeding birds in BNP, and the results have been presented and discussed in an extensive series of research papers (Tomiałoć et al. 1984, Tomiałoć & Wesołowski 1990, 1994, 1996, Wesołowski et al. 2002, 2006, 2010, 2015, 2022), and the monitoring has continued to the present time.

At the beginning, I learned that each study plot in the BNP had a specific name. In the papers, the names of the study plots were just abbreviations, but the full names that we used in the field were special, and always associated with a particular story. One of them (known as "MS" in papers) is called the 'Dream' plot (*Marzenie* in Polish). It got its name from the fact that in 1975 Ludwik Tomiałoć really wanted to have one plot resembling an old city park, full of large deciduous trees but virtually lacking any spruce *Picea abies* (spruce is quite ubiquitous even in deciduous stands of the BF). When he found his ideal place, he exclaimed that his "dream" had come true. This

plot is still covered with a beautiful old-growth forest with huge oaks *Quercus robur*, limes *Tilia cordata*, maples *Acer platanoides*, hornbeams *Carpinus betulus* and very few old spruces.

The name of another plot, known from papers as "K", is derived from the word *Katorga* in Polish, which broadly means 'slavery'. It was so named because of the hard working conditions there — a riparian forest with a flowing river, numerous fallen trees and inaccessible sections flooded with deep water and millions of annoying mosquitoes in the peak season. In addition, since this study plot is the largest (33 ha), we are forced to divide it into two mornings or between two people for mapping — real "katorga".

The "NW" and "NE" plots are the Western and Eastern Bears (*Niedźwiedzice* in Polish). They took their names from a forest path called the Bear Path, at which there was a site in the 1930s associated with an attempt to reintroduce Brown Bears *Ursus arctos*. The genesis of the names of the other plots is simple: "W" is the Entrance plot (*Wejściowa* in Polish), located just behind the entrance gate to the BNP, "CM" is the Central plot (*Centralna*) due to its location in the middle of the forest. The name of the plot "L" (*Łęgoles* in Polish) reflects the character of the riparian and ash-alder forest covering it.

Over several decades of research, we all spent a lot of time in Białowieża. Monitoring of breeding birds on seven study plots started around April 10 and ended on June 20, which means that during this period at least one person was mapping one of the plots every day. For many of us, being part of the ornithological research team in the BF meant over a 2-month long presence each year. If we roughly add the territory mapping effort together — the 45 years gives more than 17 thousand hours of field work. In the early years there were even more study plots than now, and fewer people.

In April, bird mapping starts relatively late in the morning because the sun rises only after 5:30 a.m. Then, there are relatively few birds, and it is nice to have the first records of newly arrived migrant bird species. However, some mornings during this period can be very cold, and the chill penetrates to the bone. Therefore, the sight of an ornithologist in a winter jacket, cap and gloves was nothing strange in early spring (see the photograph on p. 3). Until about 20 years ago, in the swampy forests during April we walked on ice, sinking into the mud, which, fortunately, was not very deep, because there was frost below. After

that, walking on swampy stands was no longer so easy, and it was rare to return from such a trip with a dry leg. For years now, however, the swampy forests have become drier and drier, and even the mosquitoes are noticeably less numerous than before. Climate change is happening in front of our eyes.

As the season progresses, dawn becomes very early (June 20 around 3:00 a.m.) and one must get up in the dark for mapping, especially if the plot is an hour's bike ride away. Getting up so early has good points, too — probably few people other than ornithologists could see so many magnificent sunrises, or observe bison *Bison bonasus*, elk *Alces alces*, wolves *Canis lupus*, raccoon dogs *Nyctereutes procyonoides*, badgers *Meles meles* or lynxes *Lynx lynx* in their natural environments. In addition, everyone has a variety of memories concerning other natural phenomena, such as the beautiful mists or morning scents.

Until 1989, our older colleagues operated under communistic regime. The life was not easy then as almost everything was missing, even the most basic things like food and clothes. Moreover, services could cling to anything. Wiesław even spent one winter in jail for his opposition activities. Certainly, it was the ornithological passion that helped them to survive. The study plots can only be reached on foot or by bike (up to approx. 8 km). Bicycles that were worn out and often broke down caused problems, e.g. sometimes we had to quickly repair the bike in the middle of the forest or continue on foot. We used to joke that an ornithologists' bike was recognizable by the fact that it had a removable chain guard, which made it easier to put on the chain when it unexpectedly fell off in the forest. But, according to the 'Białowieża philosophy', whatever was in use, no matter how, had to work. Our bikes are a special story in themselves; when I started my research, the most popular were heavy and rather clumsy "Ukraina" fixed-wheel bikes that had no derailleur. In addition to the fact that in the time of communism it was not easy to find other brands, they were actually the most durable in forest conditions. Since the 1990s more modern bicycles began to appear, and the era of old bicycles has passed. Only Tomasz was faithful to his Ukraine-bike until the end.

In addition to not having fully reliable bicycles, rubber boots also caused problems for the team. Until around a dozen years ago, rubber boots (gumboots, or wellingtons) were the most universal outdoor footwear. But as with rubber boots, so it was with bicycle tyres — they often got

punctures, and during the communist era they both were scarce goods (like many other products). That's why one colleague wore two rubber boots of different colours — it was an early sign of pro-environmental behaviour. It also happened, when walking through a swampy area, that the muddy ground often sucked up the rubber boot and the person jumped out with a sock into the mud. These days, this happens only occasionally because the forest is getting drier.

My older colleagues told me that long ago they sometimes spent the night at 'Dziedzinka', an old forester's lodge located deep in the forest, to be ready for dawn thrush mapping in the farthest study areas. This saved time and energy, because reaching the farthest study plots took a long time. On one occasion an old donkey, a resident of Dziedzinka, ate the rubber boot of one of the team, and another time a wild boar cannibalised the remains of their open can of pork luncheon meat. That must have been a painful loss in those times when it was difficult to buy anything.

Everyone remembers at least a few interesting stories from the evening mapping, done once a season in each study plot. One of them, still from the communist period, I know from Wiesław's anecdote, when he and Ludwik were riding their bikes back through Białowieża village in the evening, without lights (which was the norm then). Wiesław rode first, Ludwik a dozen metres away, it was downhill, and suddenly a policeman with a dog on a leash jumped out of the darkness and waved a flashlight, trying to stop them. Wiesław had weak brakes on his bicycle and so he sped away, while Ludwik slowed down and was accused of "not having lights". The policeman chastised him, commenting loudly: "well, a doctor with a university education running away from a policeman with a high school education?" After which, Ludwik was given a double fine — one for him and the other for his colleague who sped away. However, the consequences of this event were much more serious, because a few months later Ludwik was not allowed to go on a foreign internship, just because of this incident. In those days, such things happened, unfortunately.

Evening mapping, especially at the beginning of the season, when we had just arrived from different parts of Poland to Białowieża, were also an opportunity for kind of a social mini-gathering. For example, we always went to the 'L' plot as a small group, and before we dispersed on the study plot we sat on a fallen log talking about plans for the coming season or mentioning various events. After we counted the birds, we again

sat briefly on the log and shared our thoughts and observations. Also unforgettable were expeditions to the farthest study plots (NE, NW and MS) with the whole team. We drove by car to Dziedzinka, and then walked a few kilometres to the plots. While walking back, we sometimes heard wolves howling or other exciting sounds of the wilderness. Sometimes we walked in complete darkness and other times a beautiful moon illuminated our route.

None of our home universities has ever had a field station in the BF, so we always had to find some accommodation before the season. I know from stories that in the early years it used to be a youth hostel or an old museum building. Since the late 1980s, the 'Hunting House', located in the Białowieża Palace Park, served as a base for ornithologists for many years. It was the best location, as the route to the BNP gate took only a few minutes. In 2014, the Hunting House was closed, and the ornithologists dispersed to various locations — the Geobotanical Station of Warsaw University, the Białowieża village Youth Hostel, the 'House under the Stork' guesthouse, and several other accommodations. This was a pity, because living together had many advantages, especially since we spent many weeks away from our homes and were all friends with each other.

Apart birds censuses in BNP, there was a parallel study on the biology of tits *Paridae* led by Tomasz (the 'Wrocław team'), and of the Collared Flycatcher, initiated and led by Wiesław in 1988. For several years Wiesław collected the data by himself, observing birds and climbing trees to check nesting holes. Then Cezary Mitrus joined him in 1992, and I joined two years later. Over the years, many people passed through our team, which we call the 'Siedlce team'. These included our masters and doctorate students, usually working for 1–3 spring seasons. Several years of research that culminated in PhDs were also spent in Siedlce team by Tomasz Stański, Anna Kapusta, Grzegorz Bednarczyk and Mustapha Sahel. For the past three seasons, our Siedlce team has included Oliwia Karpińska, who studies the niche selection of birds in BNP, and by Kasia Kamionka-Kanclerska, who studies with tree-related microhabitats (TreMs).

Working on the flycatchers has consumed a lot of time. Starting around April 20, almost every day one must go around the plot in search of the nest-holes, which then have to be inspected repeatedly. At the beginning Wiesław climbed the trees himself to look into the holes with the help

of a small mirror and a torch. Then we bought the first ladder, which we could use to check many holes (an average height of flycatcher's nesting hole is 8.5 m), but many of the higher holes, located up to 17 m, were still checked by tree-climbing. Wiesław was very efficient in climbing trees, but the best climber in BF was Arek Szymura, who helped us more than once, especially when it came to the most difficult trees. Arek, having worked closely with the ornithologists, still visited us occasionally, and coloured our free time with interesting stories. In the past he has also been involved in other ornithological research with Ludwik, Tomasz and Wiesław.

Then a specialized telescopic device with a camera to check the holes appeared on the market, but its price was prohibitive. Once we managed to raise the funds, it turned out that the device was no longer produced. That's when Wiesław assembled his own device at little cost from a laptop, telescopic rod, a webcam, cables, LEDs, a spruce stick, and adhesive tape. We simply called it a fishing rod and it was used very successfully. The device had plenty of advantages, being cheap, lightweight, safe, with long range and relatively easy to repair, even in the forest. It was possible to check most holes with it. With the help of this device, we also conducted surveys of nest-hole resources in the forests of Nepal (Bhusal et al. 2016, Baral et al. 2018), Sweden and the US.

Wiesław used to say that he did not want to be a researcher who was identified with just one species of bird (although he produced several papers on the Collared Flycatcher, to mention just Walankiewicz 1991, 2002a, 2002b, Walankiewicz et al. 1997, 2007), and that he wanted to get out of the "ornithological ghetto". Therefore, our team expanded its interests to a variety of topics, including woodpeckers (e.g. Czeszczewik 2009, Walankiewicz et al. 2011, Czeszczewik et al. 2013, 2007), tree stand dynamics (e.g. Walankiewicz 2002c, Czeszczewik et al. 2022), tree cavity resources (Walankiewicz et al. 2014), mammals (Czeszczewik et al. 2008, Zub et al. 2014) and natural resource evaluation (Czeszczewik et al. 2019).

Sometimes the ideas for future work arose in quite unusual circumstances. In 2017, Wiesław, Grzegorz Mikusiński and I were returning from the Congress for Conservation Biology in Cartagena, Colombia, and on the plane we came up with the idea for a paper on the threats posed by logging in the BF as part of the fight against the bark beetle (Mikusiński et al. 2018). This was not a strictly ornithological paper, but the protection of

the Forest means protection for bird habitats, too. In contrast, another paper that focused on the relationships between climate, predation, rodents, birds and insect cycles (Czeszczewik et al. 2020) took at least several years to mature from an idea to implementation, but the data series has grown because of it.

Wiesław and Tomasz were very much involved with the BF until the end of their lives and doubtless still had plans to spend many more seasons there. Ludwik withdrew from Białowieża field-research around 2000, but visited the Forest many more times and published several papers based on data he gathered in BNP (e.g. Tomiałojć 2012, Tomiałojć & Wesołowski 2005, Tomiałojć & Neubauer 2017). Ludwik last visited the forest in the summer of 2019, where we celebrated his 80th birthday with a few months delay. This was his farewell meeting with the BF.

All three great ornithologists left us suddenly, in order of seniority, within the space of just 13 months. Wiesław, after his last season in Białowieża, still came back several times in the summer and autumn of 2020, as if he didn't want to be detached from it. And then suddenly he passed away, in that terrible Covid year. And then Tomasz also died completely unexpectedly, soon after a season spent in his beloved Białowieża. Maybe they have already started new research in some undiscovered Primeval Forest of another world. After their passing in 2020–2021, there was an unanticipated exchange of generations.

In total, the intensive Białowieża Forest ornithological research has spanned 48 breeding seasons. I hope to celebrate half a century of Białowieża Forest ornithology two years' time. Unfortunately, this will be without its founding fathers, but their legacy will continue.

ACKNOWLEDGEMENTS

Many thanks to Grzegorz Mikusiński and Marta Maziarz for their comments on an early version of the manuscript, and many thanks to Richard Broughton for his language proofreading and also for his comments on the manuscript.

REFERENCES

- Baral R., Czeszczewik D., Walankiewicz W., Churski M., Bhusal P., Thapa T. B., Mikusiński G. 2018. Characteristic of tree cavities in sal *Shorea robusta* forest, Nepal. J. For. Res. 23: 214–220.
- Bhusal P., Czeszczewik D., Walankiewicz W., Churski M., Baral R., Lamichhane B. R., Mikusiński G. 2016. Availability of tree cavities in a sal forest of Nepal. iForest 9: 217–225.
- Cooper C., Larson L., Dayer A., Stedman R., Decker D. 2015. Are wildlife recreationists conservationists? Linking hunting, birdwatching, and pro-environmental behaviour. J. Wildl. Manage. 79: 446–457.
- Czeszczewik D. 2009. Foraging behaviour of White-backed Woodpeckers *Dendrocopos leucotos* in a primeval forest (Białowieża National Park, NE Poland): Dependence on habitat resources and season. Acta Ornithol. 44: 109–118.
- Czeszczewik D., Czortek P., Jaroszewicz B., Zub K., Rowiński P., Walankiewicz W. 2020. Climate change has cascading effects on tree masting and the breeding performance of a forest songbird in a primeval forest. Sci. Total Environ. 747: 142084.
- Czeszczewik D., Ginter A., Mikusiński G., Pawłowska A., Kałuża H., Smithers R. J., Walankiewicz W. 2019. Bird-watching, logging and the local economy in the Białowieża Forest, Poland. Biodivers. Conserv. 28: 2967–2975.
- Czeszczewik D., Kosiński Z., Kubicka K., Puła W., Walankiewicz W. 2022. Mortality and regeneration of elms in primeval stands of Białowieża National Park, Poland. For. Pathol. 52: e12731.
- Czeszczewik D., Walankiewicz W., Mitrus C., Tumił T., Stański T., Sahel M., Bednarczyk G. 2013. Importance of dead wood resources for woodpeckers in coniferous stands of the Białowieża Forest. Bird Conserv. Int. 23: 414–425.
- Czeszczewik D., Walankiewicz W., Stańska M. 2008. Small mammals in nests of cavity nesting birds: Why should ornithologists study rodents? Can. J. Zool. 86: 286–293.
- Mikusiński G., Bubnicki J. W., Churski M., Czeszczewik D., Walankiewicz W., Kuijper D. 2018. Is the impact of loggings in the last primeval lowland forest in Europe underestimated? The conservation issues of Białowieża Forest. Biol. Conserv. 227: 266–274.
- Niedziałkowski K., Jaroszewicz B., Kowalczyk R., Kuijper D., Mikusiński G., Selva N., Walankiewicz W., Wesołowski T. 2019. Effective mitigation of conservation conflicts requires more than participatory governance — Response to Kuboń and colleagues. Conserv. Biol. 33: 962–965.
- Tomiałojć L. 1972. The birds of Poland — checklist and distribution. PWN, Warszawa.
- Tomiałojć L. 1990. The birds of Poland — their distribution and abundance. PWN, Warszawa.
- Tomiałojć L. 2000. An east-west gradient in the breeding distribution and species richness of the European woodland avifauna. Acta Ornithol. 35: 3–17.
- Tomiałojć L. 2012. Reproduction and population dynamics of Hawfinches *Coccothraustes coccothraustes* in the primeval forest of Białowieża National Park (NE Poland) 47: 63–78.
- Tomiałojć L., Neubauer G. 2017. Song Thrush *Turdus philomelos* and Hawfinch *Coccothraustes coccothraustes* exhibit non-random nest orientation in dense temperate forest. Acta Ornithol. 52: 209–220.
- Tomiałojć L., Wesołowski T. 1990. Bird communities of the primeval temperate forest of Białowieża, Poland. In: Keast A. (ed.). Biogeography and ecology of forest bird communities. Hague, pp. 141–165.
- Tomiałojć L., Wesołowski T. 1994. Die Stabilität der Vogelgemeinschaft in einem Urwald der gemäßigten Zone: Ergebnisse einer 15jährigen Studie aus dem Nationalpark von Białowieża (Polen). Ornithol. Beobachter 91: 73–110.
- Tomiałojć L., Wesołowski T. 1996. Structure of a primeval forest bird community during 1970s and 1990s (Białowieża National Park, Poland). Acta Ornithol. 31: 133–154.

- Tomiałojć L., Wesołowski T. 2005. The avifauna of Białowieża Forest: A window into the past. *British Birds* 98: 174–193.
- Tomiałojć L., Wesołowski T., Walankiewicz W. 1984. Breeding bird community of a primeval temperate forest (Białowieża National Park, Poland). *Acta Ornithol.* 20: 241–310.
- Walankiewicz W. 1991. Do secondary cavity-nesting birds suffer more from competition for cavities or from predation in a primeval deciduous forest? *Nat. Areas J.* 11: 203–211.
- Walankiewicz W. 2002a. Breeding losses in the Collared Flycatcher *Ficedula albicollis* caused by nest predators in the Białowieża National Park (Poland). *Acta Ornithol.* 37: 21–26.
- Walankiewicz W. 2002b. Nest predation as a limiting factor to the breeding population size of the Collared Flycatcher *Ficedula albicollis* in Białowieża National Park (NE Poland). *Acta Ornithol.* 37: 91–106.
- Walankiewicz W. 2002c. The number and composition of snags in the pine-spruce stands of the Białowieża National Park. In: *Proceedings of the Symposium on the Ecology and Management of Dead Wood in Western Forests: November 2–4, 1999, Reno, Nevada* / William F. Laudenslayer, Jr. [et al.], Albany, U.S. Dept. of Agriculture Forest Service, Pacific Southwest Research Station, pp. 489–500.
- Walankiewicz W., Czeszczewik D., Mitrus C. 2007. Natural nest sites of the Collared Flycatcher *Ficedula albicollis* in lime-hornbeam-oak stands of a primeval forest. *Ornis Fennica* 84: 155–162.
- Walankiewicz W., Czeszczewik D., Mitrus C., Szymura A. 1997. How the territory mapping technique reflects yearly fluctuations in the Collared Flycatcher *Ficedula albicollis* numbers? *Acta Ornithol.* 32: 201–207.
- Walankiewicz W., Czeszczewik D., Stański T., Sahel M., Ruczyński I. 2014. Tree cavity resources in spruce-pine managed and protected stands of the Białowieża Forest, Poland. *Nat. Areas J.* 34: 423–428.
- Walankiewicz W., Czeszczewik D., Tumił T., Stański T. 2011. Woodpeckers abundance in the Białowieża Forest — a comparison between deciduous, strictly protected and managed stands. *Ornis Polonica* 52: 161–168.
- Wesołowski T., Czeszczewik D., Hebda G., Maziarz M., Mitrus C., Rowiński P. 2015. 40 years of breeding bird community dynamics in a primeval temperate forest (Białowieża National Park, Poland). *Acta Ornithol.* 50: 95–120.
- Wesołowski T., Czeszczewik D., Hebda G., Maziarz M., Mitrus C., Rowiński P., Neubauer G. 2022. 45 years of bird censuses in a primeval temperate forest (Białowieża National Park, Poland). *Acta Ornithol.*
- Wesołowski T., Czeszczewik D., Hebda G., Maziarz M., Mitrus C., Rowiński P., Neubauer G. 2022. Long-term changes in breeding bird community in the primeval temperate forest: 45 years of censuses in Białowieża National Park (Poland). *Acta Ornithol.* 57: 71–100.
- Wesołowski T., Mitrus C., Czeszczewik D., Rowiński P. 2010. Breeding bird dynamics in a primeval temperate forest over 35 years: variation and stability in a changing world. *Acta Ornithol.* 45: 209–232.
- Wesołowski T., Rowiński P., Mitrus C., Czeszczewik D. 2006. Breeding bird community of a primeval temperate forest (Białowieża National Park, Poland) at the beginning of the 21st century. *Acta Ornithol.* 41: 55–70.
- Wesołowski T., Tomiałojć L., Mitrus C., Rowiński P., Czeszczewik D. 2002. Breeding bird community of a primeval temperate forest (Białowieża National Park, Poland) at the end of XXth century. *Acta Ornithol.* 37: 27–45.
- Zub K., Czeszczewik D., Ruczyński I., Kapusta A., Walankiewicz W. 2017. Silence is not golden: the hissing calls of tits affect the behaviour of a nest predator. *Behav. Ecol. Sociobiol.* 71: 1–7.

LIST OF MAJOR PUBLICATIONS DEVOTED TO THE BNP BY
LUDWIK TOMIAŁOJC, WIESŁAW WALANKIEWICZ
AND TOMASZ WESOŁOWSKI

(in chronological order)

- Tomiałoć L., Walankiewicz W., Wesołowski T. 1977. Methods and preliminary results of the bird census work in primeval forest of Białowieża National Park. *Pol. Ecol. Stud.* 3 (4): 215–223.
- Wesołowski T. 1981. Population restoration after removal of wrens (*Troglodytes troglodytes*) breeding in primeval forest. *J. Anim. Ecol.* 50: 809–814.
- Wesołowski T. 1983. The breeding ecology and behaviour of Wrens *Troglodytes troglodytes* living under primeval and secondary conditions. *Ibis* 125: 499–515.
- Tomiałoć L., Wesołowski T., Walankiewicz W. 1984. Breeding bird community of a primeval temperate forest (Białowieża National Park, Poland). *Acta Ornithol.* 20: 241–310.
- Wesołowski T. 1985. The breeding ecology of the Wood Warbler *Phylloscopus sibilatrix* in primeval forest. *Ornis Scand.* 16: 49–60.
- Wesołowski T., Tomiałoć L. 1986. The breeding ecology of woodpeckers in a temperate primeval forest — preliminary data. *Acta Ornithol.* 22: 1–22.
- Wesołowski T. 1987. Polygyny in three temperate forest Passerines (with a critical reevaluation of hypotheses for the evolution of polygyny). *Acta Ornithol.* 23: 273–302.
- Wesołowski T., Tomiałoć L., Stawarczyk T. 1987. Why low numbers of *Parus major* in Białowieża Forest — removal experiments. *Acta Ornithol.* 23: 303–316.
- Piotrowska M., Wesołowski T. 1989. The breeding ecology and behaviour of the Chiffchaff *Phylloscopus collybita* in primeval and managed stands of Białowieża Forest (Poland). *Acta Ornithol.* 25: 25–76.
- Wesołowski T. 1989. Nest-sites of hole-nesters in a primeval temperate forest (Białowieża National Park, Poland). *Acta Ornithol.* 25: 321–351.
- Tomiałoć L., Wesołowski T. 1990. Bird communities of the primeval temperate forest of Białowieża, Poland. In: Keast A., Blondel J., Helle P., Kikkawa J., Recher H. W., Holmes R. T. (eds). *Biogeography and ecology of forest bird communities*. SPB Academic Publ. bv, The Hague, pp. 141–165.
- Walankiewicz W. 1991. Do secondary cavity-nesting birds suffer more from competition for cavities or from predation in a primeval deciduous forest? *Nat. Areas J.* 11: 203–212.
- Wesołowski T., Stawarczyk T. 1991. Survival and population dynamics of Nuthatches *Sitta europaea* breeding in natural cavities in a primeval temperate forest. *Ornis Scand.* 22: 143–154.
- Tomiałoć L. 1991. Characteristics of old growth in the Białowieża Forest, Poland. *Nat. Areas J.* 11: 7–18.
- Tomiałoć L. 1993. Breeding ecology of the Blackbird *Turdus merula* studied in the primeval forest of Białowieża (Poland). P.1. Breeding numbers, distribution and nest sites. *Acta Ornithol.* 27: 131–157.
- Tomiałoć L., Wesołowski T. 1994. Die Stabilität der Vogelgemeinschaft in einem Urwald der gemäßigten Zone: Ergebnisse einer 15jährigen Studie aus dem Nationalpark von Białowieża (Polen). *Ornithologische Beobachter* 91: 73–110.
- Wesołowski T. 1994. Variation in the numbers of resident birds in a primeval temperate forest: Are winter weather, seed crop, caterpillars and interspecific competition involved? In: Hagemijer E. J. M., Verstrael T. J. (eds). *Bird numbers 1992*. Statistics Netherlands, Heerlen, pp. 203–211.
- Wesołowski T., Tomiałoć L. 1995. Ornithologische Untersuchungen im Urwald von Białowieża — eine Übersicht. *Ornithologische Beobachter* 92: 111–146.
- Wesołowski T. 1995. Value of Białowieża Forest for the conservation of white-backed woodpecker (*Dendrocopos leucotos*) in Poland. *Biol. Conserv.* 71: 69–75.
- Wesołowski T. 1995. The loss of avian cavities by injury compartmentalization in a primeval European forest. *Condor* 97: 256–257.
- Wesołowski T. 1995. Ecology and behaviour of White-backed Woodpecker (*Dendrocopos leucotos*) in a primeval temperate forest (Białowieża National Park, Poland). *Die Vogelwarte* 38: 61–75.
- Wesołowski T. 1995. Birds from a primeval temperate forest hardly use feeders in winter. *Ornis Fennica* 72: 132–134.

- Tomiałojć L. 1995. Breeding ecology of the Blackbird *Turdus merula* studied in the primeval forest of Białowieża (Poland). P.2. Reproduction and mortality. *Acta Ornithol.* 29: 101–121.
- Tomiałojć L. 1995. The birds of the Białowieża Forest — additional data and summary. *Acta zool. crac.* 38: 363–397.
- Mitrus C., Walankiewicz W., Czeszczewik D., Jabłoński P. M. 1996. Age and arrival date of Collared Flycatcher *Ficedula albicollis* males do not influence quality of natural cavities used. *Acta Ornithol.* 31: 101–106.
- Wesołowski T. 1996. Natural nest sites of Marsh Tits *Parus palustris* in a primeval forest (Białowieża National Park, Poland). *Die Vogelwarte* 38: 235–249.
- Tomiałojć L., Wesołowski T. 1996. Structure of a primeval forest bird community during 1970s and 1990s (Białowieża National Park, Poland). *Acta Ornithol.* 31: 133–154.
- Wesołowski T., Tomiałojć L. 1997. Breeding bird dynamics in a primeval temperate forest: Long-term trends in Białowieża National Park (Poland). *Ecography* 20: 432–453.
- Walankiewicz W., Czeszczewik D., Mitrus C., Szymura A. 1997. How the territory mapping technique reflects yearly fluctuations in the Collared Flycatcher *Ficedula albicollis* numbers? *Acta Ornithol.* 32: 201–207.
- Walankiewicz W., Mitrus C. 1997. How nest-box data have led to erroneous generalizations: the case of the competition between Great Tit *Parus major* and *Ficedula* flycatchers. *Acta Ornithol.* 32: 209–212.
- Walankiewicz W., Mitrus C., Czeszczewik D., Jabłoński P. M. 1997. Is the Pied Flycatcher *Ficedula hypoleuca* overcompeted by the Collared Flycatcher *Ficedula albicollis* in the natural forest of Białowieża? *Acta Ornithol.* 32: 213–217.
- Wesołowski T. 1998. Timing and synchronisation of breeding in a Marsh Tit *Parus palustris* population from a primeval forest. *Ardea* 86: 89–100.
- Wesołowski T. 1999. Marsh Tits (*Parus palustris*) are not excavators. *Ibis* 141: 149.
- Czeszczewik D., Walankiewicz W. 1999. Nest-cavity inspections by male Pied Flycatchers *Ficedula hypoleuca* under natural conditions in Białowieża National Park. *Vogelwelt* 120: 367–369.
- Czeszczewik D., Walankiewicz W., Mitrus C., Nowakowski W. 1999. Nest-box data of Pied Flycatcher *Ficedula hypoleuca* may lead to erroneous generalizations. *Vogelwelt* 120: 361–365.
- Wesołowski T. 2000. What happens to old nests in natural cavities? *Auk* 117: 498–500.
- Wesołowski T. 2000. Time saving mechanisms in the reproduction of Marsh Tits *Parus palustris* J. *Ornithol.* 141: 309–318.
- Wesołowski T., Stańska M. 2001. High ectoparasite loads in hole nesting birds — a nest box bias? *J. Avian Biol.* 32: 281–285.
- Wesołowski T. 2001. Ground checks — an efficient and reliable method to monitor holes' fate. *Ornis Fennica* 78: 193–197.
- Wesołowski T. 2001. Host-parasite interactions in natural holes: marsh tits (*Parus palustris*) and blow flies (*Protophormia falcipennis*). *J. Zool.* 255: 495–503.
- Pagenkopf K., Wesołowski T. 2002. Do male nuthatches *Sitta europaea* guard their mates? *J. Ornithol.* 143: 145–154.
- Wesołowski T., Tomiałojć L., Mitrus C., Rowiński P., Czeszczewik D. 2002. Breeding bird community of a primeval temperate forest (Białowieża National Park, Poland) at the end of XXth century. *Acta Ornithol.* 37: 27–45.
- Wesołowski T. 2002. Antipredator adaptations in nesting marsh tits *Parus palustris* — the role of nest site security. *Ibis* 144: 593–601.
- Walankiewicz W. 2002. Breeding losses in the Collared Flycatcher *Ficedula albicollis* caused by nest predators in the Białowieża National Park (Poland). *Acta Ornithol.* 37: 21–26.
- Walankiewicz W. 2002. Nest predation as a limiting factor to the breeding population size of the Collared Flycatcher *Ficedula albicollis* in the Białowieża National Park. *Acta Ornithol.* 37: 73–89.
- Wesołowski T., Czeszczewik D., Rowiński P., Walankiewicz W. 2002. Nest soaking in natural holes — a serious cause of breeding failure? *Ornis Fennica* 79: 132–138.
- Walankiewicz W., Czeszczewik D., Mitrus C., Bida E. 2002. [Snag importance for woodpeckers in deciduous stands of the Białowieża Forest]. *Not. Orn.* 43: 61–71.
- Wesołowski T. 2003. Clutch size and breeding performance of Marsh Tits *Parus palustris* in relation to hole size in a primeval forest. *Acta Ornithol.* 38: 65–72.
- Czeszczewik D., Mitrus C., Walankiewicz W. 2003. A hole or nest box: what is better for flycatcher? *Die Vogelwarte* 42: 59.
- Walankiewicz W., Czeszczewik D. 2003. Forest rodent cycles and fluctuations of the collared flycatcher *Ficedula albicollis*. *Die Vogelwarte* 42: 62.

- Czeszczewik D., Walankiewicz W. 2003. Natural nest sites of the Pied Flycatcher *Ficedula hypoleuca* in a primeval forest. *Ardea* 91: 221–230.
- Wesołowski T., Czeszczewik D., Mitrus C., Rowiński P. 2003. [Birds of the Białowieża National Park]. *Not. Orn.* 44: 1–31.
- Wesołowski T. 2003. Bird community dynamics in a primeval forest — is interspecific competition important? *Ornis Hungarica* 12–13: 51–62.
- Wesołowski T., Rowiński P. 2004. Breeding behaviour of the Nuthatch *Sitta europaea* in relation to natural hole attributes in a primeval forest. *Bird Study* 51: 143–155.
- Tomiałojć L., Wesołowski T. 2004. Diversity of the Białowieża Forest avifauna in space and time. *J. Ornithol.* 145: 81–92.
- Tomiałojć L. 2004. Accuracy of the mapping technique for a dense breeding population of the Hawfinch *Coccothraustes coccothraustes* in a deciduous forest. *Acta Ornithol.* 39: 67–74.
- Wesołowski T. 2005. Virtual conservation: how the European Union is turning a blind eye on its vanishing primeval forests. *Conserv. Biol.* 19: 1349–1358.
- Tomiałojć L., Wesołowski T. 2005. The avifauna of the Białowieża Forest: a window into the past. *Brit. Birds* 98: 174–193.
- Wesołowski T., Czeszczewik D., Rowiński P. 2005. Effects of forest management on Three-toed Woodpecker *Picoides tridactylus* distribution in the Białowieża Forest (E Poland): conservation implications. *Acta Ornithol.* 40: 53–60.
- Wesołowski T., Tomiałojć L. 2005. Nest sites, nest predation, and productivity of avian broods in a primeval temperate forest: Do the generalisations hold? *J. Avian Biol.* 36: 361–367.
- Tomiałojć L. 2005. Distribution, breeding density and nest sites of Hawfinches *Coccothraustes coccothraustes* in the primeval forest of Białowieża National Park. *Acta Ornithol.* 40: 127–138.
- Wesołowski T., Rowiński P. 2006. Is there a cost of reproduction for Marsh Tit *Parus palustris* in a primeval forest? *Ibis* 148: 126–132.
- Wesołowski T., Rowiński P. 2006. Tree defoliation by winter moth *Operophtera brumata* L. during an outbreak affected by structure of forest landscape. *For. Ecol. Manage.* 221: 299–305.
- Wesołowski T., Rowiński P. 2006. Timing of bud burst and tree-leaf development in a multispecies temperate forest. *For. Ecol. Manage.* 237: 387–393.
- Wesołowski T., Rowiński P., Mitrus C., Czeszczewik D. 2006. Breeding bird community of a primeval temperate forest (Białowieża National Park, Poland) at the beginning of the 21st century. *Acta Ornithol.* 41: 55–70.
- Wesołowski T. 2006. Nest-site re-use: Marsh Tit *Poecile palustris* decisions in a primeval forest. *Bird Study* 53: 199–204.
- Czeszczewik D., Walankiewicz W. 2006. Logging and distribution of the White backed Woodpecker *Dendrocopos leucotos* in the Białowieża Forest. *Ann. Zool. Fenn.* 43: 221–227.
- Walankiewicz W., Czeszczewik D. 2006. Woodpecker holes and nest-boxes as ecological traps for *Ficedula* flycatchers. *J. Ornithol.* 147 (Suppl. 1): 269.
- Czeszczewik D., Walankiewicz W. 2006. Foraging behavior of the White-backed Woodpecker during spring and winter in a primeval forest. *J. Ornithol.* 147 (Suppl. 1): 152–153.
- Mitrus C., Walankiewicz W., Czeszczewik D. 2007. Frequency of nest-hole occupation and breeding success of Collared Flycatchers *Ficedula albicollis*. *Ibis* 149: 414–418.
- Walankiewicz W., Czeszczewik D., Mitrus C. 2007. Natural nest sites of the Collared Flycatcher *Ficedula albicollis* in lime-hornbeam-oak stands of a primeval forest. *Ornis Fennica* 84: 155–162.
- Wesołowski T. 2007. Primeval conditions — what can we learn from them? *Ibis* 149 (Suppl. 2): 64–77.
- Wesołowski T. 2007. Lessons from long-term hole-nester studies in a primeval temperate forest. *J. Ornithol.* 148 (Suppl. 2): S 395–S405.
- Wesołowski T., Rowiński P. 2008. Late leaf development in pedunculate oaks *Quercus robur* L.: An anti-herbivore defence? *Scand. J. Forest Res.* 23: 386–394.
- Czeszczewik D., Walankiewicz W., Stańska M. 2008. Small mammals in nests of cavity nesting birds: Why should ornithologists study rodents? *Can. J. Zool.* 86: 286–293.
- Stański T., Walankiewicz W., Czeszczewik D. 2008. Absence of edge effects on nest predation in the Collared Flycatcher *Ficedula albicollis* in the primeval forest of Białowieża National Park, NE Poland. *Acta Ornithol.* 43: 92–96.
- Wesołowski T., Cholewa M. 2009. Climate variation and birds' breeding seasons in a primeval temperate forest. *Clim. Res.* 38: 199–208.
- Wesołowski T., Rowiński P., Maziarz M. 2009. Wood warbler *Phylloscopus sibilatrix* — a nomadic insectivore in search of safe breeding grounds? *Bird Study* 56: 26–33.

- Wesołowski T., Maziarz M. 2009. Changes in breeding phenology and performance of Wood Warbler *Phylloscopus sibilatrix* in a primeval forest: a thirty-years perspective. *Acta Ornithol.* 44: 69–80.
- Maziarz M., Wesołowski T. 2010. Timing of breeding and nestling diet of Wood Warbler *Phylloscopus sibilatrix* in relation to changing food supply. *Bird Study* 57: 540–552.
- Wesołowski T., Mitrus C., Czeszczewik D., Rowiński P. 2010. Breeding bird dynamics in a primeval temperate forest over 35 years: variation and stability in a changing world. *Acta Ornithol.* 45: 209–232.
- Wesołowski T. 2011. Blackcap *Sylvia atricapilla* numbers, phenology and reproduction in a primeval forest — a 33-year study. *J. Ornithol.* 152: 319–329.
- Cockle K. L., Martin K., Wesołowski T. 2011. Woodpeckers, decay, and the future of cavity-nesting vertebrate communities worldwide. *Frontiers Ecol. Environ.* 9: 377–382.
- Wesołowski T. 2011. "Lifespan" of woodpecker-made holes in a primeval temperate forest: a thirty year study. *For. Ecol. Manage.* 262: 1846–1852.
- Cholewa M., Wesołowski T. 2011. Nestling food of European hole-nesting passerines: do we know enough to test the adaptive hypotheses on breeding seasons? *Acta Ornithol.* 46: 105–116.
- Walankiewicz W., Czeszczewik D., Tumiel T., Stański T. 2011. Woodpeckers abundance in the Białowieża Forest — a comparison between deciduous, strictly protected and managed stands. *Ornis Polonica* 52: 161–168.
- Czeszczewik D., Ruczyński I., Zięba-Schraven K., Wiśniewska J., Walankiewicz W. 2012. The Pied and the Collared Flycatcher do not compete for microhabitats in the Białowieża Forest. *Belgian J. Zool.* 142: 133–137.
- Wesołowski T. 2012. "Lifespan" of non-excavated holes in a primeval temperate forest: A 30 year study. *Biol. Conserv.* 153: 118–126.
- Hebda G., Wesołowski T. 2012. Low flea loads in birds' nests in tree holes. *Ornis Fennica* 89: 139–144.
- Wesołowski T., Maziarz M. 2012. Dark tree cavities — a challenge for hole nesting birds? *J. Avian Biol.* 43: 454–460.
- Wesołowski T., Rowiński P. 2012. The breeding performance of Blue Tits *Cyanistes caeruleus* in relation to the attributes of natural holes in a primeval forest. *Bird Study* 59: 437–448.
- Wesołowski T., Fuller R. J. 2012. Spatial variation and temporal shifts in habitat use by birds at the European scale. In: Fuller R. (ed.). *Birds and habitat: relationships in changing landscapes*. Cambridge University Press, Cambridge, pp. 63–92.
- Tomiałojć L. 2012. Reproduction and population dynamics of Hawfinches *Coccothraustes coccothraustes* in the primeval forest of Białowieża National Park (NE Poland). *Acta Ornithol.* 47: 63–78.
- Czyż B., Rowiński P., Wesołowski T. 2012. No evidence for offspring sex ratio adjustment in Marsh Tits breeding in a primeval forest. *Acta Ornithol.* 47: 111–118.
- Hebda G., Pochrzast K., Mitrus S., Wesołowski T. 2013. Disappearance rates of old nest material from tree cavities: an experimental study. *Scand. J. For. Res.* 28: 445–450.
- Wesołowski T. 2013. Timing and stages of nest building by Marsh Tits *Poecile palustris* in a primeval forest. *Avian Biol. Res.* 6: 31–38.
- Maziarz M., Wesołowski T. 2013. Microclimate of tree cavities used by Great Tits (*Parus major*) in a primeval forest. *Avian Biol. Res.* 6: 47–56.
- Czeszczewik D., Walankiewicz W., Mitrus C., Tumiel T., Stański T., Sahel M., Bednarczyk G. 2013. Importance of dead wood resources for woodpeckers in coniferous stands of the Białowieża Forest. *Bird Conserv. Int.* 23: 414–425.
- Mettler R., Schaefer H. M., Chernetsov N., Fiedler W., Hobson K. A., Ilieva M., Imhof E., Johnsen A., Renner S. C., Rolshausen G., Serrano D., Wesołowski T., Segelbacher G. 2013. Contrasting patterns of genetic differentiation among Blackcaps (*Sylvia atricapilla*) with divergent migratory orientations in Europe. *PLOS ONE* 8: e81365.
- Walankiewicz W., Czeszczewik D., Stański T., Sahel M., Ruczyński I. 2014. Tree cavity resources in spruce-pine managed and protected stands of the Białowieża Forest, Poland. *Nat. Areas J.* 34: 423–428.
- Wesołowski T., Rowiński P. 2014. Do Blue Tits *Cyanistes caeruleus* synchronise reproduction with caterpillar peaks in a primeval forest? *Bird Study* 61: 235–245.
- Maziarz M., Wesołowski T. 2014. Does darkness limit the use of tree cavities for nesting by birds? *J. Ornithol.* 155: 793–799.
- Wesołowska W., Wesołowski T. 2014. Do *Leucochloridium* sporocysts manipulate the behaviour of their snail hosts? *J. Zool.* 292: 151–155.
- Kampichler Ch., Angeler D., Holmes R. T., Leito A., Svensson S., van der Jeugd H., Wesołowski T. 2014. Temporal dynamics of bird community composition: an analysis of base-line conditions from long-term data. *Oecologia* 175: 1301–1313.

- Hobson K. A., Van Wilgenburg S. L., Wesolowski T., Maziarz M., Bijlsma R. G., Grendelmeier A., Mallord J. W. 2014. A multi-isotope ($\delta^2\text{H}$, $\delta^{13}\text{C}$, $\delta^{15}\text{N}$) approach to establishing migratory connectivity in Palearctic-Afrotropical migrants: An example using Wood warblers. *Acta Ornithol.* 49: 57–69.
- Wesolowski T., Rowiński P., Maziarz M. 2015. Interannual variation in tree seed production in a primeval temperate forest: does masting prevail? *Eur. J. For. Res.* 134: 99–112.
- Wesolowski T. 2015. Dispersal in an extensive continuous forest habitat: Marsh Tit *Poecile palustris* in the Białowieża National Park. *J. Ornithol.* 156: 349–361.
- Maziarz M., Wesolowski T., Hebda G., Cholewa M. 2015. Natural nest-sites of Great Tits (*Parus major*) in a primeval temperate forest (Białowieża National Park, Poland). *J. Ornithol.* 156: 613–623.
- Wesolowski T., Czeszczewik D., Hebda G., Maziarz M., Mitrus C., Rowiński P. 2015. 40 years of breeding bird community dynamics in a primeval temperate forest (Białowieża National Park, Poland). *Acta Ornithol.* 50: 95–120.
- Czeszczewik D., Zub K., Stanski T., Sahel M., Kapusta A., Walankiewicz W. 2015. Effects of forest management on bird assemblages in the Białowieża Forest, Poland. *iForest* 8: 377–385.
- Czeszczewik D., Walankiewicz W. 2016. [Ecology and biology of birds in the Białowieża Forest: a 40-year perspective]. *Leśne Prace Badawcze* 77: 332–340.
- Kujawa A., Orczewska A., Falkowski M., Blicharska M., Bohdan A., Buchholz L., Chylarecki P., Gutowski J. M., Latałowa M., Mysłajek R. W., Nowak S., Walankiewicz W., Zalewska A. 2016. [The Białowieża Forest – a UNESCO National Heritage Site – protection priorities]. *Leśne Prace Badawcze* 77: 302–323.
- Walankiewicz W., Szymura M., Czeszczewik D. 2016. [A passive one-hundred years conservation of post-clear-cut tree stands in the Białowieża National Park led to restore of a breeding avifauna]. *Parki Narodowe i Rezerваты Przyrody* 35: 75–88.
- Wesolowski T., Cholewa M., Hebda G., Maziarz M., Rowiński P. 2016. Immense plasticity of timing of breeding in a sedentary forest passerine, *Poecile palustris*. *J. Avian Biol.* 47: 129–133.
- Mitrus S., Hebda G., Wesolowski T. 2016. Cohabitation of tree holes by ants and breeding birds in a temperate deciduous forest. *Scand. J. For. Res.* 31: 135–139.
- Maziarz M., Wesolowski T., Hebda G., Cholewa M., Broughton R. K. 2016. Breeding success of Great Tits *Parus major* in relation to attributes of natural nest-cavities in a primeval forest. *J. Ornithol.* 157: 343–354.
- Hebda G., Wesolowski T., Rowiński P. 2016. Nest sites of Middle Spotted Woodpeckers *Leiopicus medius* in a primeval forest. *Ardea* 104: 119–128.
- Zub K., Czeszczewik D., Ruczyński I., Kapusta A., Walankiewicz W. 2017. Silence is not golden: the hissing calls of tits affect the behaviour of a nest predator. *Behav. Ecol. Sociobiol.* 71: 1–7.
- Tomiałojć L., Neubauer G. 2017. Song Trush *Turdus philomelos* and Hawfinch *Coccothraustes coccothraustes* exhibit non random nest orientation in dense temperate forest. *Acta Ornithol.* 52: 209–220.
- Maziarz M., Broughton R. K., Wesolowski T. 2017. Microclimate in tree cavities and nest-boxes: implications for hole-nesting birds. *For. Ecol. Manage.* 389: 306–313.
- Hebda G., Wesolowski T., Rowiński P. 2017. Nest sites of a strong excavator, the Great Spotted Woodpecker *Dendrocopos major*, in the Białowieża National Park (Poland). *Ardea* 105: 61–71.
- Wesolowski T. 2017. Failed predator attacks: a direct test of security of tree cavities used by Marsh Tits *Poecile palustris*. *Auk* 134: 802–810.
- Wesolowski T., Neubauer G. 2017. Diet of Marsh Tit *Poecile palustris* nestlings in a primeval forest in relation to food supply and age of young. *Acta Ornithol.* 52: 105–118.
- Wesolowski T., Fuller R. J., Flade M. 2018. Temperate forests: A European perspective on variation and dynamics in bird assemblages. In: Mikusiński G., Roberge J.-M., Fuller R. J. (eds). *Ecology and conservation of forest birds*. Cambridge University Press, Cambridge, pp. 253–317.
- Wesolowski T., Martin K. 2018. Tree holes and hole-nesting birds in European and North American forests. In: Mikusiński G., Roberge J.-M., Fuller R. J. (eds). *Ecology and conservation of forest birds*. Cambridge University Press, Cambridge, pp. 79–134.
- Wesolowski T., Wierzcholska S. 2018. Tits as bryologists: patterns of moss use in nests of three species co-habiting in a primeval forest. *J. Ornithol.* 159: 733–745.
- Maziarz M., Broughton R. K., Hebda G., Wesolowski T. 2018. Occupation of songbird (*Phylloscopus sibilatrix*) nests by ants. *Insectes Sociaux* 65: 351–355.
- Mikusiński G., Bubnicki J. W., Churski M., Czeszczewik D., Walankiewicz W., Kuijper D. P. J. 2018. Is the impact of loggings in the last primeval lowland forest in Europe underestimated? The conservation issues of Białowieża Forest. *Biol. Conserv.* 227: 266–274.
- Kowalczyk R., Gutowski J. M., Jaroszewicz B., Niedziałkowski K., Rok J., Wesolowski T., Wójcik J. M. 2018. Puszcza Białowieska. Park Narodowy. Ochrona i rozwój regionalny. Białowieża.

- Maziarz M., Grendelmeier A., Wesołowski T., Arlettaz R., Broughton R. K., Pasinelli, G. 2019. Patterns of predator behaviour and Wood Warbler *Phylloscopus sibilatrix* nest survival in a primeval forest. *Ibis* 161: 854–866.
- Wesołowski T., Rowiński P., Neubauer G. 2019. Food of Nuthatch *Sitta europaea* young in a primeval forest: effects of varying food supply and age of nestlings. *Acta Ornithol.* 54: 85–104.
- Czeszczewik D., Ginter A., Mikusiński G., Pawłowska A., Kałuża H., Smithers R. J., Walankiewicz W. 2019. Birdwatching, logging and the local economy in the Białowieża Forest, Poland. *Biodiv. and Conserv.* 28: 2967–2975.
- Czeszczewik D., Czortek P., Jaroszewicz B., Zub K., Rowiński P., Walankiewicz W. 2020. Climate change has cascading effects on tree masting and the breeding performance of a forest songbird in a primeval forest. *Sci. Total Environ.* 747: 142084.
- Stański T., Czeszczewik D., Stańska M., Walankiewicz W. 2020. Foraging behaviour of the Great Spotted Woodpecker *Dendrocopos major* in relation to sex in primeval stands of the Białowieża National Park. *Acta Ornithol.* 55: 120–128.
- Wesołowski T., Hebda G., Rowiński P. 2021. Variation in timing of breeding of five woodpeckers in a primeval forest over 45 years: role of food, weather and climate. *J. Ornithol.* 162: 89–128.
- Yatsiuk Y., Wesołowski T. 2020. Diversity and abundance of large tree holes used by Tawny Owls *Strix aluco* in lowland temperate forests. *Bird Study* 67: 331–343.
- Stański T., Czeszczewik D., Stańska M., Walankiewicz W. 2021. Anvils of the Great Spotted Woodpecker (*Dendrocopos major*) in primeval oak-lime-hornbeam stands of the Białowieża National Park. *Eur. Zool. J.* 88: 1–8.
- Wesołowski T. 2021. Natural nest sites of European Starling *Sturnus vulgaris* in a primeval temperate forest. *Bird Study* 68: 145–156.
- Tomiałojć L. 2021. Impact of nest predators on migratory Woodpigeons *Columba palumbus* in Central Europe — breeding densities and nesting success in urban versus natural habitats. *Acta Ornithol.* 55: 139–154.
- Czeszczewik D., Kosiński Z., Kubicka K., Puła W., Walankiewicz W. 2022. Mortality and regeneration of elms in primeval stands of Białowieża National Park, Poland. *For. Pathol.* 52: e12731.
- Neubauer G., Wolska A., Rowiński P., Wesołowski T. 2022. N-mixture models estimate abundance reliably: a field test on Marsh Tit using time-for space substitution. *Ornithol. Applications* 124: duab054
- Wesołowski T., Czeszczewik D., Hebda G., Maziarz M., Mitrus C., Rowiński P., Neubauer G. 2022. 45 years of bird censuses in a primeval temperate forest (Białowieża National Park, Poland). *Acta Ornithol.* 57: 71–100.
- Maag N., Burgess M.D., Maziarz M., Lüpold S., Mallord J. W., Broughton R. K., Cristinacce A., Arlettaz R., Carlotti S., Castello J., Davis T., Gerber M., Grendelmeier A., Orsman C.J., Riess M., Stelbrink P., Wesołowski T., Züst Z., Pasinelli G. 2022. Reproduction of the wood warbler *Phylloscopus sibilatrix* varies across Europe. *J. Avian Biol.* e03033
- Wesołowski T. in press. Weather, food and predation shape the timing of Marsh Tit breeding in primeval conditions: a long-term study. *J. Ornithol.* DOI: 10.1007/s10336-022-02003-1