

The Euro+Med treatment of Hedera (Araliaceae) — recent studies and a new name

Author: Jury, Stephen L.

Source: Willdenowia, 34(2): 351-352

Published By: Botanic Garden and Botanical Museum Berlin (BGBM)

URL: https://doi.org/10.3372/wi.34.34202

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Willdenowia 34 – 2004 351

doi:10.3372/wi.34.34202 (available via http://dx.doi.org/)

Notulae ad floram euro-mediterraneam pertinentes No. 13

STEPHEN L. JURY

The Euro+Med treatment of *Hedera* (*Araliaceae*) – recent studies and a new name

Abstract

Jury, S. L.: The Euro+Med treatment of *Hedera (Araliaceae)* − recent studies and a new name. − Willdenowia 34: 351-352. − ISSN 0511-9618: © 2004 BGBM Berlin-Dahlem.

For the treatment of the *Hedera* taxa present in the territories included in the Euro+Med PlantBase project a new name is required and here validated, *Hedera rhizomatifera*, based on *Hedera helix* subsp. *rhizomatifera*.

The genus *Hedera* has been subject to a number of studies in recent years, chiefly by McAllister (in Watsonia 18: 7-15. 1990), Rutherford & al. (in Plantsman 15: 115-128. 1993), Vargas & al. (in Pl. Syst. Evol. 219: 165-179. 1999), Valcárcel & Vargas (in Anales Jard. Bot. Madrid 59: 363-368. 2002), Ackerfield & Wen (in Adansonia, ser. 3, 24: 197-212. 2002), Valcárcel & al. (in Molec. Phylog. Evol. 27: 1-20. 2003) and Jury & al. (in press). These have stirred up considerable controversy amongst ivy enthusiasts in the horticultural world, though supporting the radical treatment presented by McAllister & Rutherford (in Cullen & al., Eur. Gard. Fl. 5: 375-380. 1997) and Valcárcel & al. (in Castroviejo, Fl. Iber. 10: 3-12. 2003). Current editorial work for an EU-sponsored project to work on Euro+Med PlantBase, a new electronic plant information system for Europe, the extra countries bordering the Mediterranean Sea and Macaronesia, has brought to light some further anomalies in the nomenclature and taxonomy of the genus.

Hedera helix L. subsp. rhizomatifera was described by McAllister in 1993 for an unusual plant cultivated in the Liverpool University Botanic Garden, UK, originating from Prov. Huelva, Aracena area, April 1974, N. & J. Rutherford (LIV), and is known so far only from the provinces of Huelva, Sevilla, Cadíz, Málaga, Granada, Jaén, Cuenca, Castellón and Tarragona in southern and eastern Spain. It produces underground stolons, often of considerable length, clearly seen in pot cultivation when the plant produces numerous stems through the drainage holes in the base. The juvenile leaves are dark green, more or less uniform in size and shape, usually with three lobes rather than five, not showing the considerable variation to the adult type, with raised white silver veins on the surface and with trichomes appressed to the leaf surface. A clear separation is strongly

supported by the work of Ackerfield & Wen (2002) and I believe this taxon is best represented as a species in its own right, if the related *H. hibernica* (Kirchner) Bean and *H. azorica* Carrière are also separated and recognized at the specific level:

Hedera rhizomatifera (McAllister) Jury, **stat. nov.** ≡ *Hedera helix* subsp. *rhizomatifera* McAllister in Plantsman 15: 118, 119. 1993.

Address of the author:

Stephen L. Jury, Centre for Plant Diversity and Systematics, The School of Plant Sciences, The University of Reading, Whiteknights, Reading, RG6 6AS, UK; e-mail: s.l.jury@rdg.ac.uk