



Development and Characterization of Microsatellite Loci in the Pantropical Fern *Hypolepis punctata* (Dennstaedtiaceae)

Authors: Shang, Hui, Wang, Ying, and Yan, Yue-Hong

Source: Applications in Plant Sciences, 3(9)

Published By: Botanical Society of America

URL: <https://doi.org/10.3732/apps.1500047>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

DEVELOPMENT AND CHARACTERIZATION OF MICROSATELLITE LOCI IN THE PANTROPICAL FERN *HYPOLEPIS PUNCTATA* (DENNSTAEDTIACEAE)¹

HUI SHANG², YING WANG^{2,3}, AND YUE-HONG YAN^{2,4}

²Shanghai Chenshan Botanical Garden, Shanghai Chenshan Plant Science Research Center, Chinese Academy of Sciences, Shanghai 201602, People's Republic of China; and ³College of Life and Environmental Sciences, Shanghai Normal University, Shanghai 200234, People's Republic of China

- *Premise of the study:* Microsatellite loci were isolated in *Hypolepis punctata* (Dennstaedtiaceae) to further study the reproductive ecology of this species.
- *Methods and Results:* We developed 16 microsatellite loci from one sample of *H. punctata* using an enriched genomic library. These loci were characterized in 28 individuals. The number of alleles per locus ranged from two to 10, and the expected heterozygosity ranged from 0.036 to 0.845.
- *Conclusions:* The results indicate that the microsatellite markers can facilitate further studies on inferring the phylogeography and population genetics of *H. punctata* and related species.

Key words: Dennstaedtiaceae; *Hypolepis punctata*; microsatellite; phylogeography; population genetics.

Hypolepis punctata (Thunb.) Mett. ex Kuhn (Dennstaedtiaceae), i.e., downy ground fern, is a green, densely hairy, and glandulous fern that is widely distributed in tropical and subtropical regions in Asia and the Pacific (Brownsey, 1987). This plant is used in Chinese traditional medicine and contains pterisin, which has a cytotoxic effect on cancer cells (Lai, 2003). However, this species is often confused with *H. polypodioides* (Blume) Hook. and *H. resistens* (Kunze) Hook. (Xing and Wang, 2013), and using chloroplast markers (*rbcl*, *matK*, *trnL-F*, and *psbA-trnH*) is ineffective in improving identification accuracy (Shang et al., unpublished data). Moreover, *H. punctata* is an ideal species for studying mating system and sexual resource allocation because it exhibits high spore production and cloning habit. In addition, the wide distribution of this species may provide insight into the long-distance dispersal of homosporous ferns. Nuclear microsatellite markers are known as versatile molecular tools for ferns to solve the problem of inferring phylogeography or population genetics (Jiménez et al., 2008). In this study, we report on the development of 16 microsatellite markers for *H. punctata* to contribute to reproductive ecology and species differentiation research in the genus *Hypolepis* Bernh.

METHODS AND RESULTS

Total genomic DNA was extracted from the silica gel-dried leaves of an individual *H. punctata* specimen (voucher no.: JSL-WLSQ522; Appendix 1)

¹Manuscript received 27 April 2015; revision accepted 4 June 2015.

The authors thank C.-X. Wang, H.-J. Wei, Y.-F. Gu, and X.-L. Zhou for their assistance in field surveys and sample collection. This study was supported by the Shanghai Municipal Administration of Forestation and City Appearances (grant no. G152419 and F132421).

⁴Author for correspondence: yhyan@sibs.ac.cn

doi:10.3732/apps.1500047

collected from Wuling Mountain in Sangzhi County, Hunan Province, China, using a Plant Genomic DNA Kit (Tiangen Biotech, Beijing, China).

A microsatellite-enriched library was built following the method presented by Glenn and Schable (2005) with slight modifications. Genomic DNA was digested with *RsaI* and *XmnI* (New England Biolabs, Ipswich, Massachusetts, USA) at 37°C overnight and subsequently ligated to the double-stranded adapter (forward 5'-GTTTAAGGCCTAGCTAGCAGAATC-3', reverse 5'-pGATTCTGC-TAGCTAGGCCTTAAACAAA-3'). The ligated DNA was randomly linked to one of the two single-stranded biotinylated microsatellite probes (5'-(CA)₁₅-Biotin, 5'-(GA)₁₅-Biotin). The hybridized DNA was then captured by streptavidin-coated paramagnetic beads (Dynabeads M-280 Streptavidin; DYNAL BIOTECH, Oslo, Norway) and gathered using a magnetic particle-collecting unit (DynaMag-2 Magnet 12321D; Invitrogen, Waltham, Massachusetts, USA). The enriched DNA was amplified using the forward adapter as the primer. The product was then purified, ligated into the pGEM-T Easy Vector System (Promega Corporation, Madison, Wisconsin, USA), and cloned in Chemically Competent TOP10 *E. coli* cells (Tiangen Biotech). A total of 135 clones were selected and sequenced, in which 107 (~80%) contained simple sequence repeats. Among these, 83 had suitable lengths for primer design using Premier 5.0 (PREMIER BIOSOFT INTERNATIONAL, Palo Alto, California, USA). PCR amplifications were performed in 15-μL total volume with ~70 ng of genomic DNA, 10 μM of each primer, and 1× PCR mix (Tiangen Biotech). The PCR program consisted of 5 min of initial denaturation at 95°C, followed by 10 cycles of pre-PCR processing that involved 30 s of denaturation at 94°C, 30 s of annealing at 60°C, and 30 s of primer extension reaction at 72°C. The annealing temperature was reduced by 1°C per cycle. PCR amplification was continued for 25 more cycles at a constant annealing temperature of 50°C, and a final extension was performed at 72°C for 10 min. Finally, 16 pairs of primers (Table 1) were selected because they showed the clear bands of a single locus after agarose gel electrophoresis. The forward primer was labeled using one of the fluorescent dyes (FAM, TAMRA, or HEX) to detect polymorphism on an ABI 3730xl DNA analyzer (Applied Biosystems, Foster City, California, USA).

To test marker efficiency, we used 28 individuals of *H. punctata* from three different populations (five individuals from Wuling Mountain [voucher: JSL-WLSQ522]; 16 from Nanling Mountain [voucher: YYH13169]; and seven from Bawangling Mountain, Hainan Island, China [voucher: SG2984]; Appendix 1). Samples were collected from different individuals, with a minimum interval of 100 m between them, to avoid sampling the same clone. The numbers of alleles per locus, observed heterozygosity, and expected heterozygosity were estimated using CERVUS 3.0 (Kalinowski et al., 2007). In addition, cross-amplifications

TABLE 1. Characteristics of 16 microsatellite loci developed in *Hypolepis punctata*.

Locus	Primer sequences (5'–3')	Repeat motif	Allele size range (bp)	GenBank accession no.
SHH02	F: GTTGTGCGTAATCCGCAAAGTGG R: CAGATATGAGCGTCATTATCTCGGT	(TC) ₁₉	292–294	KR270806
SHH13	F: CATGGATTTGTTCTCCCTATCTGC R: TGGCCTTTGGGGAACCTTAGTA	(GA) ₃₉	362–378	KR270807
SHH17	F: CAGCAGCAGAGGAACCTGACA R: ATTCGAACCACCCATTGAC	(CT) ₁₅	445–447	KR270808
SHH19	F: TTGATGCCTCCATGACTATGCT R: TCACCTGTCTCCCTAACTTCT	(CA) ₂₄	264–288	KR270809
SHH23	F: CGGAGCGGAAAGGTAGAACA R: TTTTGCCACTATTGCTGATGAA	(AC) ₆ AT(AC) ₃	235–245	KR270810
SHH33	F: TCTCCCTCCCTCGATCTCCTT R: ATGTGGTGCTTCTAGCTGCTGAC	(CT) ₁₇	246–258	KR270811
SHH34	F: AACCGTAACAGACGTGCAAACC R: TGTGAGAAGCAGCAAGTCCAAA	(CT) ₂₀ (CA) ₉	441–453	KR270812
SHH44	F: TGGTATCATAGGCCATTTTGTCC R: TAGAGGAGGAGATGCATTGAGA	(CT) ₁₇ (CA) ₁₃	172–196	KR270813
SHH46	F: GGAATAAACCATGTAGGCAAGAGC R: CCAACGAGCCATGTGGACAA	(GA) ₁₃	121–123	KR270814
SHH51	F: TAGCAGTAAATAGTTTGTACGTGCC R: CCATCCGTTGTGCCCAT	(CA) ₆ AA(CA) ₃	246–249	KR270815
SHH55	F: GGAATCGCCAAGGAGATAATAA R: CCTCTTTTCTCAATCTATGTCCC	(AG) ₁₂	413–422	KR270816
SHH56	F: AGAAGATGCTTGTCTATAAGTAGGG R: AATGCTCAAGTCAAAGTGCC	(CT) ₂₀	421–438	KR270817
SHH65	F: TCGATAGTGTTCGCGGGTAA R: GGGCATGGTGGTGACAAAGT	(CT) ₂₃ (CA) ₁₁	271–283	KR270818
SHH71	F: TTTCTGTCTAAATCATGCTCTTTCC R: GCCTGTCTCGCTACCCGTAT	(CT) ₁₆	293–301	KR270819
SHH77	F: GATGAATAAAGAAGTAAACCAAC R: AGCAAGAAAGGGAGAACGAG	(CA) ₁₀	439–451	KR270820
SHH78	F: ACAGTGATGGAAGGCTGAAAGTC R: GAGATGGCGTACCTATGGATGG	(CT) ₁₀	238–242	KR270821

were performed to test the transferability of the marker to five other *Hypolepis* species (two individuals of *H. polypodioides* [vouchers: SG765, SG767], one individual of *H. resistens* [voucher: SG2900], one individual of *H. tenuifolia* (G. Forst.) Bernh. [voucher: HN31], two individuals of *H. pallida* (Blume) Hook. [vouchers: YYH11628, YYH11629], and one individual of *H. brooksiae* Alderw. [voucher: SIWS19]; Appendix 1).

The number of alleles per locus ranged from two to 10, with an average of 4.75 (Table 2). Meanwhile, 14 of the loci presented a significant bias between the observed and expected heterozygosities, which might indicate selfing in these populations (Table 2). Furthermore, at least six loci were interspecifically

amplifiable in each of the other five species. In particular, all 16 loci were amplifiable for *H. polypodioides* (Table 3).

CONCLUSIONS

A total of 16 polymorphic microsatellite loci were newly developed and characterized for *H. punctata*. These polymorphic microsatellite loci may provide good references for analyzing

TABLE 2. Genetic properties of the 16 newly developed microsatellites of *Hypolepis punctata*.

Locus	Total (n = 28)			Hainan (n = 7)			Wuling (n = 5)			Nanling (n = 16)		
	A	H _o	H _e	A	H _o	H _e	A	H _o	H _e	A	H _o	H _e
SHH02	2	0.036	0.036	1	0.000	0.000	1	0.000	0.000	2	0.063	0.063
SHH13	10	0.143	0.843	3	0.143	0.385	3	0.200	0.644	9	0.125	0.815
SHH17	3	0.071	0.491	2	0.000	0.264	2	0.200	0.556	3	0.063	0.179
SHH19	6	0.143	0.805	3	0.143	0.648	3	0.200	0.733	4	0.125	0.667
SHH23	3	0.143	0.137	1	0.000	0.000	2	0.400	0.356	3	0.125	0.123
SHH33	7	0.179	0.760	4	0.286	0.396	2	0.000	0.533	4	0.188	0.692
SHH34	6	0.143	0.732	2	0.000	0.440	3	0.400	0.644	3	0.125	0.573
SHH44	5	0.143	0.759	2	0.143	0.495	2	0.200	0.556	4	0.125	0.718
SHH46	2	0.143	0.468	2	0.143	0.143	2	0.200	0.556	3	0.188	0.534
SHH51	3	0.071	0.390	2	0.143	0.143	1	0.000	0.000	2	0.063	0.063
SHH55	5	0.143	0.606	4	0.143	0.495	2	0.200	0.200	2	0.125	0.444
SHH56	8	0.143	0.845	6	0.000	0.879	4	0.400	0.711	5	0.125	0.810
SHH65	6	0.143	0.779	2	0.143	0.143	2	0.200	0.200	4	0.125	0.756
SHH71	4	0.071	0.538	3	0.143	0.275	2	0.200	0.200	3	0.000	0.331
SHH77	4	0.071	0.610	1	0.000	0.000	1	0.000	0.000	3	0.125	0.486
SHH78	2	0.036	0.363	2	0.143	0.143	1	0.000	0.000	1	0.000	0.000

Note: A = number of sampled alleles; H_e = expected heterozygosity; H_o = observed heterozygosity.

TABLE 3. Cross-amplification length (in base pairs) of 16 microsatellite loci from *Hypolepis punctata* in other *Hypolepis* species.

Locus	<i>H. polypodioides</i> (n = 2)	<i>H. resistens</i> (n = 1)	<i>H. tenuifolia</i> (n = 1)	<i>H. pallida</i> (n = 2)	<i>H. brooksiae</i> (n = 1)
SHH02	292	292	278–292	—	292
SHH13	366–368	382	382	372	—
SHH17	443	—	—	485	447–449
SHH19	264–272	290	—	—	—
SHH23	235–243	233–235	225	227–239	239
SHH33	248	276	—	—	—
SHH34	449	457	—	—	—
SHH44	150	—	140	152	—
SHH46	121	111–113	149	115	121–123
SHH51	246	287	—	—	—
SHH55	416	—	—	—	421–423
SHH56	431–433	—	425	417	429–431
SHH65	273–281	—	—	277	281–283
SHH71	293	—	—	—	293
SHH77	443	449	—	—	445
SHH78	238	—	—	—	240

Note: — = failed amplification; n = number of individuals sampled.

mating systems and population structures, identifying clones, estimating gene flow, and identifying related species. This research will considerably improve knowledge on the life history of ferns. In addition, the high transferability of these loci to other species from the genus *Hypolepis* is essential for future research on hybridization or speciation.

LITERATURE CITED

- BROWNSEY, P. J. 1987. A review of the fern genus *Hypolepis* (Dennstaedtiaceae) in the Malesian and Pacific regions. *Blumea* 32: 227–276.
- GLENN, T. C., AND N. A. SCHABLE. 2005. Isolating microsatellite DNA loci. *Methods in Enzymology* 395: 202–222.
- JIMÉNEZ, A., L. G. QUINTANILLA, S. PAJARÓN, AND E. PANGUA. 2008. Reproductive and competitive interactions among gametophytes of the allotetraploid fern *Dryopteris corleyi* and its two diploid parents. *Annals of Botany* 102: 353–359.
- KALINOWSKI, S. T., M. L. TAPER, AND T. C. MARSHALL. 2007. Revising how the computer program CERVUS accommodates genotyping error increases success in paternity assignment. *Molecular Ecology* 16: 1099–1106.
- LAI, K. 2003. Studies on the cytotoxic principles from *Hypolepis punctata*. Master's thesis, Taipei Medical University, Taipei, Taiwan.
- XING, F., AND F. WANG. 2013. *Hypolepis*. In Z. Wu, P. H. Raven, and D. Hong [eds.], *Flora of China*, Vols. 2–3, 152–154. Science Press, Beijing, China, and Missouri Botanical Garden Press, St. Louis, Missouri, USA.

APPENDIX 1. Voucher and locality information of all *Hypolepis* samples used in this study.^a

Species	Voucher no.	Locality	Geographic coordinates
<i>H. punctata</i> (Thunb.) Mett. ex Kuhn	JSL-WLSQ522	Wuling Mountain, Hunan, China	29°18'31"N, 110°6'52"E
	YYH13169	Nanling Mountain, Guangdong, China	24°43'24"N, 114°15'46"E
	SG2984	Bawangling Mountain, Hainan, China	19°5'49"N, 109°13'32"E
<i>H. polypodioides</i> (Blume) Hook.	SG765, SG767	Fanjingshan Mountain, Guizhou, China	27°55'44"N, 108°41'17"E
<i>H. resistens</i> (Kunze) Hook.	SG2900	Bawangling Mountain, Hainan, China	19°5'28"N, 109°10'59"E
<i>H. tenuifolia</i> (G. Forst.) Bernh.	HN31	Wuzhishan Mountain, Hainan, China	18°55'1"N, 109°42'13"E
<i>H. pallida</i> (Blume) Hook.	YYH11628, YYH11629	Nantou County, Taiwan, China	NA
<i>H. brooksiae</i> Alderw.	SIWS19	Celebes Island, Indonesia	NA

Note: NA = not available.

^aSpecimens are deposited at the Shanghai Chenshan Botanical Garden Herbarium (CSH), except for voucher SIWS19, which is deposited at the Chinese National Herbarium, Institute of Botany, Chinese Academy of Sciences (PE).