



Isolation and Characterization of Microsatellite Loci for *Smilax sieboldii* (Smilacaceae)

Authors: Ru, Yalu, Cheng, Ruijing, Shang, Jing, Zhao, Yunpeng, Li, Pan, et al.

Source: Applications in Plant Sciences, 5(3)

Published By: Botanical Society of America

URL: <https://doi.org/10.3732/apps.1700001>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

ISOLATION AND CHARACTERIZATION OF MICROSATELLITE LOCI FOR *SMILAX SIEBOLDII* (SMILACACEAE)¹

YALU RU^{2,3,5}, RUIJING CHENG^{2,4,5}, JING SHANG^{2,4}, YUNPENG ZHAO^{2,3,6}, PAN LI^{2,3},
AND CHENGXIN FU^{2,3}

²Key Laboratory of Conservation Biology for Endangered Wildlife of the Ministry of Education, College of Life Sciences, Zhejiang University, Hangzhou 310058, People's Republic of China; ³Laboratory of Systematic and Evolutionary Botany and Biodiversity, Institute of Ecology and Conservation Center for Gene Resources of Endangered Wildlife, Zhejiang University, Hangzhou 310058, People's Republic of China; and ⁴College of Agriculture and Biotechnology, Zhejiang University, Hangzhou 310058, People's Republic of China

- **Premise of the study:** Polymorphic microsatellite markers were developed for *Smilax sieboldii* (Smilacaceae), a member of the *S. hispida* group with a biogeographic disjunction between eastern Asia and North America, to study the phylogeography and incipient speciation of this species and its close relatives.
- **Methods and Results:** Transcriptome sequencing produced 47,628 unigenes. Seventeen loci were developed from 122 randomly selected primer pairs. Polymorphism and genetic variation were evaluated for 68 accessions representing five populations of *S. sieboldii*. The number of alleles per locus ranged from four to 18; the expected heterozygosity ranged from 0.59 to 0.92. Twelve loci were successfully amplified in five related species: *S. scobinicaulis*, *S. californica*, *S. hispida*, *S. moranensis*, and *S. jalapensis*.
- **Conclusions:** These novel expressed sequence tag–derived microsatellite markers will facilitate further population genetic research of *S. sieboldii* and its close allies of the *S. hispida* group.

Key words: eastern Asian and North American disjunction; microsatellite primers; Smilacaceae; *Smilax sieboldii*; transcriptome sequencing.

The *Smilax hispida* group is a well-supported clade including six species in Smilacaceae (Qi et al., 2013) with a disjunct distribution including eastern Asia (*S. sieboldii* Miq. and *S. scobinicaulis* C. H. Wright), western North America (*S. californica* (A. DC.) A. Gray), eastern North America (*S. hispida* Raf.), and Mexico (*S. moranensis* M. Martens & Galeotti and *S. jalapensis* Schltdl.). *Smilax sieboldii* is a typical element of temperate broad-leaved forests that occurs widely in mainland China, Taiwan, Japan, and Korea. Previous studies based on two cpDNA intergenic regions indicated that at least four biogeographic lineages exist, with each lineage containing at least one private haplotype. This phylogeographic structure is considered to be related to the historical fluctuation of climate and sea level (Zhao et al., 2013). However, this study was limited by the lack of nuclear markers. Therefore, polymorphic microsatellite markers will enhance our understanding of population genetic diversity and

historical demography (e.g., gene flow, genetic bottlenecks) and will allow for connecting these patterns to geological and environmental changes.

Existing microsatellite markers for *Smilax* species (Xu et al., 2011; Martins et al., 2013) showed limited transferability and polymorphism for the *S. hispida* group due to phylogenetic distance. Therefore, in the current study we aimed to develop more polymorphic and transferable expressed sequence tag–simple sequence repeat (EST-SSR) markers from the transcriptome, which contains abundant ESTs, based on a high-throughput sequencing approach.

METHODS AND RESULTS

Transcriptome sequencing—Fresh young leaves of one wild accession of *S. sieboldii* were collected at Tianmu Mountain, Zhejiang Province, China (Appendix 1), and frozen in liquid nitrogen. RNA was extracted using TRIzol Reagent (Invitrogen Life Technologies, Carlsbad, California, USA) and treated with DNase (TaKaRa Bio, Shuzo, Kyoto, Japan) following the manufacturer's instructions. A 2 × 150-bp paired-end RNA-Seq library was prepared following the normalized eukaryote transcriptome library preparation protocol of the Beijing Genomics Institute (Shenzhen, China) and sequenced on the Illumina HiSeq 2500 platform (Illumina, San Diego, California, USA). A total of 65,863,062 raw reads were generated and uploaded to the National Center for Biotechnology Information (NCBI) Sequence Read Archive (accession SRP095761). The raw data were filtered using FASTX-TOOLKIT version 0.0.14 (Gordon and Hannon, 2010) by removing adapter sequences and low-quality reads with >5% unknown bases and/or >15% low-quality bases (quality value <20). Remaining reads were assembled into 66,482 transcripts using TRINITY version 2.3.2

¹Manuscript received 4 January 2017; revision accepted 13 February 2017.

The authors thank the editor and anonymous reviewers for their constructive comments that substantially improved the manuscript. This work was supported by the National Natural Science Foundation of China (no. 31461123001, 3151101152) and the National Project for Basic Work of Science and Technology of China (no. 2015FY110200).

⁵These authors contributed to this work equally.

⁶Author for correspondence: ypzhaoy@zju.edu.cn

doi:10.3732/apps.1700001

Applications in Plant Sciences 2017 5(3): 1700001; <http://www.bioone.org/loi/apps> © 2017 Ru et al. Published by the Botanical Society of America. This is an open access article distributed under the terms of the Creative Commons Attribution License (CC-BY-NC-SA 4.0), which permits unrestricted noncommercial use and redistribution provided that the original author and source are credited and the new work is distributed under the same license as the original.

TABLE 1. Characteristics of 17 newly developed microsatellite loci in *Smilax sieboldii*.^a

Locus	Primer sequences (5'–3')	Repeat motif	Allele size range (bp) ^b	Fluorescent dye ^c	GenBank accession no.	Function ^d	Organism	E-value
SS2	F: ACTGTAGGAGTTGACACAGAGG R: AGATTCTGGGAAACACAGGAAT	(GA) ₁₇	60–100	FAM	KY404961	Auxin response factor 15	<i>Oryza sativa</i> subsp. <i>japonica</i>	0
SS5	F: CAACCAAAACAAACAGAGAG R: GATACACGGGTAAACACACACC	(AG) ₁₂	96–132	TAMRA	KY404962	Hydrolase protein 30	<i>Arabidopsis thaliana</i>	5E-24
SS19	F: ACTTTGCCTATTAGCATCCGTT R: AGTACTGCTTCTCCACAAACAG	(CT) ₁₀	116–154	ROX	KY404963	Polygalacturonase inhibitor	<i>Pyrus communis</i>	5E-98
SS20	F: AACACACGATCTCAAGAAGAGC R: CGTCGTCATCTTCTCTCTGTTT	(GAA) ₁₅	89–122	FAM	KY404964	Protein FAF-like, chloroplastic	<i>Arabidopsis thaliana</i>	6E-18
SS21	F: GAATCCTTTTCGTTAGGGAAT R: CACAAGAAATAAAGAACGCTCG	(CT) ₁₂	107–137	TAMRA	KY404965	Probable ADP-ribosylation factor GTPase-activating protein AGD14	<i>Arabidopsis thaliana</i>	2E-15
SS33	F: AGTAGATCCAGCTTCTTTGAG R: CTCTCTCATCCCAATGTTCT	(AG) ₁₁	141–179	HEX	KY404966	Uncharacterized protein At4g08330, chloroplastic	<i>Arabidopsis thaliana</i>	2E-32
SS43	F: CAAGTATCCACACGAAACCAT R: GTGGAGGAACATGCAGTTGAT	(GA) ₁₁	154–180	HEX	KY404967	Oxygen-evolving enhancer protein 2, chloroplastic	<i>Fritillaria agrestis</i>	7E-119
SS74	F: GACGGCACCAAGAGAAGAT R: GTGGATATCATCACTCGGG	(CTG) ₈	181–241	FAM	KY404968	—	—	—
SS95	F: GTAGAGCGCTGGGTTC R: GCCAAGCTCTGGAAGAACAC	(TGG) ₈	135–180	ROX	KY404969	Sulfated surface glycoprotein 185	<i>Vohox carteri</i> PE	3E-06
SS100	F: GATTAGTGAGCITGGCGG R: ATGACCAACTCCTTCCAAC	(GAG) ₉	137–170	TAMRA	KY404970	Threonine-protein kinase-like protein At5g23170	<i>Arabidopsis thaliana</i>	2E-64
SS103	F: ACCACTGTCCCAGTTGCAT R: CTCCGAGGTTGTCAAAGAG	(TGG) ₁₀	263–281	ROX	KY404971	E3 ubiquitin-protein ligase At1g12760	<i>Arabidopsis thaliana</i>	8E-24
SS108	F: AAAGCCCCCAATTATCATC R: CGGCTGGAGAAGATGAATC	(TGC) ₁₃	106–124	FAM	KY404972	Formin-like protein 5	<i>Oryza sativa</i> subsp. <i>japonica</i>	5E-29
SS109	F: CCGCAAGTATTGAGGATGT R: GGTGAAGAGCTCAAAGACG	(ATC) ₁₄	139–175	HEX	KY404973	—	—	—
SS113	F: CTGATTCTCTCTCTGTTACGTTG R: CAATAAACCCAGCTTCAGCTCCTA	(CTGT) ₆	132–172	TAMRA	KY404974	—	—	—
SS114	F: TATTCGTGTAAGATACGTGGC R: TCGGCATTATTTTAATCACATC	(GTGTGA) ₉	137–167	ROX	KY404975	DNA-directed RNA polymerase II subunit 1	<i>Arabidopsis thaliana</i>	6E-09
SS120	F: ATATCCCGTCGAGTATCGTCTT R: GAGGAGGTGGGTACAGGGTAAG	(GCAGTA) ₄	146–200	ROX	KY404976	ABC transporter G family member 14	<i>Arabidopsis thaliana</i>	0
SS122	F: GACGGACTGACTGACTTTGGAT R: GGAATACTCAAGTTGCGCGTATC	(TAGCAC) ₄	125–185	HEX	KY404977	Protein PHLOEM PROTEIN 2-LIKE A1	<i>Arabidopsis thaliana</i>	7E-13

^a An annealing temperature of 58°C was used for all loci.
^b Size range values based on 68 individuals.
^c Forward 5' label.
^d The unigenes containing microsatellite loci were searched against the SWISS-PROT database (<http://www.expasy.ch/sprot/>); — = not found.

(Grabherr et al., 2011), which were then clustered into 47,628 unigenes with TGICL version 2.1 (Pertea et al., 2003).

Microsatellite development—Using the MicroSAteLLite identification tool (MISA) (Thiel et al., 2003), microsatellite regions in the unigenes were screened according to the following criteria for repeat numbers: dinucleotide repeats ≥ 6 , trinucleotide repeats ≥ 5 , and tetranucleotide, pentanucleotide, and hexanucleotide repeats ≥ 4 . Primers were designed for the screened microsatellite loci using Primer3 (Untergasser et al., 2012) with the default parameter settings. A total of 9263 microsatellite sequences were obtained, from which 2252 primer pairs were designed. Of these, 122 primer pairs were randomly selected and their forward primers were synthesized with one of three different universal primers (5'-CACGACGTTGTAAAAACGAC-3', 5'-TGTGGAATTGTGAGCGG-3', or 5'-CTATAGGGCACGCGTGGT-3') (Boutin-Ganache et al., 2001; Sakaguchi and Ito, 2014). To prevent primer dimers, hairpin structures, and mismatches, the best matches of forward primers and universal primers were selected using OLIGO version 6.67 (Molecular Biology Insights, Cascade, Colorado, USA).

We selected 12 accessions from various populations (Appendix 1) to test the effectiveness of primer amplification and to preliminarily assess genetic variation. Total genomic DNAs were extracted from silica-dried leaves using Plant DNAzol (Invitrogen Life Technologies). PCR amplifications were performed following the standard protocol of the Tsingke PCR kit (Tsingke Biotech Company, Beijing, China) in a final volume of 10 μ L, which contained approximately 5 ng of DNA, 5 μ L of 2 $\times$ PCR Master Mix, 0.1 μ M of forward primer, 0.4 μ M of reverse primer, and 0.3 μ M of fluorescently labeled universal primer (FAM, ROX, HEX, TAMRA; Table 1). The PCR thermal profile involved an initial denaturation at 95°C for 5 min; followed by 35 cycles of 94°C for 40 s, 58°C for 30 min, 72°C for 30 s; and a final 10-min extension step at 72°C. Fragment lengths of PCR products were analyzed on a 3730xl DNA Analyzer (Applied Biosystems, Foster City, California, USA) with GeneScan 500 LIZ as an internal reference (Applied Biosystems). Electrophoresis peaks were scored using GeneMarker version 2.2.0 (SoftGenetics, State College, Pennsylvania, USA). A total of 17 primer pairs with stable repeatability and high variation were selected for further analysis. All primer sequences obtained from this study were submitted to GenBank (Table 1).

Polymorphism assessment—To further evaluate the applicability of these primers, 68 individuals from five representative populations from China, Korea, and Japan (Appendix 1) were used to calculate genetic variation parameters. DNA extraction, PCR amplification, and length assessment of PCR products were performed following the procedures described above. The presence of null alleles and their bias on genetic diversity were evaluated based on the expectation maximization method implemented in FreeNA (Chapuis and Estoup, 2007). Deviation from Hardy–Weinberg equilibrium for each population and linkage disequilibrium for each primer pair were tested using GENEPOP version 4.0.7 (Rousset, 2008). The number of alleles, observed heterozygosity, expected heterozygosity, and polymorphism information content were calculated to assess the genetic polymorphism at each locus using CERVUS version 3.0.3 (Kalinowski et al., 2007).

Two loci (SS20, SS95) with high occurrence of null alleles ($>5\%$) were excluded from the following analysis. No significant deviation from Hardy–Weinberg equilibrium ($P < 0.001$) was observed for the remaining 15 loci except SS5 in populations CZJ and JFS; SS19 in population KMJ; and SS21, SS100, and SS109 in population JFS, which might be caused by Wahlund effect of specific populations. There was no evidence of significant linkage disequilibrium in any pair of loci. We detected 156 alleles in total, and the number of alleles at each locus ranged from four to 18, suggesting a moderate to high level of polymorphism. The observed heterozygosity, expected heterozygosity, and polymorphism information content for each locus ranged from 0.36 to 0.97, 0.59 to 0.92, and 0.53 to 0.91, respectively (Table 2).

Transferability evaluation—Transferability of the 15 primers was examined in the accessions of the five related species, i.e., five accessions each for *S. californica*, *S. hispida*, *S. moranensis*, and *S. jalapensis* and 10 accessions for *S. scobinicaulis* (Appendix 1). All loci were successfully amplified except two loci (SS21 and SS100) for *S. hispida* and one (SS33) for *S. moranensis* (Table 3). Polymorphism was detected in all but two loci (SS21 and SS100) for *S. californica*, five (SS2, SS19, SS103, SS120, and SS122) for *S. hispida*, four (SS21, SS74, SS103, and SS114) for *S. moranensis*, and one (SS100) for *S. jalapensis* (Table 3). The levels of both cross-amplifiability and polymorphism largely decreased with increasing phylogenetic distance. In total, 12 loci were amplifiable across the other five species in the *S. hispida* group.

TABLE 2. Genetic properties of the 15 newly developed microsatellite loci for *Smilax sieboldii*. Loci SS20 and SS95 are not included due to a high proportion ($>5\%$) of null alleles.^a

Locus	CTW ($n = 6$)				CZJ ($n = 14$)				CJS ($n = 15$)				KMJ ($n = 16$)				JFS ($n = 17$)				Total ($n = 68$)			
	A	H _o	H _e	PIC	A	H _o	H _e	PIC	A	H _o	H _e	PIC	A	H _o	H _e	PIC	A	H _o	H _e	PIC	A	H _o	H _e	PIC
SS2	4	0.83	0.77	0.65	8	1.00	0.86	0.80	8	0.87	0.85	0.80	9	0.88	0.86	0.82	10	0.82	0.84	0.79	17	0.88	0.92	0.90
SS5	4	1.00	0.78	0.65	10	0.86	0.88	0.83*	7	0.93	0.83	0.78	8	1.00	0.81	0.76	6	0.94	0.82	0.76*	14	0.94	0.90	0.88
SS19	4	0.83	0.74	0.62	7	0.79	0.74	0.67	4	0.60	0.71	0.63	8	0.81	0.87	0.83*	7	0.65	0.79	0.74	13	0.72	0.87	0.85
SS21	2	0.67	0.49	0.35	5	0.71	0.73	0.65	6	0.67	0.81	0.75	1	0.00	0.00	0.00	2	0.00	0.51	0.37*	9	0.36	0.79	0.76
SS33	5	1.00	0.74	0.64	10	0.86	0.89	0.84	4	1.00	0.76	0.68	7	1.00	0.77	0.71	10	1.00	0.88	0.83	18	0.97	0.92	0.91
SS43	5	1.00	0.82	0.70	5	0.71	0.77	0.71	4	1.00	0.72	0.64	5	1.00	0.78	0.71	8	0.71	0.70	0.66	12	0.87	0.87	0.85
SS74	8	1.00	0.91	0.81	5	0.46	0.63	0.55	5	0.40	0.36	0.34	4	0.56	0.60	0.50	9	0.77	0.79	0.74	16	0.60	0.77	0.75
SS100	4	0.67	0.71	0.60	7	0.86	0.77	0.71	6	0.73	0.73	0.67	5	0.86	0.74	0.67	6	0.38	0.56	0.70*	10	0.69	0.82	0.79
SS103	3	0.60	0.69	0.55	4	0.46	0.64	0.54	4	0.73	0.72	0.63	5	0.94	0.80	0.73	4	0.50	0.56	0.48	5	0.66	0.76	0.71
SS108	4	0.83	0.76	0.64	5	1.00	0.68	0.59	2	0.87	0.51	0.37	4	0.94	0.60	0.50	4	0.94	0.64	0.54	7	0.93	0.63	0.55
SS109	4	0.50	0.56	0.48	5	0.71	0.75	0.68	3	0.67	0.67	0.58	5	0.69	0.80	0.74	4	0.47	0.64	0.56*	9	0.62	0.84	0.82
SS113	4	1.00	0.76	0.64	4	0.71	0.55	0.45	5	0.87	0.63	0.56	4	0.44	0.47	0.43	3	0.53	0.42	0.34	9	0.66	0.59	0.53
SS114	3	0.67	0.55	0.45	3	0.79	0.62	0.53	4	0.87	0.66	0.57	3	0.31	0.46	0.40	3	0.77	0.55	0.47	4	0.68	0.69	0.62
SS120	3	1.00	0.67	0.54	6	0.50	0.72	0.65	2	0.40	0.41	0.32	4	0.69	0.73	0.65*	2	0.31	0.27	0.23	8	0.52	0.67	0.64
SS122	4	0.83	0.77	0.65	3	0.33	0.45	0.37	4	0.33	0.41	0.37	4	0.86	0.66	0.57	4	0.67	0.65	0.57	5	0.58	0.70	0.64

Note: A = number of alleles sampled; H_e = expected heterozygosity; H_o = observed heterozygosity; n = number of individuals sampled; PIC = polymorphism information content.
^a Voucher and locality information are provided in Appendix 1.
* Significant deviation from Hardy–Weinberg equilibrium ($P < 0.001$).

TABLE 3. Fragment sizes detected in cross-amplification tests of the 15 newly developed microsatellite markers in the remaining five species of the *Smilax hispida* group.^a

Locus	<i>S. scobinicaulis</i> (n = 10)	<i>S. californica</i> (n = 5)	<i>S. hispida</i> (n = 5)	<i>S. moranensis</i> (n = 5)	<i>S. jalapensis</i> (n = 5)
SS2	66–80	72–84	72	72–76	66–84
SS5	98–124	114–116	114–116	114–124	114–116
SS19	116–150	132–136	140	132–138	128–138
SS21	125–127	125	—	131	123–129
SS33	157–179	167–177	167–179	—	167–179
SS43	164–176	168–188	168–170	164–176	166–168
SS74	184–241	193–199	202–217	196	199–214
SS100	152–170	164	—	152–164	164
SS103	263–278	272–278	272	278	257–278
SS108	106–118	109–118	106–118	106–109	94–118
SS109	172–175	127–142	136–142	127–136	127–151
SS113	132–164	160–164	160–164	156–164	132–156
SS114	137–155	137–149	143–149	137	137–155
SS120	146–164	170–182	158	152–170	164–182
SS122	125–179	167–191	173	173–191	167–179

Note: — = amplification failed.
^aVoucher and locality information are provided in Appendix 1.

CONCLUSIONS

Using high-throughput sequencing, we sequenced and assembled the transcriptome of *S. sieboldii* without a reference genome. Fifteen EST-SSR markers were successfully developed to evaluate the genetic structure and demography of *S. sieboldii*, of which 12 are likely to be useful for all six species of the *S. hispida* group.

LITERATURE CITED

BOUTIN-GANACHE, I., M. RAPOSO, M. RAYMOND, AND C. F. DESCHÉPPER. 2001. M13-tailed primers improve the readability and usability of microsatellite analyses performed with two different allele-sizing methods. *BioTechniques* 31: 24–28.

CHAPUIS, M., AND A. ESTOUP. 2007. Microsatellite null alleles and estimation of population differentiation. *Molecular Biology and Evolution* 24: 621–631.

GORDON, A., AND G. J. HANNON. 2010. FASTX-TOOLKIT, version 0.0.14. Computer program and documentation distributed by the author.

Website http://hannonlab.cshl.edu/fastx_toolkit [accessed 1 February 2017].

GRABHERR, M. G., B. J. HAAS, M. YASSOUR, J. Z. LEVIN, D. A. THOMPSON, I. AMIT, X. ADICONIS, ET AL. 2011. Full-length transcriptome assembly from RNA-Seq data without a reference genome. *Nature Biotechnology* 29: 644–652.

KALINOWSKI, S. T., M. L. TAPER, AND T. C. MARSHALL. 2007. Revising how the computer program CERVUS accommodates genotyping error increases success in paternity assignment. *Molecular Ecology* 16: 1099–1106.

MARTINS, A. R., A. G. ABREU, M. M. BAJAY, P. M. S. VILLELA, C. E. A. BATISTA, M. MONTEIRO, A. ALVES-PEREIRA, ET AL. 2013. Development and characterization of microsatellite markers for the medicinal plant *Smilax brasiliensis* (Smilacaceae) and related species. *Applications in Plant Sciences* 1: 1200507.

PERTEA, G., X. Q. HUANG, F. LIANG, V. ANTONESCU, R. SULTANA, S. KARAMYCHEVA, Y. LEE, ET AL. 2003. TIGR gene indices clustering tools (TGICL): A software system for fast clustering of large EST datasets. *Bioinformatics (Oxford, England)* 19: 651–652.

QI, Z. C., K. M. CAMERON, P. LI, Y. P. ZHAO, S. C. CHEN, G. C. CHEN, AND C. X. FU. 2013. Phylogenetics, character evolution, and distribution patterns of the greenbriers, Smilacaceae (Liliales), a near-cosmopolitan family of monocots. *Botanical Journal of the Linnean Society* 173: 535–548.

ROUSSET, F. 2008. GENEPOP’007: A complete re-implementation of the GENEPOP software for Windows and Linux. *Molecular Ecology Resources* 8: 103–106.

SAKAGUCHI, S., AND M. ITO. 2014. Development and characterization of EST-SSR markers for the *Solidago virgaurea* complex (Asteraceae) in the Japanese archipelago. *Applications in Plant Sciences* 2: 1400035.

THIEL, T., W. MICHALEK, R. K. VARSHNEY, AND A. GRANER. 2003. Exploiting EST databases for the development and characterization of gene-derived SSR-markers in barley (*Hordeum vulgare* L.). *Theoretical and Applied Genetics* 106: 411–422.

UNTERGASSER, A., I. CUTCUTACHE, T. KORESSAAR, J. YE, B. C. FAIRCLOTH, M. REMM, AND S. G. ROZEN. 2012. Primer3: New capabilities and interfaces. *Nucleic Acids Research* 40: e115.

XU, X. H., Y. WAN, Z. C. QI, Y. X. QIU, AND C. X. FU. 2011. Isolation of compound microsatellite markers for the common Mediterranean shrub *Smilax aspera* (Smilacaceae). *American Journal of Botany* 98: e64–e66.

ZHAO, Y. P., Z. C. QI, W. W. MA, Q. Y. DAI, P. LI, K. M. CAMERON, J. K. LEE, ET AL. 2013. Comparative phylogeography of the *Smilax hispida* group (Smilacaceae) in eastern Asia and North America: Implications for allopatric speciation, causes of diversity disparity, and origins of temperate elements in Mexico. *Molecular Phylogenetics and Evolution* 68: 300–311.

APPENDIX 1. Voucher information for *Smilax* species used in this study.

Species	Population code	Voucher specimens ^a	Collection locality	Geographic coordinates	n
<i>Smilax sieboldii</i> Miq.	CTW	Xiaoxian Liu, 0812003	Mt. Zhu, Taiwan, China	23.31000N, 120.50000E	6
<i>Smilax sieboldii</i>	CZJ	Yalu Ru, Ru150921001	Mt. Tianmu, Zhejiang, China	30.37809N, 119.42061E	14
<i>Smilax sieboldii</i>	CJS	Yunpeng Zhao, HZU00441	Mt. Longchi, Jiangsu, China	31.24818N, 119.74551E	15
<i>Smilax sieboldii</i>	KMJ	Joongku Lee, GG13	Myeongjisan, Gyeonggi-do, Korea	37.93458N, 127.47325E	16
<i>Smilax sieboldii</i>	JFS	Chengxin Fu & Xinjie Jin, Fu1505092	Fujiyama, Tokyo, Japan	35.50281N, 138.76985E	17
<i>Smilax scobinicaulis</i> C. H. Wright		Pan Li, LP150444	Mt. Wuzhi, Hubei, China	31.08961N, 110.88390E	10
<i>Smilax californica</i> (A. DC.) A. Gray		Pan Li, LP150436	Near Shasta Lake, CA, USA	40.75954N, 122.03657W	5
<i>Smilax hispida</i> Raf.		Yunpeng Zhao, 090834	Croatian National Forest, NC, USA	36.20339N, 86.98333W	5
<i>Smilax jalapensis</i> Schldtl.		Pan Li, US10041	Teopisca, Chiapas, Mexico	16.57310N, 92.50445W	5
<i>Smilax moranensis</i> M. Martens & Galeotti		Pan Li, US10031	Mexico City, Mexico	19.30541N, 99.30743W	5

Note: n = number of individuals sampled.
^aVouchers were deposited in the Herbarium of Zhejiang University (HZU), Hangzhou, Zhejiang, China.