



Isolation and Characterization of Microsatellite Primers for the Critically Endangered Shrub *Styphelia longissima* (Ericaceae)

Authors: Anthony, Janet M., Allcock, Richard J. N., Dobrowolski, Mark P., and Krauss, Siegfried L.

Source: Applications in Plant Sciences, 5(11)

Published By: Botanical Society of America

URL: <https://doi.org/10.3732/apps.1700108>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

ISOLATION AND CHARACTERIZATION OF MICROSATELLITE PRIMERS FOR THE CRITICALLY ENDANGERED SHRUB *STYPHELIA* *LONGISSIMA* (ERICACEAE)¹

JANET M. ANTHONY^{2,3,6}, RICHARD J. N. ALLCOCK⁴, MARK P. DOBROWOLSKI⁵, AND
SIEGFRIED L. KRAUSS^{2,3}

²Kings Park and Botanic Garden, Botanic Garden and Parks Authority, Fraser Avenue, West Perth, 6005 Western Australia, Australia; ³School of Biological Sciences, The University of Western Australia, Crawley, 6009 Western Australia, Australia;

⁴Lotterywest State Biomedical Facility: Genomics, School of Pathology and Laboratory Medicine, The University of Western Australia, Crawley, Western Australia 6009, Australia; and ⁵Iluka Resources Limited, Level 23, 140 St. Georges Terrace, Perth, Western Australia 6000, Australia

- **Premise of the study:** Microsatellite markers were developed for population genetic analysis in the rare shrub *Styphelia longissima* (Ericaceae).
- **Methods and Results:** We generated ca. 2.5 million sequence reads using a Personal Genome Machine semiconductor sequencer. Using the QDD pipeline, we designed primers for >12,000 sequences with PCR product lengths of 80–480 bp. From these, 30 primer pairs were selected and screened using PCR; of these, 16 loci were found to be polymorphic, four loci were monomorphic, and 10 loci did not amplify reliably for *S. longissima*. For a sample of 57 plants from the only known population, the number of alleles observed for these 16 loci ranged from two to 21 and expected heterozygosity ranged from 0.49 to 0.91. These markers were also amplified in *Astroloma xerophyllum*, a closely related species.
- **Conclusions:** These markers will be used to characterize population genetic variation, spatial genetic structure, mating system parameters, and dispersal to aid in the management and conservation of the rare shrub *S. longissima*.

Key words: *Astroloma xerophyllum*; Ericaceae; microsatellite primers; shotgun sequencing; *Styphelia longissima*.

Styphelia longissima Hislop & Puente-Lel. (Ericaceae) is a newly described species (Hislop and Puente-Lelièvre, 2017) and found only in a single population on sand within remnant kwongan vegetation near Eneabba, in the South West Australian Floristic Region (SWAFR), an international biodiversity hotspot (Hopper and Gioia, 2004). Until very recently, this taxon was assigned to the genus *Leucopogon* R. Br. (Ericaceae) with the temporary name of *Leucopogon* sp. ciliate Eneabba (F. Obbens & C. Godden s.n. 3/7/2003). Recent taxonomic revision has placed this and some other *Leucopogon* species within the Styphelieae, with sister taxa *Leucopogon* sp. Ongerup and *Astroloma* sp. sessile leaf (Puente-Lelièvre et al., 2015; Hislop and Puente-Lelièvre, 2017). In 2007, the population consisted of just 1993 individuals (Woodman Environmental Consulting Pty. Ltd., 2008). Since then, however, mortality has substantially exceeded recruitment (Harris, 2013), and the species is currently listed as Rare Flora under the Wildlife Conservation Act 1950 (Western Australian Minister for the Environment, 2015). *Styphelia longissima* is a spindly to dense shrub, to 0.5(–0.8) m high, with cream-white colored flowers in July that are most likely insect pollinated, and

seed dispersal is myrmecochorous (Harris, 2013). Microsatellite markers were developed for *S. longissima* to enable an assessment of population genetic variation, spatial genetic structure, mating system parameters, and dispersal for management and conservation. *Astroloma xerophyllum* (DC.) Sond., a sister taxon, was chosen for cross-amplification based on molecular phylogeny of the Styphelieae (Puente-Lelièvre et al., 2015).

METHODS AND RESULTS

Genomic DNA was extracted from a single tissue-cultured plant, sampled from the only known population (Appendix 1), using a Carlsons method (Carlson et al., 1991) with modifications outlined in Anthony et al. (2016). Next-generation sequencing was performed on a Personal Genome Machine (PGM) semiconductor sequencer (Life Technologies, Carlsbad, California, USA) at the Lotterywest State Biomedical Facility Genomics Node in Perth, Western Australia, as described previously (Anthony et al., 2016). After sequencing, signal processing, base-calling, and quality trimming were performed using the default settings of TorrentSuite 4.0 (Thermo Fisher Scientific, Waltham, Massachusetts, USA), and library-specific FASTQ files were also generated. Sequencing resulted in >2.5 million reads, with a modal read length of 344 bp and a total data output of 766 Mb (National Center for Biotechnology Information [NCBI] Sequence Read Archive Bioproject no. PRJNA397350).

The raw sequences were screened using QDD version 3.1 pipeline (Megléc et al., 2014) to remove redundant sequences and design primers for >12,000 sequences with PCR product lengths of 80–480 bp. The default parameters of the program were used both for the screening steps and for primer design. The

¹Manuscript received 6 September 2017; revision accepted 11 October 2017.

⁶Author for correspondence: janet.anthony@dbca.wa.gov.au

doi:10.3732/apps.1700108

Applications in Plant Sciences 2017 5(11): 1700108; <http://www.bioone.org/loi/apps> © 2017 Anthony et al. Published by the Botanical Society of America. This is an open access article distributed under the terms of the Creative Commons Attribution License (CC-BY-NC-SA 4.0), which permits unrestricted noncommercial use and redistribution provided that the original author and source are credited and the new work is distributed under the same license as the original.

resultant sequences were filtered to ensure that the primer was not overlapping the repeat sequence, there were no poly-‘A’ or poly-‘T’ runs for more than seven base pairs within the sequence, and there was only one repeat motif between the primers. Subsequently, 30 primer pairs were selected based on the suggestions of Megléc (2014).

Initial screening was performed with CFX96 Touch Real-Time PCR Detection System (Bio-Rad Laboratories, Hercules, California, USA) using 5 µL of SsOAdvanced Universal SYBR Green Supermix (Bio-Rad Laboratories), 0.2 µM each of forward and reverse primers, and 5–10 ng of genomic DNA in a 10-µL reaction volume. Initially, screening included reliable amplification of a single sample across a range of temperatures to determine the most appropriate annealing temperature, followed by evidence of polymorphism among eight individuals. Consequently, 16 polymorphic loci (including three loci that were fixed heterozygotes) and four monomorphic loci were selected to complete the study (Table 1),

and the remaining 10 loci did not amplify reliably for *S. longissima*. Amplification of 57 individuals using 16 polymorphic loci was performed using a Veriti thermocycler (Life Technologies) within four multiplex mixes containing 6.25 µL of 2× Multmix and 2.25 µL of 5× Q-Solution (Type-It Microsatellite PCR kit; QIAGEN, Hilden, Germany), 1.25 µL of primer mix, and 2.25 µL of 5–100 ng DNA in a 12-µL reaction. Primer Mix (PM) 1 contained the primers SI36, SI65, SI53; PM 2 contained SI17, SI18, SI57, SI60; PM 3 contained SI6, SI26; and PM 4 contained SI01, SI47, SI67, SI71 using the following PCR conditions: an initial 1-min denaturation at 95°C; 35 cycles of 94°C for 10 s, 62°C (PM 1, 2, 3) or 56°C (PM 4) for 30 s, and 72°C for 45 s; followed by a final extension of 15 min at 72°C. Electrophoresis was performed using the ABI 3500 sequencer (Life Technologies), and allele sizes were determined using Geneious version 7.1 (Biomatters Ltd., Auckland, New Zealand). Multiple replicate runs were performed to ensure the accuracy of the final data set. Genetic diversity parameters were

TABLE 1. Characteristics of microsatellite loci developed for *Styphelia longissima* and cross-amplified in *Astroloma xerophyllum*.^a

Locus	Primer sequences (5′–3′)	Repeat motif	Allele size range (bp)		<i>T_a</i> (°C)	Fluorescent label	GenBank accession no.
			<i>S. longissima</i> (N = 57)	<i>A. xerophyllum</i> (N = 5)			
SI01	F: TGGTTATACTCAGCTTATGTATTGG R: AAGATTTACTTGCATTCTGCCA	(CT) ₁₄	203–223	201–205	56	6-FAM	KY559296
SI06	F: CACAAGTGGTTGATGAGTCCC R: CATCCTTCTCCCTTCCCTTC	(AG) ₁₄	87–111	187–228	56	VIC	KY559297
SI16	F: GGCTCACCAATTGGCTACAT R: CACTTGTGTACTTGTGACTTCATTTG	(AG) ₁₃	—	326 ^b	62	NED	MF405897
SI17	F: CGCTGCTTTACATCTTTGCTC R: CCTGATTTCCCTGAACCTCA	(CA) ₁₃	113–133	159–233	62	VIC	KY559298
SI18	F: CCAGAAAGTCTTCTCTCCCT R: ATCGGGAAGTGGAAGGTCAG	(AC) ₁₃	98–112	187–194	62	PET	KY559299
SI22	F: AGCCGCGGAACATATACAG R: GGACCAATTCATTCACTTGGAG	(AAG) ₁₂	—	163 ^b	56	6-FAM	MF405898
SI24	F: TCTCGACAGGAAGTGGAGT R: GGTACCAATATGTACACGGCA	(AG) ₁₂	345 ^b	254–262	56	NED	MF405899
SI26	F: AGCGTTCAAGTAAGGCGAAA R: GCGGATGGCCACTTTACTAC	(AG) ₁₂	155–213	153–161	62	NED	KY559300
SI29	F: GGAACACATCGTAACGTGA R: ACCACGCAAACTGTGAAA	(GA) ₁₂	—	274–276	56	6-FAM	MF405900
SI32	F: CGATCGATCAAACTCTCAGCA R: GACCAAGAAGACAAGAAGACGA	(AG) ₁₁	207 ^b	—	56	6-FAM	MF405901
SI36	F: CCCACGATAGCTCAGAGGT R: GGGATACAGCAGGTAAGTTCA	(GAA) ₁₁	136–185	136–148	62	VIC	KY559301
SI45	F: TTGGTCGTAATCACTTGGGAG R: GGCATTGGCACAACCTTCAG	(AG) ₁₁	153 ^b	—	56	NED	MF405902
SI47	F: TTTGTTTACAGAAGCTTAGCGG R: TTTGTAGTTCTTTCAGAATTGAGGA	(GA) ₁₀	238–246	233–250	56	VIC	KY559302
SI51	F: AAATGAGCTGAGCTGAATGC R: GGGTGTTCCTCACTTAACCTGC	(TG) ₁₀	—	138–150	56	VIC	MF405903
SI53	F: GGAAATCACAATCAGACCA R: TAGACATCCCTTCCGGATCA	(CA) ₁₀	115–141	114 ^b	62	PET	KY559303
SI57	F: AACACCAAACTAGGAAAGCCG R: ATTCAGAAATCAGGACCTGCC	(AG) ₁₀	89–107	—	62	VIC	KY559304
SI60	F: AATTGGGTACGTGACATCGG R: ATGAACGTCAGCCCATTTCC	(GA) ₁₀	172–262	—	62	NED	KY559305
SI62	F: GAGAGGGACTTCGGAACAAA R: CCCACTCTCCAAATCCTTCA	(AG) ₁₀	180, 182 ^c	—	56	VIC	MF405904
SI65	F: ACTGGTCAGCCTGTTAGCGT R: CGCAAGTGAAATCAAGTACGA	(CT) ₉	290–338	—	62	NED	KY559306
SI67	F: TTCCAAGATAACTAAATCATATACACA R: TGAATGTTGTCTATGGTTCTG	(TTG) ₉	143–176	133–153	56	PET	KY559307
SI68	F: TGCAATCAAATTACTTACAGCCA R: CAACCCTGTTCAATTCCACC	(AAG) ₉	240 ^b	—	56	PET	MF405905
SI70	F: TGTGTAGGTTCTGATTCTTATTCG R: CTCTGGACGCAAGGTACTGA	(AC) ₉	164, 168 ^c	—	56	PET	MF405906
SI71	F: ACCTCCAACACCAATGAACC R: ACCTCCAACACCAATGAACC	(CA) ₉	315–349	314–329	56	VIC	KY559308
SI75	F: TTGCACTTTGAGCCTCTCATTT R: TTGCTTCAATAAGTGGTAATTT	(AGT) ₈	125, 129 ^c	133 ^b	56	6-FAM	MF405907

Note: — = no amplification; N = number of individuals used in this study; *T_a* = annealing temperature.

^aVoucher and locality information are provided in Appendix 1.

^bMonomorphic loci.

^cFixed heterozygotes.

TABLE 2. Results of primer screening with 16 polymorphic primers for *Styphelia longissima*.

Locus	A	H _o	H _e	HWE ^a
SI01 ^b	7	0.421	0.727	***
SI06	11	0.825	0.825	ns
SI17	12	0.825	0.857	ns
SI18 ^b	7	0.368	0.692	***
SI26 ^b	21	0.691	0.914	***
SI36 ^b	8	0.456	0.591	***
SI47	5	0.439	0.491	ns
SI53 ^b	5	0.246	0.647	***
SI57 ^{b,c}	8	0.614	0.771	*
SI60 ^b	18	0.386	0.821	***
SI62	2	1.00	0.500	***
SI65 ^b	12	0.386	0.711	***
SI67	7	0.737	0.776	ns
SI70	2	1.00	0.500	***
SI71 ^{b,c}	9	0.474	0.690	***
SI75	2	1.00	0.500	***

Note: A = number of alleles; H_e = expected heterozygosity; H_o = observed heterozygosity; HWE = Hardy–Weinberg equilibrium.
^aSignificant departures from HWE are indicated as *P ≤ 0.05, ***P ≤ 0.001, ns = not significant.
^bEvidence suggesting null alleles.
^cEvidence of stuttering.

calculated using GenAEx version 6.4 (Peakall and Smouse, 2006). Departure from Hardy–Weinberg equilibrium (HWE) was assessed for each locus and population by χ^2 tests, and the possibility of null alleles was checked using MICRO-CHECKER version 2.2.3 (van Oosterhout et al., 2004).
The number of alleles observed for the 16 polymorphic loci ranged from two to 21, and the observed and expected heterozygosities ranged from 0.25 to 1.00 and 0.49 to 0.91, respectively (Table 2). There were significant departures from HWE for 12 of the 16 loci, with four loci showing no evidence of stuttering, large allele dropout, or null alleles (Table 2). Two loci showed evidence of stuttering indicated as a deficit of heterozygote genotypes with alleles of one base pair repeat difference. These loci have been thoroughly checked by repeat PCR and analysis scoring, and the one base pair difference shown to be real. These departures from HWE are most likely due to inbreeding within a small isolated population.
Using the same extraction method, five individuals were cross-amplified for the sister taxon *A. xerophyllum*. Amplification was performed using the same initial screening method outlined above and resulted in 12 polymorphic and four monomorphic loci (Table 1).

CONCLUSIONS

These 16 polymorphic microsatellites will be used for conservation genetic studies in the rare *S. longissima* to underpin

management and conservation. These microsatellites are likely to be useful for genetic studies in other related species given the initial success in cross-amplification for *A. xerophyllum*.

LITERATURE CITED

ANTHONY, J. M., R. J. N. ALLCOCK, AND S. L. KRAUSS. 2016. Isolation and characterization of 13 microsatellites for the rare endemic shrub *Tetradlea erubescens* (Elaeocarpaceae). *Applications in Plant Sciences* 4: 1500102.
CARLSON, J. E., L. K. TULSIERAM, J. C. GLAUBITZ, V. W. K. LUK, C. KAUFFELDT, AND R. RUTLEDGE. 1991. Segregation of random amplified DNA markers in F₁ progeny of conifers. *Theoretical and Applied Genetics* 83: 194–200.
HARRIS, B. 2013. Development of biotechnology-based ex situ conservation approaches for the critically endangered *Leucopogon* sp. ciliate Eneabba (Ericaceae). Ph.D. Thesis, The University of Western Australia, Perth, Western Australia, Australia.
HISLOP, M., AND C. PUENTE-LELIEVRE. 2017. Five new species of *Styphelia* (Ericaceae: Epacridoideae: Stypheliaceae) from the Geraldton Sandplains, including notes on a new, expanded circumscription for the genus. *Nuytsia* 28: 95–114.
HOPPER, S. D., AND P. GIOIA. 2004. The Southwest Australian Floristic Region: Evolution and conservation of a global hot spot of biodiversity. *Annual Review of Ecology Evolution and Systematics* 35: 623–650.
MEGLÉCZ, E. 2014. QDD: Program for microsatellite selection and primer design. Website <http://net.imbe.fr/~emeglec/qdd.html#choice> [accessed 17 October 2016].
MEGLÉCZ, E., N. PECH, A. GILLES, V. DUBUT, P. HINGAMP, A. TRILLES, R. GRENIER, AND J. F. MARTIN. 2014. QDD version 3.1: A user-friendly computer program for microsatellite selection and primer design revisited: Experimental validation of variables determining genotyping success rate. *Molecular Ecology Resources* 14: 1302–1313.
PEAKALL, R., AND P. E. SMOUSE. 2006. GenAEx 6: Genetic analysis in Excel. Population genetic software for teaching and research. *Molecular Ecology Notes* 6: 288–295.
PUENTE-LELIEVRE, C., M. HISLOP, M. HARRINGTON, E. A. BROWN, M. KUZMINA, AND D. M. CRAYN. 2015. A five-marker molecular phylogeny of the Stypheliaceae (Epacridoideae, Ericaceae) supports a broad concept of *Styphelia*. *Australian Systematic Botany* 28: 368–387.
VAN OOSTERHOUT, C., W. F. HUTCHINSON, D. P. M. WILLS, AND P. SHIPLEY. 2004. MICRO-CHECKER: Software for identifying and correcting genotyping errors in microsatellite data. *Molecular Ecology Notes* 4: 535–538.
WESTERN AUSTRALIAN MINISTER FOR THE ENVIRONMENT. 2015. Wildlife Conservation (Rare Flora) Notice 2015. In Western Australian Government Gazette, November 2015. State Law Publisher, Government of Western Australia, Perth, Western Australia, Australia.
WOODMAN ENVIRONMENTAL CONSULTING PTY LTD. 2008. Monitoring of *Leucopogon* sp. ciliate Eneabba (F. Obbens and C. Godden sn 3/7/2003). Unpublished report prepared for Iluka Resources. Woodman Environmental Consulting Pty Ltd., Applecross, Western Australia, Australia.

APPENDIX 1. Locality and voucher information for species used in this study.^a

Species	Voucher specimen accession no.	Collection locality (Population ID)	Geographic coordinates	N
<i>Styphelia longissima</i> Hislop & Puente-Lel.	PERTH 0709170	Eneabba	Rare flora	57
<i>Astroloma xerophyllum</i> (DC.) Sond.	PERTH 8448108	16 km N of Eneabba	–29.683333, 115.483333	5

Note: N = number of individuals used in this study.
^aVouchers are stored in the Western Australian Herbarium (PERTH), Perth, Western Australia, Australia.