

Toxoplasma gondii ANTIBODIES IN FREE-LIVING AFRICAN MAMMALS

Authors: RIEMANN, H. P., BURRIDGE, M. J., BEHYMER, D. E., and Franti, C. E.

Source: Journal of Wildlife Diseases, 11(4) : 529-533

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-11.4.529>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Toxoplasma gondii ANTIBODIES IN FREE-LIVING AFRICAN MAMMALS

H. P. RIEMANN, M. J. BURRIDGE, D. E. BEHYMER and C. E. FRANTI,

Department of Epidemiology and Preventive Medicine, School of Veterinary Medicine,
University of California, Davis, California 95616, USA

Abstract: Twelve species of free-living African mammals from Kenya, Tanzania, Uganda and Zambia were tested for antibodies to *Toxoplasma gondii* using the indirect hemagglutination test. Of 157 animals sampled, 20 (13%) were seropositive. *T. gondii* antibodies were detected in Burchell's zebra, (*Equus burchelli*), hippopotamus (*Hippopotamus amphibius*), African elephant (*Loxodonta africana*), defassa waterbuck (*Kobus defassa*), lion (*Panthera leo*), and rock hyrax (*Procavia capensis*). The highest titers were found in elephants, two having titers of 1:4096 and one of 1:8192. These results are discussed in relation to the maintenance of *T. gondii* among African wildlife.

INTRODUCTION

In a recent study in California, 32% (35 of 109) of the captive exotic animals tested had serologic evidence of exposure to *Toxoplasma gondii*. However, it was not possible to determine if some of these animals had been infected prior to importation into America. It was important, therefore, to examine free-living animals and, in the present study, sera collected from African mammals in their native habitat were tested for antibodies to *T. gondii*.

MATERIALS AND METHODS

Animals

One hundred and fifty-seven serum samples were collected from 12 species of wild mammals in four areas of Africa (Table 1). These areas were as follows: a cattle ranch at Elmenteita in the Rift Valley of Kenya; the northern shore of Lake Victoria in the Busoga District of Uganda;⁷ the Serengeti National Park and the adjacent area of Loliondo, both

in Tanzania; and the Southern Luangwa Valley Game Reserve in Zambia. All animals were shot and then bled immediately for serum collection. Aqueous merthiolate (1:10,000) was added to each serum sample and all sera were stored at -20 C. The sera were heated at 56 C for 1 h prior to shipment to this laboratory.

The ages of the elephant and hippopotamus sampled were recorded as adult, juvenile, or calf, and the sex of every animal tested was noted.

Serology

The serum samples were tested by the microtiter method of the indirect hemagglutination (IHA) test,⁶ using a commercially available antigen.[†] Sera giving a 2+ reaction at a 1:64 dilution were considered positive and were titrated to their endpoints. The 1:32 dilution of serum was used as a nonsensitized cell control and any serum that reacted nonspecifically was omitted from the study.

[†] Industrial Biological Laboratories, Inc., 481 South Ave., Rockville, Maryland 20851, USA.

TABLE 1. Wild mammals studied, by area of capture.

Order	Species	Number sampled by area				
		Kenya	Uganda	Serengeti	Loliondo	Zambia
Camivora	Lion (<i>Panthera leo</i>)			1		1
Proboscidea	African elephant (<i>Loxodonta africana</i>)		1			62
Hyracoidea	Rock hyrax (<i>Procavia capensis</i>)			1		1
Perissodactyla	Burchell's zebra (<i>Equus burchelli</i>)				29	29
	Bushpig (<i>Potamochoerus porcus</i>)		1			1
	Warthog (<i>Phacochoerus aethiopicus</i>)			2		2
	Hippopotamus (<i>Hippopotamus amphibius</i>)					49
Artiodactyla	Defassa waterbuck (<i>Kobus defassa</i>)		2			2
	Impala (<i>Aepyceros melampus</i>)	1				2
	Grant's gazelle (<i>Gazella granti</i>)	1				1
	Thomson's gazelle (<i>Gazella thomsoni</i>)			4		4
	Duiker (undetermined sp.)	1				1
Totals		3	4	8	29	113
						157

RESULTS

Of 157 serum samples tested, 20 (13%) had *T. gondii* antibodies (Table II). The highest prevalence of antibodies was detected among zebra where eight of 29 samples were seropositive. Other species showing *T. gondii* antibodies were hippopotamus (4 of 49), elephant (4 of 63), defassa waterbuck (2 of 2), and lion and rock hyrax (both 1 of 1). The highest titers were found in elephants, two having titers of 1:4096 and one of 1:8192. There was no evidence of *T. gondii* antibodies in either the wild pig or the small antelope species tested.

T. gondii antibodies were detected in wild mammals from all areas studied, except the ranch in Kenya from which only three samples were collected.

All the elephant and hippopotamus sera containing *T. gondii* antibodies were collected from adult animals (for both species, 4 of 39 adults were seropositive). The prevalence of *T. gondii* antibodies was similar for each sex of elephant (1 of 13 males and 3 of 49 females were seropositive), hippopotamus (1 of 18 and 3 of 31), and zebra (6 of 20 and 2 of 9).

TABLE 2. Distribution of HI antibody titers to *Toxoplasma gondii* among African wildlife.

Species	No. tested	No. positive	Reciprocal titer								
			<64	64	128	256	512	1024	2048	4096	8192
Lion	1	1	0	1							
African elephant	63	4	59		1					2	1
Rock hyrax	1	1	0			1					
Burchell's zebra	29	8	21	3	2	2			1		
Bushpig	1	0	1								
Warthog	2	0	2								
Hippopotamus	49	4	45	4							
Defassa waterbuck	2	2	0	2							
Impala	3	0	3								
Grant's gazelle	1	0	1								
Thomson's gazelle	4	0	4								
Duiker	1	0	1								
Totals	157	20	137	10	3	3	0	0	1	2	1

DISCUSSION

Our results show that *T. gondii* is present in wild mammals in three different areas of Africa. Many of the captive exotic animals studied by Riemann *et al.*⁸ were from species indigenous to Africa. Direct comparisons between the African and U.S. groups of animals

cannot be made because different species were studied in each group. There were, however, certain similarities between the groups. For example, about 10% of the Artiodactyla in each group were seropositive. Since these are herbivores, exposure to *T. gondii* would likely be from ingestion of oocysts shed by Felidae rather than by ingestion of cyst-

infected meat.^{2,3,5,10} If one can extend the observations that certain Felidae can serve as definitive hosts,^{4,5} one might postulate that lions and genets could also serve as such hosts and oocyst shedders. The potential ability of wild Felidae to harbor *T. gondii* has been shown by the study of captive exotic animals in California where 75% of the 12 feline species tested were found to have *T. gondii* antibodies.⁸

Another similarity between the wild and the captive groups of animals was the high antibody titers found in some elephants. Three of the seropositive African elephants had titers of 1:4096 or 1:8192, and were the highest found among the 12 species tested. One of the three captive Indian elephants tested and reported in our earlier study,⁸ had a titer of 1:524,288, which was the second highest titer among the 47 captive species studied.

In general, titers for seropositive captive animals were higher than seropositive animals in the present study, the conditional median antibody titer for the seropositive captive animals being 1:256 compared to a median titer of 1:64 for the seropositive African wildlife. High antibody titers are often an indication of active chronic infection. Many investiga-

tors have found that a high frequency of isolations can be made from animals with titers $\geq 1:256$.⁹ However, the parasites also can be present in animals with much lower antibody titers and, in some cases, particularly in birds, the organism can be isolated from individuals with no detectable antibodies.⁵

This study provides evidence that *T. gondii* is being maintained in free-living African wildlife. The organism appears to be maintained among chronically infected herbivores and carnivores, with various wild Felidae amplifying the infection chain among these free-living wild mammals. All of the necessary conditions for the perpetuation of *T. gondii* were present in the areas that were studied. Small species of Felidae such as genets (*Genetta genetta*) were found in all areas, while larger species such as lion were also present in Tanzania and Zambia. These cats probably act as the definitive hosts for the parasite, with herbivores becoming infected by ingestion of oocysts. In all probability, the complete cycle of *T. gondii* can be maintained in Africa by an infection chain that includes the lion and zebra, which have a common predator-prey relationship, as well as in domestic cat-rodent cycle as has been shown in less remote areas.¹

Acknowledgements

We are indebted to Dr. M. P. Cunningham and Mr. C. D. Kimber of the FAO/UNDP Immunological Research on Tick-borne Cattle Diseases and Tick Control Project, East African Veterinary Research Organization, Muguga, Kenya, for provision of the serum samples.

LITERATURE CITED

1. DUBEY, J. P., N. L. MILLER and J. K. FRENKEL. 1970. *Toxoplasma gondii* life cycle in cats. J. Am. vet. med. Ass. 157: 1767-1770.
2. FRENKEL, J. K. and J. P. DUBEY. 1972. Toxoplasmosis and its prevention in cats and man. J. Infect. Dis. 126: 664-673.
3. JACOBS, L. 1961. Toxoplasmosis in man and animals. N.Z. Vet. J. 9: 85-91.
4. JEWELL, M. L., J. K. FRENKEL, K. M. JOHNSON, V. REED and A. RUIZ. 1972. Development of *Toxoplasma* oocysts in neotropical Felidae. Am. J. trop. Med. Hyg. 21: 512-517.
5. MILLER, N. L., J. K. FRENKEL and J. P. DUBEY. 1972. Oral infections with *Toxoplasma* cysts and oocysts in felines, other mammals, and in birds. J. Parasit. 58: 928-937.

6. PARK, H. Y. 1961. *Toxoplasma* hemagglutination test using alcohol-formalin fixed sensitized lyophilized erythrocytes. Arch. Ophthal., Chicago 65: 184-191.
7. REID, H. W., M. J. BURRIDGE, N. B. PULLAN, R. W. SUTHERST and E. B. WAIN. 1970. Survey for trypanosome infections in domestic cattle and wild animals in areas of East Africa. I. Introduction. Br. Vet. J. 126: 622-626.
8. RIEMANN, H. P., D. E. BEHYMER, M. E. FOWLER, T. SCHULZ, A. LOCK, J. G. ORTHOEFER, S. SILVERMAN and C. E. FRANTI. 1974. Prevalence of antibodies to *Toxoplasma gondii* in captive exotic mammals. J. Am. vet. med. Ass. 165: 798-800.
9. WALLACE, G. D. 1973. Intermediate and transport hosts in the natural history of *Toxoplasma gondii*. Am. J. trop. Med. Hyg. 22: 456-464.
10. YILMAZ, S. M. and S. H. HOPKINS. 1972. Effects of different conditions on duration of infectivity of *Toxoplasma gondii* oocysts. J. Parasit. 58: 938-939.

Received for publication 31 March 1975
