



## **BLOOD PARASITES OF SOME BIRDS FROM COLOMBIA**

Authors: BENNETT, GORDON F., and JOSE IGNACIO, BORRERO H.

Source: Journal of Wildlife Diseases, 12(3) : 454-458

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-12.3.454>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

## BLOOD PARASITES OF SOME BIRDS FROM COLOMBIA

GORDON F. BENNETT<sup>[1]</sup> and JOSE IGNACIO BORRERO H.<sup>[2]</sup>

**Abstract:** Blood films from 421 birds of 142 species, representing 29 avian families, from the environs of Cali, Colombia, were examined for blood parasites. Only 30 (7.1%) birds of 26 species harbored hematozoa. Species of *Haemoproteus* (3.1%) and microfilaria (2.3%) were the most commonly encountered blood parasites; species of *Leucocytozoon*, *Plasmodium*, *Akiba* and *Lankesterella* were found in a few birds. Mixed infections with more than one genus of blood parasite were rare; most infections encountered were of low intensity.

### INTRODUCTION

Although the ornithological fauna of South and Central America is the richest and most diverse of any continent, the blood parasites of this avifauna have had only limited study, and then usually for rather specific regions. Recently, Gabaldon and his colleagues<sup>7,8</sup> conducted large-scale surveys of the avian hematozoa of Venezuela and Beltran and his co-workers conducted surveys on the blood parasites of Mexican birds.<sup>2,3</sup> Lainson *et al.*<sup>15</sup> reported on the blood parasites of some birds from Brasil while Jorg<sup>14</sup> reported on the blood parasites of some birds from Argentina. Hewitt,<sup>12</sup> Huff and Wetmore<sup>13</sup> and Galindo and Sousa<sup>9</sup> have all contributed to our knowledge of the avian hematozoa of Central American birds. Few, if any, surveys of avian hematozoa of birds from the western slopes of the Andes have been recorded, including Colombia. Studies conducted by the second author on the distribution of birds on the mountain slopes of the Andes in an area between Cali and Buenaventura Colombia, provided the opportunity to sample some of them for the hematozoan burden. This report summarizes the prevalence of hematozoa in 421 birds of 142 species.

### MATERIALS AND METHODS

Birds were captured in a variety of ways from a number of locations ranging from sea level on the Pacific coast to an altitude of 2140 m on the western slopes of the Cordillera Occidental. Blood smears were air-dried and fixed in 100% methanol or ethanol in Colombia and subsequently stained in Giemsa's (pH 7.2) and examined for blood parasites at the International Reference Centre for Avian Hematozoa. The sample was collected at monthly intervals throughout the period 1971-1972, with larger samples collected in May and October, 1971 and January, 1972. The birds were not aged at time of collection. Most of the birds sampled represent indigenous species.

### RESULTS AND DISCUSSION

A total of 421 birds of 142 species, representing 29 avian families, were examined for blood parasites (Table 1). Only 30 (7.1%) birds of 26 species were found to harbor hematozoa. The most commonly occurring blood parasite genus was *Haemoproteus* (3.1%), followed by microfilaria<sup>2</sup> (2.3%); other parasite genera represented in small numbers were *Leucocytozoon*, *Plasmodium*, *Akiba* and *Lankesterella*. No infections with *Trypanosoma* were found. Most of the infec-

[1] Department of Biology and International Reference Centre for Avian Hematozoa, Memorial University of Newfoundland, St. John's, Newfoundland, Canada.

[2] Departamento de Biología, Universidad del Valle, Cali, Colombia.

TABLE 1. Blood parasites of birds from Colombia.

Bird Family and Species	Birds examined		L.	H.	Pl.	M.	A./La.
	Total	Infected					
COLUMBIDAE							
<i>Columbigallina talpacoti</i>	22	2		1	1		
CUCULIDAE							
<i>Crotophaga ani</i>	1	1		1			
GALBULIDAE							
<i>Galbula ruficauda</i>	1	1		1			
PARULIDAE							
<i>Basileuterus rivularis</i>	2	1					A
ICTERIDAE							
<i>Psarocolius angustifrons</i>	1	1			1		
PIPRIDAE							
<i>Chloropipo holochlora</i>	2	2		2			
PSITTACIDAE							
<i>Pionopsitta haematotis</i>	1	1				1	
<i>Pionopsitta pulchra</i>	1	1				1	
THRAUPIDAE							
<i>Chlorothraupis olivacea</i>	2	1		1			
<i>Tachyphonus delatrii</i>	10	2		1			La.
<i>Tangara nigro-cincta</i>	3	1		1			
<i>Tangara ruficervix</i>	1	1	1				
<i>Tangara xanthogastra</i>	5	1				1	
<i>Thraupis palmarum</i>	3	1				1	
<i>Thraupis virens</i>	3	1		1		1	
TROCHILIDAE							
<i>Coeligena wilsoni</i>	1	1				1	
<i>Eutoxeres aquila</i>	7	2		1		1	
TROGLODYTIDAE							
<i>Thryothorus nigricapillus</i>	2	1					A
TURDIDAE							
<i>Catharus ustulatus</i>	6	1	1			1	
<i>Myadestes ralloides</i>	3	1				1	
<i>Turdus serranus</i>	1	1	1				
TYRANNIDAE							
<i>Contopus virens</i>	2	1		1			
<i>Mionectes olivaceous</i>	5	1		1			
<i>Myiozetetes cayanensis</i>	9	1		1			
<i>Schiffornis turdinus</i>	3	1			1		
<i>Tyrannus melancholicus</i>	3	1				1	
NEGATIVE BIRDS (see below)	391						
TOTAL:	421	30	3	13	3	10	3
PERCENT INFECTED:		7.1	0.7	3.1	0.7	2.3	0.7

NOTE: L. = *Leucocytozoon*; H. = *Haemoproteus*; Pl. = *Plasmodium*; M. = *Microfilaria*;  
 A. = *Akiba*; La. = *Lankesterella*.

TABLE 1 (Continued)

NEGATIVE BIRDS (number examined in parentheses): ACCIPITRIDAE: *Buteo magnirostris* (1); *B. platypterus* (1). ARDEIDAE: *Bubulcus ibis* (55); *Butorides striatus* (8). BUCCONIDAE: *Galbula ruficauda* (2); *Hapaloptila castanea* (1); *Monasa morphoeus* (1); *Notharcus tectus* (1). CAPITONIDAE: *Ebucco bourcierii* (1). CAPRIMULGIDAE: *Nyctodromus albicollis* (1). COLUMBIDAE: *Columba goodsoni* (2). COTINGIDAE: *Lipaugus unirufus* (1); *Pachyramphus cinnamomeus* (3); *Rytipterna holerythra* (1); *Tityra semifasciata* (1). CUCULIDAE: *Coccyzus pumilus* (2). DENDROCOLAPTIDAE: *Dendrocincla fuliginosa* (2); *Dendrocolaptes picumnus* (2); *Glyphorhynchus spirurus* (6); *Lepidocolaptes affinis* (1); *Xiphorhynchus erythropygius* (2). FORMICARIIDAE: *Hylophylax naevioides* (1); *Microrhopias quixensis* (1); *Myrmotherula surinamensis* (1); *Thamnophilus multistriatus* (1); *T. punctatus* (1). FRINGILLIDAE: *Arremon aurantirostris* (3); *Atlappetes brunneinucha* (2); *Oryzoborus angolensis* (4); *O. crassirostris* (1); *Passerina cyanea* (1); *P. cyanoides* (1); *Pitylus grossus* (4); *Saltator albicollis* (3); *S. atripennis* (2); *S. maximus* (1); *Spinus psaltria* (12); *Sporophila americana* (3); *S. intermedia* (2); *S. luctuosa* (2); *S. minuta* (19); *S. nigricollis* (2); *S. obscura* (2); *Tiaris olivacea* (1); *Volatinia jacarina* (15); *Zonotrichia capensis* (1). FURNARIIDAE: *Automolus ochrolaemus* (2); *Syndactyla subalaris* (2); *Synallaxis brachyura* (1); *Thripadectes ignobilis* (1). HIRUNDINIDAE: *Stelgidopteryx ruficollis* (4). ICTERIDAE: *Agelaius icterocephalus* (1); *Cacicus uropygialis* (1). PARULIDAE: *Basileuterus coronatus* (2); *B. nigrocristatus* (1); *Coereba flaveola* (4); *Dendroica petechia* (1); *Geothlypis semiflava* (1); *Oporornis philadelphia* (2). PHASIANIDAE: *Odontophorus hyperythrus* (1). PICIDAE: *Chrysomitris punctigula* (1); *Picumnus granadensis* (2); *Veniliornis fumigatus* (1). PIPRIDAE: *Manacus manacus* (9); *M. vitellinus* (1); *Pipra coronata* (4). PSITTACIDAE: *Aratinga wagleri* (3); *Forpus conspicillatus* (4); *Pionus menstruus* (1). RAMPHASTIDAE: *Aulacorhynchus haematopygius* (1); *Ramphastos swainsoni* (1). STEATORNITHIDAE: *Steatornis caripensis* (2). SYLVIIDAE: *Microbates cinereiventris* (1). THRAUPIDAE: *Chlorospingus flavigularis* (1); *Chlorothraupis stolzmanni* (1); *Habia cristata* (1); *Mitrospingus cassinii* (2); *Pipraeidea melanonota* (1); *Ramphocelus dimidiatus* (1); *R. flamigerus* (8); *Tachyphonus rufus* (1). TROCHILIDAE: *Agelaiocercus kingi* (1); *Amazilia franciae* (1); *Coeligena coeligena* (1); *C. wilsoni* (1); *Metallura tyrianthina* (1); *Phaethornis guy* (2); *P. longuemareus* (1); *P. syrmatophorus* (1); *P. yaruqui* (5); *Thalurania furcata* (2); *Threnetes ruckeri* (2). TROGLODYTIDAE: *Cyphorinus thoracicus* (1); *Microcerculus marginatus* (1); *Troglodytes aedon* (5). TURDIDAE: *Microbates cinereiventris* (1); *Turdus ignobilis* (8). TYRANNIDAE: *Camptostoma obsoletum* (1); *Colonia colonus* (1); *Contopus* sp. (1); *Elaenia flavogaster* (6); *Empidonax virescens* (2); *Leptopogon rufipectus* (1); *L. superciliosus* (2); *Myiobius atricaudus* (1); *M. miniatus* (1); *M. villosus* (2); *Myiarchus tyrannulus* (1); *Myiobius ornatus* (1); *Phaemoyias murina* (7); *Pitangus sulphuratus* (2); *Phynchocyphus brevirostris* (1); *Pyrocephalus rubinus* (1); *Todirostrum cinereum* (1); *Tyranniscus viridiflavus* (1). VIREONIDAE: *Vireo olivaceus* (1).

tions were extremely light, frequently represented by only one or two parasites per smear. Such a low intensity precluded specific identification of most of the hematozoans. Among the haemoproteids, infections of *Haemoproteus maccallumi* and *H. archilocus* in a columbid and a trochilid respectively, were noted. Several haemoproteids were noted which did not correspond to any known

species but taxonomic evaluation of these species is beyond the scope of this report. Two thrushes harbored *Leucocytozoon dubreuilii* and a thraupid harbored *Leucocytozoon fringillinarum*; one tyrannid was infected with *Plasmodium relictum*. The monotypic genus *Akiba* has so far been limited to galliforms in South-East Asia. Two undoubted infections of *Akiba* in a parulid and a wren (Table 1)

TABLE 2. Altitudinal distribution of birds infected with blood parasites.

Altitude metres	Total birds examined	Total birds infected	
		Number	percent
0	43	0	
100	80	11	13.8
1000	52	0	
1035	75	2	2.7
1120	59	7	11.9
1220	57	1	1.8
1845	13	5	38.5
2140	14	1	7.1

were diagnosed—parasites differing from *Akiba cauleryi* in a number of ways. However, lack of material adequate for taxonomic treatment precludes their further consideration at this time.

The birds sampled were collected at various altitudes from sea level to 2140 m up the mountain slopes. The distribution of infected birds (Table 2) was highly variable and no trend could be determined.

The most striking aspect of this survey is the paucity of hematozoan infections in the avian population. This generally low prevalence is in accord with the results of other workers in South and Central America,<sup>2,3,8,9,12,13,15</sup> but in sharp contrast with the prevalence of blood parasites in North America<sup>10</sup> and in Africa<sup>4,5</sup> and Asia.<sup>6,11</sup> One would anticipate that the prevalence of blood parasites in the unique but rich and diversified avifauna of tropical Central and South America would be at least equal to that noted in similar tropical areas of Africa and Asia. The exceptionally low prevalence recorded is difficult to explain with complete satisfaction. The fact that some blood parasites do occur indicates that there

are no physiological barriers in the bird population. One is forced to conclude, therefore, that the low prevalence is the result of a dearth of suitable vector species. This is particularly surprising for *Plasmodium*, a genus generally believed to be plentiful in the wet tropical regions—a belief predicated on the fact that the culicine fauna of such regions is both rich and diverse. It is of interest to note that all three birds harboring *Leucocytozoon* were collected at elevations of 1845–2140 m. This situation might be anticipated if it is considered that the simuliid vectors of this haemosporozoan require rapid running water—a condition more likely to be encountered in mountains. On the other hand, two of the three birds (taken in October–November) were *Catharus ustulatus*, a migratory North American species known to overwinter as far south as Peru and Bolivia.<sup>1</sup> Their infections could have been contracted on their North American breeding grounds. This survey, taken with similar reports from South America indicate that a thorough survey of the species composition and density of the ornithophilic biting fly fauna of the regions is urgently needed.

#### Acknowledgements

The financial support of the National Research Council of Canada to the first author is gratefully acknowledged.

## LITERATURE CITED

1. American Ornithologist' Union. 1957. *The A. O. U. Checklist of North American Birds*. 5th Edition. The American Ornithologist' Union. The Lord Baltimore Press, Baltimore, Maryland. 691 pp.
2. BELTRAN, E. 1939. Investigacion protozoologica de la sangre de 276 aves des mercado en la ciudad de Mexico. *Rev. Inst. Salubr. Enferm. Trop. (Mexico)* 1: 67-72.
3. ——— and A. PARDINAS. 1953. Los protozarios sanguineos de las aves mexicanas. *Mem. Cong. Cien. Mex.* 7: 117-138.
4. BENNETT, G. F., N. O. OKIA and M. F. CAMERON. 1974. Avian hematozoa of some Ugandan birds. *J. Wildl. Dis.* 10: 458-465.
5. ——— and C. M. HERMAN. Blood parasites of some birds from Kenya, Tanzania and Zaire. *J. Wildl. Dis.* 12: in press.
6. DISSANAIKE, A. S., P. NELSON, M. A. FERNANDO and W. J. NILES. 1965. Studies on the haemosporidia of Ceylon birds with special reference to plasmodia. *Ceylon Vet. J.* 13: 65-75.
7. GABALDON, A., G. ULLOA and A. G. de MONTCOURT. 1974. Encuesta sobre Malaria Aviaria en Venezuela; resultados del Primer Ano. *Bol. Dir. Malar. San. Amb.* 19: 80-103.
8. ———. 1975. Encuesta sobre Malaria Aviaria en Venezuela; resultados del Segundo Ano. *Bol. Dir. Malar. San. Amb.* 20: 73-92.
9. GALINDO, P. and O. SOUSA. 1966. Blood parasites of birds from Almirante, Panama with ecological notes on the hosts. *Rev. Biol. (Univ. Costa Rica)* 14: 27-46.
10. GREINER, E. C., G. F. BENNETT, E. M. WHITE and R. F. COOMBS. 1975. Distribution of the avian hematozoa of North America. *Can. J. Zool.* 53: 1762-1787.
11. HEGNER, R. W. and H. J. CHU. 1930. A survey of protozoa parasitic in plants and animals of the Philippine Islands. *Philipp. J. Sci.* 43: 451-482.
12. HEWITT, R. 1940. Studies on blood protozoa obtained from Mexican wild birds. *J. Parasit.* 26: 287-295.
13. HUFF, C. G. and A. WETMORE. 1967. Blood parasites of birds collected in four successive years in Panama. *Bull. Wildl. Dis. Ass.* 3: 178-181.
14. JORG, M. E. 1931. Neuvs protozoos parasitos de la Republica Argentina. *Arch. Soc. Biol. Montevideo* 5 (Suppl.): 1142-1165.
15. LAINSON, R., J. J. SHAW and P. S. HUMPHREY. 1970. Preliminary survey of blood-parasites of birds of the area de pesquisas ecologicas do Guamá, Belém, Pará, Brasil. *Proc. II nd. Int. Cong. Parasitol. in J. Parasit.* 56 (Suppl.): 197-198.

Received for publication 19 February 1976