

Dascyllus SPP.: NEW HOSTS FOR LYMPHOCYSTIS, AND A LIST OF RECENT HOSTS

Authors: LAWLER, ADRIAN R., OGLE, JOHN T., and DONNES, CHAD

Source: Journal of Wildlife Diseases, 13(3): 307-312

Published By: Wildlife Disease Association

URL: https://doi.org/10.7589/0090-3558-13.3.307

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Dascyllus SPP.: NEW HOSTS FOR LYMPHOCYSTIS, AND A LIST OF RECENT HOSTS

ADRIAN R. LAWLER and JOHN T. OGLE, Gulf Coast Research Laboratory, Ocean Springs, Mississippi 39564, USA

CHAD DONNES, Silent World, 317 Huey Long, Gretna, Louisiana 70053, USA

Abstract: Lymphocystis disease is reported for the first time from the white-tailed damselfish, Dascyllus aruanus, and the black-tailed humbug, Dascyllus melanurus. Both species are commercially important exotic aquarium fish from the Indo-Australian Pacific. Lymphocystis is recorded again from the copper banded angelfish, Chelmon rostratus. An updated host list for the disease is included.

INTRODUCTION

Lymphocystis is an infectious viral disease of freshwater and marine teleosts. The virus causes hypertrophy of connective tissue cells resulting in macroscopic nodules often located on the fins and skin. Lymphocystis disease usually is non-fatal, and the nodules eventually disappear; there is no known treatment. Many additional fishes have been reported with lymphocystis (Table 1) since the summary by Nigrelli and Ruggieri.21 To present, lymphocystis infects 96 species of fish belonging to 32 families and 6 orders. The unconfirmed observation by Dawson⁶ would add one species, one family, and one order to the list.

CASE REPORTS

1. Dascyllus aruanus (Linnaeus), white-tailed damselfish

Lymphocystis has been observed routinely on this fish, imported from the Philippines through California, for the past two years. Two individuals were placed in a 300 l tank at GCRL on 14 August 1976. The tank was already occupied by three clown anemone, Amphiprion ocellaris, one high-hat, Equetus

acuminatus, and three peppermint shrimp, Lysmata wurdemanni; none of the fish showed any evidence of lymphocystis.

The two fish were moved to a 20 l tank (A) and five additional specimens were obtained from the same sources on 29 August 1976 and placed in another 20 1 tank (B). One of the five (TL = 47 mm) had a lymphocystis infection. Microscopic examination revealed nodules on all the fins, on top of the head, and on both sides of the body. No nodules were found on the gills or eyes, or internally. Viral particles were confirmed by electron microscope studies (Harold D. Howse, GCRL, personal communication). No parasitic dinoflagellates, Amyloodinium ocellatum (Brown), were found on the gills or skin; no internal parasites were found (Ronnie Palmer, GCRL, personal communication).

The two fish in tank A (34 ppt) and the four remaining in tank B (30 ppt) were anesthetized with MS-222, examined for lymphocystis nodules (none found), and injected on both sides of the body plus the bases of the dorsal, and caudal fins with fresh tissue obtained from the fish infected with lymphocystis.

Tricaine methanesulfonate, Crescent Research Chemicals Inc., Scottsdale, Arizona.

TABLE 1. Lymphocystis-infected fish reported since Nigrelli and Ruggieri:²¹ Spontaneous (S), aquarium (A), experimental (E), and tissue culture (TC) infections.

Host	Type of infection	
CLUPEIFORMES		
CLUPEIDAE		
Clupea harengus var. membras Linnaeus	S	1
LOPHIIFORMES		
OGCOCEPHALIDAE		
Ogcocephalus nasutus (Valenciennes)	?	8
GADIFORMES		
OPHIDIIDAE		
Gunterichthys longipenis Dawson	S	6*
PERCIFORMES		
CENTROPOMIDAE		
Centropomus undecimalis (Bloch	S	9
SERRANIDAE		
Doderleina berycoides Hilgendorf	?	8
LUTJANIDAE		
Lutjanus griseus (Linnaeus)	S	16
SPARIDAE		
Archosargus probatocephalus (Walbaum)	S	23
SCIAENIDAE		
Bairdiella chrysura (Lacépède)	S	7, 13, 14, 15, 27
	E TC	15 27, 28
Cynoscion arenarius Ginsburg	S	4, 7, 10, 13
	\mathbf{E}	5
Cynoscion nothus (Holbrook)	S	25
Cynoscion regalis (Bloch & Schneider)	S	24, 25
Larimus fasciatus (Holbrook)	E	14, 15
Micropogon undulatus (Linnaeus)	S E	4, 5, 10 5, 11
Pogonias cromis (Linnaeus)	E	5
Stellifer lanceolatus (Holbrook)	S	24, 25
CHAETODONTIDAE		
Chaetodon auriga Forskal	A	18
Chaetodon lunulatus Quoy & Gaimard	A	18
Chaetodon miliaris Quoy & Gaimard	A	18
Chelmon rostratus (Linnaeus)	A	18, present report
Pomacanthus annularis (Bloch)	Α	18

TABLE 1—continued

Host	Type of infection	Authority
CICHLIDAE		
Apistogramma ramirezi Myers & Harry	?	8
Haplochromis sp.	S	22
Haplochromas elegans	S	22
Pterophyllum scalare (Lichenstein)	?	8
Tilapia amphimelas	S	22
Tilapia esculenta	S	22
Tilapia variabilis	S	22
POMACENTRIDAE		
Dascyllus aruanus (Linnaeus)	Α	present report
Dascyllus melanurus Bleeker	Α	present report
GRAMMIDAE		
Gramma loreto Poey	Α	12
LABRIDAE		
Bodianus mesothorax (Schneider)	Α	18
CLINIDAE		
Acanthemblemaria crockeri Beebe & Tee-Van	S	17
BLENNIIDAE		
Hypsoblennius gilberti (Jordan)	Α	17
GOBIIDAE		
Gillichthys seta (Ginsburg)	S	19
Gobiosoma paradoxum (Gilbert)	S	19
SCORPAENIDAE		
Sebastes constellatus (Jordan & Gilbert)	Α	18
Sebastes maliger (Jordan & Gilbert)	A	18
Sebastes nebulosus Ayres	Α	18
SIGANIDAE		
Lo vulpinus (Schlegel & Müller)	Α	12, pers. comm.
Lo varpinas (Schieger & Munci)	21	12, pers. comm.
LEURONECTIFORMES		
BOTHIDAE		
Paralichthys dentatus (Linnaeus)	Α	26, pers. comm.
Paralichthys lethostigma Jordan & Gilbert	Α	26
PLEURONECTIDAE		
Pseudopleuronectes americanus (Walbaum)	S	20

TABLE 1-continued

Host	Type of infection		Authority
TETRAODONTIFORMES			
DIODONTIDAE			
Diodon holocanthus Linnaeus	S	19	
Diodon hystrix Linnaeus	E	19	

^{* &}quot;...apparently infected with lymphocystis disease" (Dawson, p. 206). Presently unconfirmed.

The four fish in tank B died on 9, 24, and 27 September 1976 from massive ciliate infections; the ciliates provisionally were identified as *Cryptocaryon irritans* Brown. None of these fish showed any signs of lymphocystis; however, large areas of hyperplastic epidermal tissue were associated with the ciliates. By 21 October 1976, the two fish in tank A had not shown signs of lymphocystis.

On 31 October 1976 we obtained another fish which had lymphocystis nodules on the fins and skin. Thus, two of eight fish examined were infected.

2. Dascyllus melanurus Bleeker, blacktailed humbug

Lymphocystis also has been observed on this species over the past two years. Two fish were placed in the same 300 l tank as the original two *D. aruanus* on 14 August 1976. Nine days later we observed white-colored lesions on the fins and skin of both fish; lymphocystis was verified microscopically, and viral particles were confirmed by electron microscope studies (Harold D. Howse, GCRL, personal communication).

These two fish were moved to tank A with the two D. aruanus, and on 1 September 1976 they were anesthetized with MS-222, and the nodules removed with the aid of forceps and a dissecting microscope. Both fishes (36, 48 mm) had nodules on all fins and on both sides of the

body. The fish were not examined internally, but were revived and placed in a 600 l tank for further observation.

On 17 September 1976 one of the two fish from which the nodules were removed was noticed to have more nodules than previously. This infection is presumed to be a continuation of the initial infection, and not a re-infection. On 25 October 1976 this fish had one small lesion on the dorsal fin; the remainder had sloughed. Thus, in this case, the course of the disease in *D. melanurus* was approximately eight weeks.

3. Chelmon rostratus (Linnaeus), copper banded angelfish

Six fish received in September 1976 were examined microscopically and lymphocystis nodules were present on the fins and skin of four.

DISCUSSION

The only other cases of lymphocystis in the family Pomacentridae were from the spine-cheeked anemone fish, Amphiprion (Premnas) biaculeatus (Bloch), and the percula anemone fish, Amphiprion percula (Lacepede)³ (probably A. ocellaris; see Allen, p. 268). We report two additional hosts from this family. Lymphocystis was previously reported from Chelmon rostratus¹⁸ and is recorded again.

Acknowledgments

W. Jeff Tolbert provided some technical assistance.

LITERATURE CITED

- ALLEN, G. R. 1972. The Anemonefishes. Their Classification and Biology. T. F. H. Publications, Inc. Ltd., Hong Kong, 288 p.
- ANEER, G. and O. LJUNGBERG. 1976. Lymphocystis disease in Baltic herring (Clupea harengus var. membras L.). J. Fish Biol. 8: 345-350.
- BENISCH, J. 1937. Uber das Auftreten der Lymphocystis-Krankheit bei einigen Korallenfischarten. Wochenschr. f. Aq. und Terrarienkunde 34: 380-382.
- CHRISTMAS, J. Y. and H. D. HOWSE. 1970. The occurrence of lymphocystis in Micropogon undulatus and Cynoscion arenarius from Mississippi estuaries. Gulf Res. Rep. 3: 131-154.
- COOK, D. W. 1972. Experimental infection studies with lymphocystis virus from Atlantic croaker. Proc. 3rd Annual Workshop, World Mariculture Society, St. Petersburg, Florida, January 26-28, 1972: 329-335.
- DAWSON, C. E. 1966. Gunterichthys longipenis, a new genus and species of ophidioid fish from the northern Gulf of Mexico. Proc. Biol. Soc. Wash. 79: 205-214.
- DUKES, T. W. and A. R. LAWLER. 1975. The ocular lesions of naturally occurring lymphocystis in fish. Can. J. Comp. Med. 39: 406-410.
- HARSHBARGER, J. C. 1973. Activities report registry of tumors in lower animals. Smithsonian Institution, Washington, D.C. 141 p.
- HOWSE, H. D. 1972. Snook (Centropomus: Centropomidae): New host for lymphocystis, including observations on the ultrastructure of the virus. Amer. Midl. Nat. 88: 476-479.
- and J. Y. CHRISTMAS. 1970. Lymphocystis tumors: Histochemical identification of hyaline substances. Trans. Am. Micro. Soc. 89: 276-282.
- and ________ 1971. Observations on the ultrastructure of lymphocystis virus in the Atlantic croaker, Micropogon undulatus (Linnaeus). Virology 44: 211-214.
- 12. KINGSFORD, E. 1975. Treatment of Exotic Marine Fish Diseases. Palmetto Publishing Company, St. Petersburg, Florida. 90 p.
- LAWLER, A. R. and T. W. DUKES. 1973. Lymphocystis in the eye. The Vet. Rec. 93: 297.
- 14. ———, H. D. HOWSE and D. W. COOK. 1974. Silver perch, Bairdiella chrysura: New host for lymphocystis. Copeia 1974: 266-269.
- 15. ———, ——— and ————. 1975. Lymphocystis infections in the silver perch, *Bairdiella chrysura*. J. Miss. Acad. Sci. 19 (1973-1974): 183.
- LOPEZ, D. M., M. M. SIEGEL, A. R. BEASLEY and L. S. DIETRICH. 1969. Biochemical and morphologic studies of lymphocystis disease. Nat. Cancer Inst. Monograph 31: 223-236.
- 17. McCOSKER, J. E. 1969. A behavioral correlate for the passage of lymphocystis disease in three blennioid fishes. Copeia 1969: 636-637.
- M. D. LAGIOS and T. TUCKER. 1976. Ultrastructure of lymphocystis virus in the quillback rockfish, Sebastes maliger, with records of infection in other aquarium-held fishes. Trans. Am. Fish. Soc. 1976: 333-337.
- and R. F. NIGRELLI. 1971. New records of lymphocystis disease in four eastern Pacific fish species. J. Fish. Res. Bd. Canada 28: 1809-1810.
- MURCHELANO, R. A. and D. W. BRIDGES. 1976. Lymphocystis disease in the winter flounder, *Pseudopleuronectes americanus*. J. Wildl. Dis. 12: 101-103.

- NIGRELLI, R. P. and G. D. RUGIERI. 1965. Studies on virus diseases of fishes. Spontaneous and experimentally induced cellular hypertrophy (lymphocystis disease) in fishes of the New York Aquarium, with a report of new cases and an annotated bibliography (1874-1965). Zoologica 50: 83-96.
- 22. PAPERNA, I. 1973. Lymphocystis in fish from East African lakes. J. Wildl. Dis. 9: 331-335.
- 23. SMITH, A. C. and R. L. TAYLOR. 1972. Preliminary results of marine pathology study off the Florida coast. Marine Tech. Soc. J. 6: 51-52.
- SMITH, F. G. 1970. A preliminary report on the incidence of lymphocystis disease in the fish of the Sapelo Island, Georgia, area. J. Wildl. Dis. (Proceedings Annual Conference) 6: 469-471.
- 1973. Lymphocystis disease in Cynoscion nothus, Cynoscion regalis and Stellifer lanceolatus from Georgia estuaries. Proc. 26th Ann. Conf. SE Assoc. Game & Fish Comm., Oct. 22-25, 1972. Knoxville, Tenn., p. 598-608.
- STICKNEY, R. R. and D. B. WHITE. 1974. Lymphocystis in tank-cultured flounder. Aquaculture 4 (1974): 307-308.
- 27. WHARTON, J. H., R. D. ELLENDER, B. L. MIDDLEBROOKS, P. K. STOCKS, A. R. LAWLER and H. D. HOWSE (in press). Gulf teleost cell culture: Characteristics of a cell line from the silver perch, Bairdiella chrysura In Vitro.
- 28. ——, —— and P. K. STOCKS. 1974. In vitro cultivation of cells from the silver perch, Bairdiella chrysura: A substrate for lymphocystis replication, p. 143-151. In: Proceedings of Gulf Coast Regional Symposium on Diseases of Aquatic Animals. R. L. Amborski, M. A. Hood and R. R. Miller (eds). Louisiana State University, Baton Rouge, Louisiana, April 16-17, 1974. Publ. No. LSU-SG-74-05.

Received for publication 1 March 1977