



STUDIES ON PRONGHORN ANTELOPE (*Antilocapra americana*) AS RESERVOIRS OF ANAPLASMOSIS IN MONTANA 1

Authors: JACOBSON, RICHARD H., WORLEY, DAVID E., and HAWKINS, WILLIAM W.

Source: Journal of Wildlife Diseases, 13(3) : 323-326

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-13.3.323>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

STUDIES ON PRONGHORN ANTELOPE (*Antilocapra americana*) AS RESERVOIRS OF ANAPLASMOSIS IN MONTANA¹

RICHARD H. JACOBSON,² DAVID E. WORLEY³ and WILLIAM W. HAWKINS⁴

Veterinary Research Laboratory, Montana State University, Bozeman, Montana 59715, USA

Abstract: Twenty-six pronghorn antelope (*Antilocapra americana*) were collected in an area of eastern Montana where bovine anaplasmosis is enzootic. Their sera were examined for evidence of anaplasmosis by the complement-fixation test. Strong false positive reactions (3+ and 4+ reactors) occurred for 19 of the sera tested; 6 sera were anticomplementary. Inoculation of antelope blood into anaplasmosis-free intact calves did not produce clinical or serological evidence of anaplasmosis, and anaplasma bodies were not found in stained blood smears of antelope or recipient calves.

INTRODUCTION

Although precise data are unavailable, an estimated 50,000 to 100,000 cattle die of anaplasmosis each year.⁹ Probably the greatest economic impact of the disease results from reduced milk production, weight loss and abortion of fetuses by infected cows.⁹ *Anaplasma marginale* is enzootic in the eastern one-third of Montana where 1.1 million cattle reside.¹⁰ Reports from veterinarians and ranchers in Montana indicate the disease is spreading westward into areas previously considered free of anaplasmosis.

The role of wildlife species as reservoirs of *A. marginale* in Montana has not been studied. Natural infections, or the latent carrier state of anaplasmosis have been observed in Columbian black-tailed deer (*Odocoileus hemionus columbianus*);^{3,5,11} crosses of their progeny with mule deer (*O. h. hemionus*) are known reservoirs of *A. marginale* in California.⁴ In a Wyoming study, blood samples from pronghorn antelope (*Antilocapra americanus*), white-tailed deer

(*O. virginianus*) and elk (*Cervus canadensis*) were negative for *A. marginale* when tested in splenectomized bovine calves.⁷ However, one serum pool, representing 35 mule deer was positive for anaplasmosis.⁷ In a subsequent study, none of 31 mule deer in Oregon were latent carriers of the disease.¹² Similarly, 27 elk in Idaho were negative for anaplasmosis.¹⁵ Of 132 American bison (*Bison bison*) from an area enzootic for anaplasmosis, none were positive for *A. marginale*.¹³

Although these reports suggest that natural infections occur infrequently in wild ruminant species, experimental infections of *A. marginale* have been produced in Columbian black-tailed deer,^{2,14} white-tailed deer,¹⁴ mule deer^{2,8} elk, bighorn sheep (*Ovis canadensis canadensis*) and pronghorn antelope.⁸ Clearly, the potential exists for several of these species, which co-inhabit rangeland with cattle in many areas of the western states, to serve as reservoirs of anaplasmosis. Because natural infections with

¹ Supported in part by Contract No. 12-14-100-11, 151 (45) Livestock Research Division, ARS, USDA. Contribution of the Agricultural Experiment Station, Montana State University, Bozeman, Montana, and published as Journal Series No. 706.

² Present address: New York State College of Veterinary Medicine, Cornell University, Ithaca, New York 14853, USA.

³ Address reprint requests to D. E. Worley.

⁴ Present address: Dillon Veterinary Hospital, Dillon, Montana 59725, USA.

A. marginale have not been identified in antelope, the current investigation was designed to determine if pronghorn antelope, which are numerous in eastern Montana, serve as reservoirs. A secondary objective was to extend a previous evaluation² of the complement-fixation test for diagnosis of anaplasmosis in antelope.

MATERIALS AND METHODS

Twenty-six antelope (16 males and 10 females) were collected from Rosebud County, Montana, in an area enzootic for bovine anaplasmosis. From 2 to 4 antelope were shot by personnel of the Montana Department of Fish and Game on each of 8 collection dates over a 13-month period (Table 1). Approximately 50 ml of blood were obtained from each antelope at the collection site, usually by cardiac puncture using a syringe containing 1% heparin solution. An additional 25 ml blood sample was collected and transported to the laboratory on ice. Following clot formation, the serum was removed and stored at -20 C until used in serological studies.

Eight Hereford calves, one for each antelope collection date, served as recipients of antelope blood. A calf was transported to the collection site in a screened enclosure to preclude exposure to natural vectors of anaplasmosis. Within an average of 15 min. after the death of each antelope, the heparinized blood was inoculated subcutaneously into the calf. Following the inoculations of blood from all the antelope on a given date, the calf was returned to the laboratory and was housed in an isolation unit.

A blood sample was obtained from each calf at weekly intervals for the first 3 weeks and thereafter twice weekly until termination of each test approximately 100 days after inoculation of antelope blood. Blood smears also were prepared on each calf bleeding date. Complement-fixation (CF) tests were conducted on antelope and calf sera,¹ and all blood films were stained with Giemsa and were examined by light microscopy for infected erythrocytes.

RESULTS

A summary of the data collected in this study is presented in Table 1. Of 25 antelope serum samples examined, 18 were 4+ reactors, 1 was a 3+ reactor, and 6 were anticomplementary in CF tests using the *A. marginale* antigen preparation supplied by the Agricultural Research Service, Beltsville, Maryland. No anaplasma bodies or any other parasites were observed in any of the antelope blood films.

All of the antelope were examined in the field for ectoparasites which might serve as vectors of *A. marginale*. The only ectoparasites recovered were two specimens of *Dermacentor* sp. (one on each of two antelope which were collected at an 11-month interval). Therefore, no attempts were made to conduct feeding tests on calves to determine if the ticks harbored *Anaplasma* sp.

As shown in Table 1, each of 8 test calves received at least one injection of blood from antelope which were 4+ reactors in the CF test. None of these calves, however, developed anaplasmosis as determined by CF tests on approximately 25 blood samples drawn serially from each calf over the subsequent 100-day period. Also, erythrocytes in blood films made at the time of each calf bleeding did not contain anaplasma bodies. Furthermore, during the test period, none of the calves developed clinical signs of anaplasmosis.

The experimental protocol provided for limited observations on seasonal prevalence of anaplasmosis in antelope. No pattern emerged either in the CF data, which showed 4+ reactivity on all collection dates, or in transmission of the disease from potentially infected antelope to calves.

DISCUSSION

In the study area of eastern Montana, outbreaks of clinical anaplasmosis characteristically occur in the spring when tick populations (primarily *Dermacentor andersoni*) are at their peak, and in the summer concurrent with the deer fly (*Chrysops* spp.) season. It has not been established experimentally that these

TABLE 1. Experimental Design and Complement Fixation Test Results of Sera from Antelope.

Animal No.	Age (yrs.)	Sex	CF Test	Date Collected	Calf no.*
1	Adult	M	4+	18 Apr	I
2	1	M	AC	18 Apr	
3	2	M	3+	18 Apr	
4	Adult	M	4+	23 May	II
5	UD	F	4+	23 May	
6	Adult	F	4+	7 July	III
7	Adult	M	AC	7 July	
8	2	M	4+	7 July	
9	1	M	AC	7 July	
10	Adult	M	4+	15 Aug	IV
11	Adult	M	4+	15 Aug	
12	2	M	4+	15 Aug	
13	Adult	F	4+	20 Sept	V
14	UD	M	4+	20 Sept	
15	Adult	F	ND	20 Sept	
16	Adult	M	4+	20 Sept	
17	<1	F	4+	15 Nov	VI
18	Adult	F	4+	15 Nov	
19	<1	F	4+	15 Nov	
20	<1	M	4+	15 Nov	
21	UD	M	AC	3 Apr	VII
22	1	M	4+	3 Apr	
23	Adult	M	4+	3 Apr	
24	Adult	F	4+	18 May	VIII
25	1	F	AC	18 May	
26	1	F	AC	18 May	

ND = not done; AC = anti-complementary; UD = undetermined.

*Each calf was given 40-60 ml of antelope blood subcutaneously.

ticks and flies are the actual vectors of anaplasmosis in this area. Clinical disease, however, is usually observed in cattle only at times when these potential vectors are at peak population densities. In our study, 4+ reactivity in CF tests on antelope sera occurred throughout the year regardless of the month of collection. Based on the historical annual pattern of anaplasmosis in the study area, it is unlikely that such strong positive serologic responsiveness can be attributed to continuous exposure of antelope to *Anaplasma* antigens throughout the year. One might argue that active or latent infections were responsible for the CF activity. This conclusion, however, seems precluded on the

basis of the consistently negative blood transfer studies.

To facilitate transmission of viable, infective organisms, every precaution was taken to insure rapid transfer of fresh whole blood from potentially infected antelope into test calves. The time interval between death of an antelope and injection of its blood into calves ranged from 5 to 45 min. with a mean of 15 min. Others¹² have observed that *A. marginale* remains viable in heparinized blood obtained from infected cattle for at least 48 h. Therefore, viable organisms should have been transmitted to test calves in our study if antelope harbored either latent or active infections.

Collectively, the data strongly suggest that antelope in our study were not infected with *A. marginale* and that all of the antelope sera, excluding those which were anticomplementary, resulted in false positive CF reactions. Such a

conclusion confirms the observation of others that the CF test as used for detection of bovine anaplasmosis is not useful for determining the status of *A. marginale* infections in antelope."

Acknowledgements

The authors thank Mr. E. J. Sheehan, USDA Serology Laboratory, Bozeman, Montana, for conducting the CF tests and gratefully acknowledge the assistance of the Montana Department of Fish and Game. Thanks are also extended to Mr. Floyd M. Seesee who assisted with laboratory aspects of the study.

LITERATURE CITED

1. AGRICULTURAL RESEARCH SERVICE, U.S.D.A. 1958. A manual for conducting the complement-fixation test for anaplasmosis. ARS, USDA, Washington, D.C.
2. BOYNTON, W. H. and G. M. WOODS. 1933. Deer as carriers of anaplasmosis. *Science* 78: 559-560.
3. ——— and ———. 1940. Anaplasmosis among deer in the natural state. *Science* 91: 168.
4. CHRISTENSEN, J. F. and D. W. McNEAL. 1967. *Anaplasma marginale* infection in deer in the Sierra Nevada foothill area of California. *Am. J. vet. Res.* 28: 599-601.
5. ———, J. W. OSEBOLD and J. R. DOUGLAS. 1962. Bovine anaplasmosis in the coastal range area of California. *J. Am. vet. med. Ass.* 141: 952-957.
6. ———, ———, J. B. HARROLD and M. N. ROSEN. 1960. Persistence of latent *Anaplasma marginale* infection in deer. *J. Am. vet. med. Ass.* 136: 426-427.
7. HOWE, D. L. and W. G. HEPWORTH. 1965. Anaplasmosis in big game animals: Tests on wild populations in Wyoming. *Am. J. vet. Res.* 26: 1114-1120.
8. ———, ———, F. M. BLUNT and G. M. THOMAS. 1964. Anaplasmosis in big game animals: Experimental infection and evaluation of serologic tests. *Am. J. vet. Res.* 25: 1271-1276.
9. McCALLON, B. R. 1976. Anaplasmosis. *J. Dairy Sci.* 59: 1171-1174.
10. MONTANA AGRICULTURAL STATISTICS, Montana Department of Agriculture and Statistical Reporting Service—USDA. 1972. Vol. 14. Helena, Montana.
11. OSEBOLD, J. W., J. F. CHRISTENSEN, W. M. LONGHURST and M. N. ROSEN. 1959. Latent *Anaplasma marginale* infection in wild deer demonstrated by calf inoculation. *Cornell Vet.* 49: 97-115.
12. PETERSON, K. J., T. P. KISTNER and H. E. DAVIS. 1973. Epizootiologic studies of anaplasmosis in Oregon mule deer. *J. Wildl. Dis.* 9: 314-319.
13. ——— and T. O. ROBY. 1975. Absence of *Anaplasma marginale* infection in American bison raised in an anaplasmosis endemic area. *J. Wildl. Dis.* 11: 395-397.
14. RISTIC, M. and A. M. WATRACH. 1961. Studies in anaplasmosis. II. Electron microscopy of *Anaplasma marginale* in deer. *Am. J. vet. Res.* 22: 109-116.
15. VAUGHN, H. W., H. W. RENSHAW and F. W. FRANK. 1976. Survey of anaplasmosis in elk of the Clearwater National Forest (Idaho). *Am. J. vet. Res.* 37: 615-617.

Received for publication 30 July 1976