

# POPULATIONS, MICROHABITAT PREFERENCE AND EFFECTS OF INFESTATION OF TWO SPECIES OF Orthohalarachne (HALARACHNIDAE: ACARINA) IN THE NORTHERN FUR SEAL 1

Authors: KIM, KE CHUNG, and HAAS, VERDA L.

Source: Journal of Wildlife Diseases, 16(1): 45-51

Published By: Wildlife Disease Association

URL: https://doi.org/10.7589/0090-3558-16.1.45

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at <u>www.bioone.org/terms-of-use</u>.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

## POPULATIONS, MICROHABITAT PREFERENCE AND EFFECTS OF INFESTATION OF TWO SPECIES OF Orthohalarachne (HALARACHNIDAE: ACARINA) IN THE NORTHERN FUR SEAL<sup>II</sup>

KE CHUNG KIM and VERDA L. HAAS, The Frost Entomological Museum, Department of Entomology, The Pennsylvania State University, University Park, Pennsylvania 16802 USA, and MARK C. KEYES, Marine Mammal Division, Northwest and Alaska Fisheries Center, National Marine Fisheries Service, Seattle, Washington 98115 USA.

Abstract: All of 116 northern fur seals examined, except black pups (up to 3 months old), had nasal mites, Orthohalarachne attenuata and O. diminuata, with the mean density of 1,808 mites per subadult male, 435 per adult female, 251 per silver pup, and 21.5 per black pup. Only 63% of black pups examined were infested with both mites. Larvae represented as much as 99% of the total mite population (total samples), and the females of both species of Orthohalarachne accounted for more than 90% of the total population of adult mites. The O. attenuata adults inhabited the nasopharynx and O. diminuata adults were found primarily in the lungs. Larvae of both species occupied the mucus-filled turbinates. The heavy infestation with these mites appeared to result in impairment of respiration in fur seals, and could also cause lesions in the lungs and secondary alveolar emphysems, predispose to more serious diseases, or even kill the host animal.

#### **INTRODUCTION**

Northern fur seals (Callorhinus ursinus) (Otariidae: Pinnipedia) are infested with two species of the nasal mites, Orthohalarachne attenuata (Banks, 1910)<sup>2</sup> and O. diminuata (Doetschmann, 1944),<sup>6</sup> in addition to the two ectoparasitic sucking lice. Antarctophthirus callorhini and Proechinophthirus fluctus, which inhabit skin and fur.<sup>11,12</sup> Orthohararachne spp., obligate parasites of the respiratory passages, are known to occur in the two families of Pinnipedia, Otariidae and Odobenidae.<sup>15</sup> In addition to the northern fur seal (type host) O. attenuata has also been known from other fur seals (Arctocephalus doriferus), sea lions (Neophoca cinerea, Zalophus californianus), and the walrus (Odobenus rosmarus).<sup>5</sup> Orthohalarachne diminuata was described initially from the California sea lion (Zalophus californianus) but has been recorded from northern fur seals,<sup>15</sup> other fur seals (Arctocephalus doriferus)<sup>5</sup> and the northern sea lion (Eumetopias jubata).<sup>15</sup>

Although the larvae,<sup>5</sup> nymphs<sup>8</sup> and adults<sup>15</sup> are now well described, published information on the biology<sup>8</sup> and populations of *Orthohalarachne* is very meager.<sup>7,10</sup>

The impact of arthropod parasites, including the nasal mites, have been considered insignificant in the health and population dynamics of marine mammalian hosts. Dunlap *et al.*<sup>7</sup> first observed some erosion and inflammation of

Paper no. 5515, Journal Series of The Pennsylvania Agricultural Experiment Station, University Park, Pennsylvania 16802, USA. This research is a contribution from the Pennsylvania Agricultural Experiment Station Project No. 2324, and was partly supported by the Marine Fisheries Service, NOAA, Northwest Fisheries Center, U.S. Department of Commerce, Seattle, Washington, 98115 USA.

the nasal turbinates and nasopharynx associated with species of Orthohalarachne. However, they concluded that O. attenuata in the northern fur seal was of minor significance since there were no clinical signs of respiratory difficulty and no evidence of occlusion of the respiratory passages. However, the death of two California sea lions was attributed to bronchitis, secondary emphysema and the lung lesions caused by O. diminuata.4 We report the first population study of Orthohalarachne including infestation rate, population densities and structure, and microhabitat preference of the two species, O. attenuata and O. diminuata, in the respiratory passages of northern fur seals. The effects of mite infestation to the host are discussed.

#### **MATERIALS AND METHODS**

Although snout samples from 154 seals were examined, this report is based on 116 complete samples of heads and respiratory organs from subadult males (66), adult females (15), and pups (35) because some samples were not complete. Adult male samples were not available for study. Heads, tracheae and lungs were collected and promptly frozen in June and July (1975-1977) during the annual harvest of surplus males (subadults of 3-4 years old) on the St. Paul Island, Pribilof Islands, Alaska. Samples were kept frozen and transported to the The Pennsylvania State University Ectoparasite Laboratory for examination.

Skulls were cut sagittally and brain cases with brains were removed. The turbinates, nasopharynx, tracheae and lungs were then examined for mites. The location and the number of mites were recorded for each seal. Mites were preserved for subsequent identification to species, stage of development and sex, in 80% ethyl alcohol. Mites from each seal were kept separate. Nasal turbinates were initially digested with 1% trypsin solution,<sup>3,12</sup> and dissolved with hot 10% KOH solution. For preliminary histopathology, two samples were fixed in 10% formalin. Sections were stained with hematoxylin-eosin.

Tests of significance were calculated pairwise for population difference by Mann-Whitney U-statistics (nonparametric). For the analysis a program MINITAB II was used in the Computation Center, The Pennsylvania State University.

#### **RESULTS AND DISCUSSION**

O. attenuata adults are large mites with a wide range of the body length, 0.9 mm -5 mm, due to the great elongation of the opisthosoma typical of gravid adults. O. attenuata nymphs are about 1.2 mm long and the larvae are 1.0 - 1.3 mm long, much longer than the larvae of O. diminuata. The adults of O. diminuata are small (0.6 mm - 0.8 mm long), without an elongated opisthosoma, and the nymphs are about 0.7 mm long.<sup>#</sup> O. diminuata larvae can be distinguished from O. attenuata by their much shorter body length (0.5 - 0.8 mm long).

All 116 seals examined, except black pups (up to 3 months of age, with black coats), were infested with both species of mites. Only 63% of the black pups examined were infested with both mites. The population density of the nasal mites in the respiratory system was unexpectedly high (Table 1). The mean population of Orthohalarachne spp. was 1,808 (range 23 to 8,170) per subadult male (3-4 years old), 435 per adult female (12-15 years old), 251 per silver pup (more than 4 months old with silvery coat after the first molt), and 21.5 per black pup. Of 66 subadult males examined, 47% (or 31) had more than 1,000 mites per seal and 30% had more than 2,000. Only 17% of the subadult males had less than 100 mites each.

Difference in mite population density was significant between subadult males and females at 0.0319, adults (males plus

I ABLE 1. I Island, Alasl standard err	opulation d sa, collected or.	lensity and during the	structure of summers o	<i>Orthonala</i> of 1972-1977	acnne spp. 11 7. Each value	n the respirat( is the avera	ory system o ge number o	i <i>Callornin</i> f individua	us ursınus ( 1 mites per	on St. Faul host±the
	SAMPLE		O. atte	enuata			O. dim	inuata		GRAND
STSOH	SIZE	LARVAE	SHAMYN	ADULTS	TOTAL	LARVAE	NYMPHS	ADULTS	TOTAL	TOTAL
Subadult &	99	555.7	0	30.1	585.8	1,214.8	0	7.7	1,222.5	1,808.2*
•		± 87.0		$\pm 3.3$	$\pm 94.8$	$\pm 199.5$		$\pm 1.4$	$\pm 199.9$	$\pm 257.8$
Adult <b>Q</b>	15	272.4	.1	121.7	394.2	34.4	0	6.9	40.8	435.1*
(cow)		$\pm$ 86.0		$\pm 27.7$	$\pm 106.5$	$\pm 9.7$		$\pm 2.5$	± 11.0	$\pm 111.0$
Silver	9	126.0	0	28.8	166.0	84.8	0	0	84.8	250.8
Pup		$\pm 21.1$		± 4.6	$\pm 23.7$	$\pm 11.2$			$\pm 11.2$	$\pm 26.6$
Black	29		0	7.1	7.1	5.8	0	8.7	14.3	$21.5^{*}$
Pup				$\pm 2.8$	$\pm 2.8$	$\pm 3.8$		$\pm 2.3$	$\pm 5.5$	$\pm 6.3$
*Difference i	s highly sig	mificant.								

females) and black pups at 0.0000, and silver pups and black pups at 0.0002. Difference between adults and silver pups was not significant at P≤0.05. This may be attributed to small sample size of silver pups. The significant difference in mite population density between subadult males and adult females may be caused by age, sex (and size) or both. Several possible factors may be enumerated for future consideration: (a) the behavioral characteristics of subadult males, which usually aggregate closely on separate beaches away from rookeries;16 (b) a higher level of immunity in mature adults due to infestation in the subadult stage<sup>14</sup> and (c) hormonal action.13

Mite population density in the silver pups increased more than 10 times from the low level in the black pups. New-born pups were free of mites. It appears that a full reproductive capacity of mites was already reached in the silver pups.

We confirmed the findings of Dunlap et  $al.^{7}$  that the larvae of O. attenuata were found on the turbinate mucosa but the adults were found attached to the nasopharyngeal mucosa (Fig. 1, 2). Dunlap et al.<sup>7</sup> made no reference to O. diminuata. We found that the larvae in the turbinate scrolls and nasal meatuses filled with mucous material represented both O. attenuata and O. diminuata (Fig. 3). The adult O. attenuata occupied the nasopharynx extending to the choanae. Since they attached to the nasopharyngeal mucosa by the mouthparts, only the elongated opisthosoma of the adult was visible on the surface (Fig. 4). O. diminuata adults were found primarily in the lungs (bronchi and bronchioles) (Fig. 1). It was assumed that some adults of O. diminuata in the turbinates had been newly molted and some adults in the tracheae would have been migrated to the lungs eventually.

During the life cycle, the hexapod larvae of both species are most active, freely motile, and it is the stage when transmission or dispersal could occur among



FIGURE 1. Distribution of *Orthohalarachne* mites in the respiratory system of northern fur seal. The shaded area in the nymph of *O. diminuata* is based on inference.



FIGURE 2. Sagittal section of the skull of northern fur seal (subadult male); A-nostril, B-turbinate mucosa with larval mites, C-nasopharynx with adults of Orthohalarachne attenuata (M).

Journal of Wildlife Diseases Vol. 16, No. 1, January, 1980



FIGURE 3. Larvae of Orthohalarachne mites in the turbinate scrolls; A-O. attenuata, D-O. diminuata.



FIGURE 4. Sagittal section of the nasopharynx of subadult male northern fur seal showing heavy infestation of *Orthohalarachne attenuata* adults (M).

hosts. Development of the larvae requires a long period of time and they may survive for longer periods.<sup>9</sup> The octopod protonymphal and deutonymphal instars, which require no food, have a very short duration, usually 16-20 hours each.<sup>9</sup>

Of the total mites collected from 116 samples, O. attenuata accounted for 46% and O. diminuata represented 54%. In the subadult males the ratio of O. diminuata to O. attenuata was more than 2 to 1. However, in the adult females and silver pups the density of O. attenuata far exceeded O. diminuata. The population size of mites on black pups was too small to be meaningful. We are not certain of the cause for this difference.

Among adult mites, females represented about 90% of the total; viz., 92% females for O. attenuata and 88% for O. diminuata. Of the total mite population the larvae represented as much as 99%. This suggests a high larval mortality during development. Transmission of these mites from host to host is accomplished by hexapod larvae which may be transferred by nose contact9 or be sneezed from the nostrils of infested animals. Orthohalarachne larvae have been found on hairs of the seal skin on several occasions. In the adult seals expiratory dyspnea or prolonged, forced exhalation is commonly observed. It is not uncommon to see mucous material extruded onto the surface of the nose. It appears that this behavior may be related to the heavy infestation of nasal mites in the turbinates. Unlike sucking lice on the seals<sup>12</sup> there is no mass transfer of mites from the nursing cow to new-born pup. New-born pups are free of the mites.

Preliminary histopathological examination based on heavily infested samples from two subadult males showed that the nasal cavities of heavily infested seals had chronic inflammation of the mucous membranes with erosion, hyperplasia and reactive mononuclear and eosinophilic cellular infiltration. In the lung focal edema and congestion, focal fibrosis and pneumonitis were evident and many bronchioles contained edematous fluid and inflammatory cells.

The heavy infestation with Orthohalarachne mites appeared to result in impairment of respiration, and probably causes expiratory dyspnea due to the blockage of respiratory passages in fur seals, in contrary to the conclusion of Dunlap et al.<sup>7</sup> The heavy mite infestation could also cause lesions in the lungs and secondary alveolar emphysema, predispose to more serious diseases<sup>10</sup> or even kill the host animal.<sup>4</sup> The causes of mortality in fur seal populations are poorly understood because of difficulties in identifying various mortality factors. particularly in the pelagic condition. Hookworms in the fur seal are considered as one of the major causes of mortality in new-born pups. Hookworm larvae occur in the tissues of fur seals of all ages, but adult hookworms occur only in pups.<sup>1,10</sup>

Damage associated with the unexpectedly high level of *Orthohalarachne* infestations must be an important factor affecting the health of northern fur seals particularly in the event that these mites could transmit other pathogens. Thus, the impact of halarachnid mites, as well as that of other internal or external parasites, should be more seriously considered in mortality studies and population analyses of those marine mammals in which they occur.

#### Acknowledgements

We thank George Y. Harry, Alton Y. Roppel and other members, for their field assistance, of the Marine Mammal Division, National Marine Fisheries Service, NOAA, Northwest and Alaska Fisheries Center, Seattle, Washington 98115 and also extend our appreciation to David J. Shetlar and R.G. Anthony of The Pennsylvania State University for their assistance on St. Paul Island, Alaska. We are indebted to Hans Rothenbacher for his histopathological examination, Peter H. Adler for his assistance in statistical analyses, and to B.F. Coon, Deane P. Furman, Hans Rothenbacher, G.L. Storm, S.N. Istvan and M.T. Quinn for reading and making valuable comments on the manuscript.

### LITERATURE CITED

- 1. BAKER, R.C., F. WILKE, and C.H. BALTZO. 1970. The Northern Fur Seal. U.S. Dept. Interior, Fish and Wildl. Serv. Circular 336, 1-19.
- BANKS, N. 1910. New American Mites (Arachnoidea, Acarina). Proc. Entomol. Soc. Wash. 12: 2-12, 3 pls.
- COOK, E.F. 1954. A modification of Hopkins' technique for collecting ectoparasites from mammalian skins. Entomol. News 65: 35-7.
- DAHME, E. and E. POPP. 1963. Todesfalle bei Seelowen (Zalophus californianus Lesson) verursacht durch eine bisher unbekannte Milbe (Orthohalarachne letalis Popp). Berl. und Munch. Tierarztl. Wochensch. 21: 441-3.
- DOMROW, R. 1974. Notes on halarachnine larval morphology and a new species of *Pseudomonyssus* Banks (Acari: Dermanyssidae). J. Aust. Ent. Soc. 13: 17-26.
- 6. DOETSCHMAN, W.H. 1944. A new species of endoparasitic mite of the family Halarachnidae (Acarina). Trans. Ameri. Microsc. Soc. 63: 68-72.
- DUNLAP, S., R.C. PIPER and M.C. KEYES. 1976. Lesions associated with Orthohalarachne attenuata (Halarachnidae) in northern fur seal (Callorhinus ursinus). J. Wildl. Dis. 12: 42-44.
- 8. FURMAN, D.P. 1977. Observations on the ontogeny of halarachnid mites (Acarina: Halarachnidae). J. Parasit. 63: 748-55.
- 9. and A.W. SMITH. 1973. In vitro development of two species of Orthohalarachne (Acarina: Halarachnidae) and adaptations of the life cycles for endoparasitism in mammals. J. Med. Ent. 10: 414-16.
- KEYES, M.C. 1965. Pathology of the northern fur seal. J. Am. vet. med. Ass. 147: 1090-5.
- KIM, K.C. 1971. The sucking lice (Anoplura: Echinophthiriidae) of the northern fur seal; descriptions and morphological adaptation. Ann. Entomol. Soc. 64: 280-92.
- 12. ——. 1972. Louse populations of the northern fur seal (Callorhinus ursinus). Am. J. Vet. Res. 33: 2027-36.
- NELSON, W.A., J.E. KEIRANS, J.F. BELL and C.M. CLIFFORD. 1975. Hostectoparasite relationships. J. Med. Entomol. 12: 143-66.
- 14. ——, J.F. BELL, C.M. CLIFFORD and J.E. KEIRANS. 1977. Interaction of ectoparasites and their hosts. J. Med. Ent. 13: 389-428.
- NEWELL, I.M. 1947. Studies on the morphology and systematics of the family Halarachnidae Oudemans 1906 (Acari: Parasitoidea). Bull. Bingham Oceanogr. Coll. Peabody Mus. Nat. Hist., Yale Univ., New Haven, 10: 235-66.
- PETERSON, R.S. 1968. 1. Social Behavior in Pinnipeds with Particular Reference to the Northern Fur Seal, p. 3-53. In: *The Behavior and Physiology* of Pinnipeds. R.C. Harrison et al. Eds. Appleton-Century-Croft, New York.

**Received** for publication 5 February 1979