

ISOLATION OF ENTEROTOXIGENIC YERSINIA ENTEROCOLITICA FROM BIRDS IN NORWAY

Authors: KAPPERUD, GEORG, and OLSVIK, ØRJAN

Source: Journal of Wildlife Diseases, 18(2) : 247-248

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-18.2.247>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

ISOLATION OF ENTEROTOXIGENIC *YERSINIA* *ENTEROCOLITICA* FROM BIRDS IN NORWAY

GEORG KAPPERUD, Norwegian Defence Microbiological Laboratory, National Institute of Public Health, Geitmyrsveien 75, Oslo 4, Norway.

ØRJAN OLSVIK, Department of Microbiology, Institute of Pharmacy, University of Oslo, Blindern, Oslo 3, Norway.

Recognition of the role of *Yersinia enterocolitica* as a human and animal pathogen has increased in recent years (Mollaret et al., 1979, Contr. Microbiol. Immunol. 5: 174-184). Despite the comprehensive information available on the occurrence of *Y. enterocolitica* and related bacteria in various habitats, the epidemiology of yersiniosis is still hypothetical. This bacterial group has been isolated from numerous animal species including some wild and domestic birds. However, the avian wildlife reservoir has been incompletely investigated.

During May to October 1979, 76 rectal swabs were collected from six avian species at four different localities in Norway. Cultivation was performed on lactose-bromothymol-blue agar and lactose-sucrose-urea (LSU) agar at room temperature for 48 h. Isolated strains were characterized biochemically and antigenically as described previously (Kapperud, 1975, Acta Path. Microbiol. Scand., Sect. B 83: 335-342). All strains were tested for their ability to produce heat-stable enterotoxin at 22 C and 37 C, using the infant mouse assay (Dean et al., 1972, J. Infect. Dis. 125: 407-411). Invasiveness was examined by means of a HeLa cell model (Lee et al., 1977, Can. J. Microbiol. 23: 1714-1722).

Six strains of *Yersinia* were isolated (Table 1) from four (5%) of 76 birds. Each of two birds harbored two antigenically distinct strains. Isolations were obtained from two of six herring gulls (*Larus argentatus*) frequenting a refuse dump heavily contaminated with sewage

sludge. Negative results were obtained from another 10 herring gulls and 12 great black-backed gulls (*Larus marinus*) captured at an ornithologic research station. However, strains were isolated from two of four carrion crows (*Corvus corone*) which were predators of eggs and nestlings within the same area. *Yersinia* was not recovered from 27 willow ptarmigan, (*Lagopus lagopus*) 13 rock ptarmigan (*Lagopus mutus*) and 4 wood pigeons (*Columba palumbus*).

Enterotoxin production at 22 C, but not at 37 C, was indicated for three of the six isolates (Table 1). Invasiveness for HeLa cells was not demonstrated for any of the strains. Enterotoxin production is widespread among *Y. enterocolitica* and *Y. enterocolitica*-like bacteria (Kapperud, 1980, Acta Path. Microbiol. Scand., Sect. B 88: 287-291; Pai et al., 1978, Infect. Immun. 22: 334-338). However, the clinical significance of this enterotoxin in man or animals has not yet been established. Some *Yersinia* strains produce enterotoxin at refrigeration temperature (Kapperud and Langeland, 1981, Curr. Microbiol. 5: 119-122) which is the common storage condition for perishable foods. A possible role of this enterotoxin in food-borne intoxication has been suggested (Boyce et al., 1979, Infect. Immun. 25: 532-537).

Gulls and crows represent a potential source of human infection through fecal contamination of the environment. In this work, *Y. enterocolitica* was isolated from a population of gulls which frequented a refuse dump where sewage sludge was deposited. The same birds

TABLE 1. Characterization of *Yersinia enterocolitica* strains isolated from birds in Norway.

| Strain ^a | Source | Serogroup ^b | Biotype ^c | Enterotoxin ^d production |
|---------------------|--------------|------------------------|----------------------|-------------------------------------|
| K3 | Carrion crow | 6 | 1 | + |
| K4A | Carrion crow | 16 | 1 | — |
| K4B | Carrion crow | 6 | 1 | + |
| K38A | Herring gull | NAG | 1 | + |
| K38B | Herring gull | 17 | 1 | — |
| K42 | Herring gull | NAG | S,SO,C | — |

^aVertical lines indicate strains isolated from the same host.

^bNAG = non-agglutinable.

^cAccording to Wauters' biotype scheme. Strain K42 did not produce acid from sucrose (S), sorbose (SO), nor cellobiose (C).

^dAt 22 C.

breed in areas close to drinking water reservoirs. On the other hand, contact with sewage or sewage sludge may represent a health risk to populations of free-

living birds and mammals. The clinical significance of *Y. enterocolitica* and *Y. enterocolitica*-like bacteria in avian populations is not known.

Received for publication 7 July 1981