

OCCURRENCE OF ANTIBODIES TO THE ETIOLOGIC AGENTS OF INFECTIOUS BOVINE RHINOTRACHEITIS, PARAINFLUENZA 3, LEPTOSPIROSIS, AND BRUCELLOSIS IN WHITE-TAILED DEER IN MINNESOTA

Authors: Ingebrigtsen, David K., Ludwig, John R., and McClurkin, Arlin W.

Source: Journal of Wildlife Diseases, 22(1) : 83-86

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-22.1.83>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

OCCURRENCE OF ANTIBODIES TO THE ETIOLOGIC AGENTS OF INFECTIOUS BOVINE RHINOTRACHEITIS, PARAINFLUENZA 3, LEPTOSPIROSIS, AND BRUCELLOSIS IN WHITE-TAILED DEER IN MINNESOTA

David K. Ingebrigtsen,¹ John R. Ludwig,^{1,3} and Arlin W. McClurkin²

ABSTRACT: A serologic survey was conducted on 628 white-tailed deer (*Odocoileus virginianus*) in 1976 and 1979-1980. Tests for antibodies to the etiologic agents of infectious bovine rhinotracheitis (IBR), parainfluenza 3 (PI3), leptospirosis, and brucellosis produced positive results of 15%, 20%, 3% and 0%, respectively. Adult deer had significantly higher prevalence of antibodies to IBR virus and PI3 virus than fawns. These data provide a basis for monitoring these disease agents in Minnesota's white-tailed deer.

INTRODUCTION

A decline in the productivity of white-tailed deer (*Odocoileus virginianus*) was observed in the farmland area of Minnesota in 1976. Disease, drought, and related stress were suspected as possible factors because adequate habitat was abundant and recent winters had been mild. In conjunction with an intensive study of bovine virus diarrhea (BVD) (Ludwig and McClurkin, unpubl. data), a serologic survey was conducted to determine the occurrence of antibodies in white-tailed deer to the etiologic agents of infectious bovine rhinotracheitis (IBR), parainfluenza 3 (PI3), leptospirosis, and brucellosis. Antibodies to these four disease agents have been reported previously in white-tailed deer in the United States (Table 1). Antibodies to IBR and PI3 viruses also have been reported in moose (*Alces alces*) in Minnesota (Johnson et al., 1972).

Infectious bovine rhinotracheitis is a herpes virus infection of cattle which primarily is a respiratory disease and is known to cause abortion (Richards, 1981). Little

is known of IBR virus in deer. The PI3 virus is a paramyxovirus which has been associated with pneumonia in calves (Richards, 1981). Adult cattle may harbor the virus. Whether cattle transmit the virus to deer is unknown. Leptospirosis is generally transmitted through the urine of carrier animals. Leptospiral organisms survive in moist soil, stagnant ponds, and slow-moving streams (Roth, 1970). Drinking contaminated water is a likely means of transmittal of leptospiral organisms. Leptospirosis has been observed under experimental conditions to cause abortion in white-tailed deer (Trainer et al., 1961). Brucellosis is a widespread disease of man, cattle, goats, and swine, but is found rarely in deer in the United States. It is a cause of abortion in cattle (Witter and O'Meara, 1970). Contact between livestock and wildlife in Minnesota was indicated by the high prevalence (19%-54%) of seropositives to BVD virus in white-tailed deer (Ludwig and McClurkin, unpubl. data).

The objectives of this study were to determine the prevalence of neutralizing antibodies to the etiologic agents of IBR, PI3, leptospirosis, and brucellosis in white-tailed deer in the farmland area of Minnesota. The data would document exposure of deer to bovine diseases and suggest areas for additional research that might explain changes in deer productivity and populations.

Received for publication 18 December 1984.

¹ Farmland Wildlife Populations and Research Group, Minnesota Department of Natural Resources, Route 1, Box 181, Madelia, Minnesota 56062, USA.

² National Animal Disease Center, U.S. Department of Agriculture, P.O. Box 70, Ames, Iowa 50010, USA.

³ Present address: Route 2, Box 981, 639 Thorobred Avenue, Gardnerville, Nevada 89410, USA.

TABLE 1. Summary of selected references on serosurveys of infectious disease agents in white-tailed deer in the United States.

Disease agent	No. positive/no. tested	% Positive	Reference
Infectious bovine rhinotracheitis virus	4/198 12/49	2 24	Friend and Halterman, 1967 Anonymous, 1976
Parainfluenza 3 virus	1/27 9/49	4 18	Shah et al., 1965 Anonymous, 1976
<i>Leptospira interrogans</i>	30/187 585/3,673 3/261 28/285 3/36	16 16 1 10 8	Wedman and Driver, 1957 Ferris et al., 1961 Friend and Halterman, 1967 Haugen, 1967 Fleming and Nusbaum, 1979
<i>Brucella abortus</i>	1/179 20/12,706 1/70	<1 <1 1	Wedman and Driver, 1957 Fay, 1961 Corey et al., 1964

MATERIALS AND METHODS

In 1976, blood samples were collected from hunter-killed deer at 36 mandatory deer registration stations in the farmland area. Blood was collected from the body cavity or visceral organs in 15-ml tubes, centrifuged, and the sera removed and frozen within 24 to 36 hr of collection. The brucellosis test was conducted at the State-Federal Brucellosis Laboratory in St. Paul, Minnesota on the sera collected in 1976 using the standard plate agglutination test (U.S. Department of Agriculture [not dated]) at dilutions of 1:25, 1:50, 1:100, and 1:200. The microscopic-agglutination microtiter test (MAMT) for leptospirosis (Galton et al., 1965) was conducted at the Veterinary Diagnostic Laboratories, University of Minnesota. Sera were tested for antibodies to *Leptospira interrogans* serovars *pomona*, *canicola*, *icterohaemorrhagiae*, *hardjo*, and *grippotyphosa*.

In 1979, a randomly selected sample ($n = 4,200$) of antlerless deer permit recipients was sent prepaid mailers containing a 15-ml plastic syringe, 15-ga needle, centrifuge tube, and instructions on how to collect a blood sample. The hunters were asked to mail the blood to the Minnesota Department of Natural Resources (DNR) at Madelia, where it was centrifuged to obtain serum for IBR and PI3 tests.

Blood specimens were also collected by DNR personnel from deer processed at a sample of mandatory registration stations during the 1979 muzzleloader season and from deer live-trapped at Whitewater Wildlife Management Area (WWMA) during December 1979–February 1980. These sera were kept frozen and subsequently sent to the National Animal Disease Center, Ames, Iowa, for testing for antibodies

to IBR and PI3 viruses. The sera were tested by microneutralization (Ross and Kiesel, 1971) using BT cells (McClurkin et al., 1974) maintained in Gibco (Grand Island Biologic Company, Grand Island, New York 14072, USA) minimum essential media with Earle's salts and 10% virus- and antibody-free bovine serum.

Neutralization endpoints for antibodies to IBR and PI3 viruses were 1:512 or greater, and titers from there to 1:4 were considered to be positive. Data were analyzed by the chi-square test using Yates' correction for the 2×2 tables (Yates, 1934). Correlations were used to compare densities.

RESULTS AND DISCUSSION

Adequate amounts of suitable sera were obtained from 124 deer in 1976, 428 deer during the 1979 regular firearms and muzzleloader seasons, and 77 deer live-trapped at WWMA the following winter. The 418 returns of acceptable serum obtained from the 1979 sample represent a 10% return for the mail survey.

Seropositive results were obtained for IBR virus, PI3 virus, and *Leptospira interrogans*; however, no positive results were obtained for *Brucella abortus* (Table 2). Seropositive titers of 1:100, 1:100, and 1:200 were found for leptospiral serovars *icterohaemorrhagiae*, *pomona*, and *grippotyphosa*, respectively, in each of three deer.

Prevalence of antibodies to IBR virus

TABLE 2. Serologic results for four disease agents in white-tailed deer in the farmland area of Minnesota, 1976–1980.

Disease agent	Years	No. positive/ no. tested	% Posi- tive
Infectious bovine rhinotracheitis virus	1979–1980	74/504	15
Parainfluenza 3 virus	1979–1980	99/504	20
<i>Leptospira interrogans</i>	1976	3/115	3
<i>Brucella abortus</i>	1976	0/124	0

($\chi^2 = 0.02$, $P = 0.90$) and PI3 virus ($\chi^2 = 0.02$, $P = 0.90$) were not different among sexes. However, adults had a significantly higher prevalence of antibody than fawns to IBR virus (18% vs. 8%, $\chi^2 = 8.28$, $P < 0.005$) and PI3 virus (24% vs. 11%, $\chi^2 = 9.78$, $P < 0.002$) (Tables 3, 4) which is probably a result of the increased chance of exposure as deer age or become more mobile. There was no significant difference between the prevalence of antibody of the hunter-killed deer and WWMA live-trapped deer for IBR virus and PI3 virus.

The occurrence of seropositive deer was widespread in the state. Twenty-one of 64 counties with at least one deer tested had

TABLE 3. Serologic results for white-tailed deer older than 1 yr by titer against infectious bovine rhinotracheitis virus and parainfluenza 3 virus in Minnesota, 1979–1980.

Titer	Infectious bovine rhinotracheitis (n = 339)		Parainfluenza 3 (n = 338)	
	No. positive	Percent positive	No. positive	Percent positive
1:4	26	8	25	7
1:8	17	5	16	5
1:16	7	2	7	2
1:32	7	2	8	2
1:64	2	1	5	1
1:128	1	<1	6	2
1:256	0	0	8	2
1:512 or >	1	<1	5	1
Total	61	18	80	24

TABLE 4. Serologic results for white-tailed deer younger than 1 yr by titer against infectious bovine rhinotracheitis virus and parainfluenza 3 virus in Minnesota, 1979–1980.

Titer	Infectious bovine rhinotracheitis (n = 165)		Parainfluenza 3 (n = 166)	
	No. positive	Percent positive	No. positive	Percent positive
1:4	6	4	4	2
1:8	3	2	3	2
1:16	2	1	3	2
1:32	1	1	1	1
1:64	1	1	0	0
1:128	0	0	0	0
1:256	0	0	6	4
1:512 or >	0	0	2	1
Total	13	8	19	11

IBR virus positives, and 30 of 64 counties had PI3 virus positives. In 1976, antibodies to *Leptospira interrogans* were less prevalent (3%) than in a 1957 survey where 16% of white-tailed deer in Minnesota were found to be seropositive (Wedman and Driver, 1957).

Brucellosis is very rare in white-tailed deer. Studies in the United States have documented seropositive results of less than 2% (Wedman and Driver, 1957; Fay, 1961; Corey et al., 1964). Wedman and Driver (1957) found only one positive sample for antibodies to *Brucellosis abortus* in Minnesota.

The present study was conducted to document exposure of white-tailed deer to bovine diseases. Historically, cattle have been prevalent in areas of Minnesota where cash-grain agriculture is not feasible and these areas provide suitable deer habitat. In 1979, deer densities estimated for Minnesota's nine farmland deer management subunits (Schultz, 1984) were correlated ($r = 0.80$, $t = 3.47$, $df = 7$, $P = 0.01$) with cattle densities (U.S. Department of Agriculture, 1980). This suggests that ample opportunity has existed for cattle–deer contact. However, we cannot rule out a density dependent effect among

deer until it is known if these bovine diseases are capable of maintaining themselves within a deer population without exposure of bovine origin.

The positive reactors provide documentation of exposure of white-tailed deer to three bovine pathogens or closely related pathogens specific to white-tailed deer. Even though it has been shown that white-tailed deer have apparently been exposed to these bovine pathogens, very little is known about their actual response to them or their role in transmittal to other species or back to cattle. This study indicated that livestock and wildlife may be exposed to common pathogens and provided baseline data for future studies of diseases of white-tailed deer.

ACKNOWLEDGMENTS

This project was funded by the Section of Wildlife, Division of Fish and Wildlife, Minnesota Department of Natural Resources and the National Animal Disease Center in Ames, Iowa. We would like to thank the hunters, students, and DNR personnel who collected the blood samples. R. Kimmel, R. Lake, D. Heisey, and A. Berner reviewed and J. Lammers typed the manuscript.

LITERATURE CITED

- ANONYMOUS. 1976. Statewide investigations project: Hematology of deer. Arizona Project W-78-R, Job No. 4, Final Report, 9 pp.
- COREY, R. R., L. J. PAULISSEN, AND D. SWARTZ. 1964. Prevalence of brucellae in the wildlife of Arkansas. *Wildl. Dis.* 36: 1-8.
- FAY, L. D. 1961. The current status of brucellosis in white-tailed and mule deer in the United States. *Trans. N. Am. Wildl. and Nat. Resour. Conf.* 26: 203-211.
- FERRIS, D. H., L. E. HANSEN, H. E. RHOADES, AND J. O. ALBERTS. 1961. Bacteriologic and serologic investigations of brucellosis and leptospirosis in Illinois deer. *J. Am. Vet. Med. Assoc.* 139: 892-896.
- FLEMING, W. J., AND S. R. NUSBAUM. 1979. A survey of deer on the Seneca Army Depot for evidence of leptospirosis, brucellosis, and toxoplasmosis. *N.Y. Fish Game J.* 26: 198.
- FRIEND, M., AND L. G. HALTERMAN. 1967. Serologic survey of two deer herds in New York State. *Bull. Wildl. Dis. Assoc.* 3: 32-34.
- GALTON, M. M., C. R. SULZER, C. A. SANTA ROSA, AND M. J. FIELDS. 1965. Application of a microtechnique to the agglutination test for leptospiral antibodies. *Appl. Microbiol.* 13: 81-85.
- HAUGEN, A. O. 1968. Incidence of some disease antibodies in Iowa deer. *Iowa Acad. Sci.* 74: 78-86.
- JOHNSON, D. W., M. WHEELLOCK, B. KOLSTAD, AND P. D. KARNs. 1973. Serologic survey of Minnesota moose for infectious bovine rhinotracheitis (IBR), bovine virus diarrhea (BVD), parainfluenza 3 (PI3), and blue tongue (BT) antibodies. *Proc. Annu. Meet. U.S. Anim. Health Assoc.* 76: 702-710.
- MCCLURKIN, A. W., E. C. PIRTLE, M. F. CORIA, AND R. L. SMITH. 1974. Comparison of low- and high-passage bovine turbinate cells for assay of bovine viral diarrhea virus. *Arch. Virol.* 45: 285-289.
- RICHARDS, S. H. 1981. Miscellaneous viral diseases. In *Diseases and Parasites of White-Tailed Deer*, W. R. Davidson et al. (eds.). Tall Timbers Research Station, Misc. Publ. No. 7, Tallahassee, Florida, pp. 108-125.
- ROSS, C. R., AND G. K. KIESEL. 1971. Microtiter tests for detecting antibody in bovine serum to parainfluenza-3 virus, infectious bovine rhinotracheitis virus, and bovine virus diarrhea virus. *Microbiology* 22: 32-36.
- ROTH, E. E. 1970. Leptospirosis. In *Infectious Diseases of Wild Mammals*, J. W. Davis et al. (eds.). Iowa State Univ. Press, Ames, Iowa, pp. 293-303.
- SCHULTZ, D. F. 1984. Status of wildlife populations, fall 1984 and 1979-1983 hunting and harvest statistics. Unpubl. Rep., Sect. of Wildl., Minn. Dept. Nat. Resour., St. Paul, 94 pp.
- SHAH, K. V., G. B. SHALLER, V. FLYGER, AND C. M. HERMAN. 1965. Antibodies to *Myxovirus parainfluenza 3* in sera of wild deer. *Bull. Wildl. Dis. Assoc.* 1: 31-32.
- TRAINER, D. O., L. KARSTAD, AND R. P. HANSON. 1961. Experimental leptospirosis in white-tailed deer. *J. Infect. Dis.* 108: 278-286.
- U.S. DEPARTMENT OF AGRICULTURE. [not dated]. Standard agglutination test procedures for the diagnosis of brucellosis. *Diagnostic Reagents Manual 65D*. National Veterinary Services Laboratories, Ames, Iowa, 9 pp.
- . 1980. Minnesota Agricultural Statistics, 1980. *Econ. Stat. Coop. Serv.*, 90 pp.
- WEDMAN, E. E., AND F. C. DRIVER. 1957. Leptospirosis and brucellosis titers in deer blood. *J. Am. Vet. Med. Assoc.* 130: 513-514.
- WITTER, J. F., AND D. C. O'MEARA. 1970. Brucellosis. In *Infectious Diseases of Wild Mammals*, J. W. Davis et al. (eds.). Iowa State Univ. Press, Ames, Iowa, pp. 249-255.
- YATES, F. 1934. Contingency tables involving small numbers and the χ^2 -test. *J. R. Stat. Soc. Suppl.* 1: 217-235.