



Prevalence of Parelaphostrongylus tenuis in a Sample of Hunter-Harvested White-tailed Deer from a Tri-County Area in Northeastern Wisconsin

Author: Dew, Terry L.

Source: Journal of Wildlife Diseases, 24(4) : 720-721

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-24.4.720>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Prevalence of *Parelaphostrongylus tenuis* in a Sample of Hunter-Harvested White-tailed Deer from a Tri-County Area in Northeastern Wisconsin

Terry L. Dew, Department of Pathobiological Sciences, University of Wisconsin, School of Veterinary Medicine, 2015 Linden Drive West, Madison, Wisconsin 53706, USA. Present address: Veterinary Specialists of Connecticut, 993 North Main Street, West Hartford, Connecticut 06117, USA

ABSTRACT: Heads from 78 hunter-harvested white-tailed deer (*Odocoileus virginianus*) were donated for study during the 1985 Wisconsin gun hunting season. The intra-cranial nervous tissue was examined for *Parelaphostrongylus tenuis*. Prevalences were calculated for the total sample and for the portion of the sample which had no intra-cranial trauma. The prevalences thus calculated were 58% and 75%, respectively.

Key words: *Parelaphostrongylus tenuis*, white-tailed deer, *Odocoileus virginianus*, prevalence, Wisconsin.

Parelaphostrongylus tenuis is an endemic, relatively innocuous parasite of white-tailed deer (*Odocoileus virginianus*). The parasite is of veterinary importance because of its ability to cause neurologic disease in hoofed species other than white-tailed deer. This nematode infection is becoming increasingly significant in states such as Wisconsin where diverse agri-industry brings numerous susceptible species into contact with a large population of white-tailed deer.

The prevalence of *P. tenuis* infection of white-tailed deer in selected areas of Wisconsin has been reported by other authors. Samuel and Trainer (1969) using post-mortem and fecal analysis reported prevalences of 38% and 15%, respectively. Trainer (1973) reported a prevalence of 50% in a captive free-ranging herd of 600 animals in Vilas County, Wisconsin. Foreyt and Trainer (1980) surveyed deer herds in central and southern Wisconsin and found prevalences of 31% and 28%, respectively.

Seventy-eight specimens of white-tailed deer were collected at two hunter registration stations maintained by the Wisconsin Department of Natural Resources (Madison, Wisconsin 53706, USA). The

deer were harvested from Vilas, Oneida and Forest counties (45°30' to 46°10'N, 88°30' to 89°50'W). The animals' sex was determined, and the heads were disarticulated at the atlanto-occipital joint and stored at -17 C until dissections were performed. Frozen heads were sawn sagittally and allowed to thaw for 12 hr. The cerebral hemispheres were removed intact and the meningeal and dural surfaces were examined for parasites. The deer were aged using dental eruption and tooth wear criteria established by Severinghaus (1949).

Forty-five of 78 specimens harbored *P. tenuis* resulting in a prevalence of 58% (Table 1). Fifty-six of the specimens had no cranial trauma induced by gunshot or trophy collection; 44 (78%) of these were infected. Nematodes were found in only one of 22 damaged heads. The age range of the sample population was 0.5 to 8.5 yr (graded to 0.5 yr), the mean age was 1.9 yr (SD = 1.8). Sixty-five percent of the sample population was female. Chi-square analysis revealed no significant difference

TABLE 1. Specimen classification and the associated prevalences of *Parelaphostrongylus tenuis* in white-tailed deer harvested in northeastern Wisconsin during 1985.

Classification	Number infected/ number examined	Prevalence (%)
Total specimens	45/78	58
Untraumatized specimens	44/56	78
Traumatized specimens	1/22	5
Males	13/27	48
Females	32/51	63
≤2 yr old	33/55	60
>2 yr old	13/23	57

in the prevalences when animals were grouped by sex ($P > 0.5$) or by age (\leq and > 2 yr) ($P > 0.1$).

The prevalence of *Parelaphostrongylus tenuis* in free-ranging white-tailed deer reported here is significantly higher than that previously reported in Wisconsin. While reasons for this may include factors such as geographic differences in population densities of the definitive and intermediate hosts, it is more likely that differences in sampling technique were responsible. Direct identification of the adult parasites eliminates the inherent difficulties associated with the field collection of feces and the definitive identification of metastrongyloid larvae (Gray et al., 1985).

The author thanks Steve Weber for aging the sample population and Gerald Henseler, Jacquelyn Jacobs and Rebecca Stepien for collecting the sample population.

LITERATURE CITED

- FOREYT, W. J., AND D. O. TRAINER. 1980. Seasonal parasitism changes in two populations of white-tailed deer in Wisconsin. *The Journal of Wildlife Management* 44: 758-764.
- GRAY, J. B., W. M. SAMUEL, A. W. SHOSTAK, AND M. J. PYBUS. 1985. *Varestrongylus alpenae* (Nematoda: Metastrongyloidea) in white-tailed deer (*Odocoileus virginianus*) of Saskatchewan. *Canadian Journal of Zoology* 63: 1449-1454.
- SAMUEL, W. M., AND D. O. TRAINER. 1969. A technique for the survey of some helminth and protozoan infections of white-tailed deer. *The Journal of Wildlife Management* 33: 888-894.
- SEVERINGHAUS, C. W. 1949. Tooth development and wear as criteria of age in white-tailed deer. *The Journal of Wildlife Management* 13: 195-216.
- TRAINER, D. O. 1973. Caribou mortality due to the meningeal worm (*Parelaphostrongylus tenuis*). *Journal of Wildlife Disease* 9: 376-378.

Received for publication 8 December 1987.