



## On the Proper Use of Succinylcholine in Elk

Authors: Pybus, M. J., and Onderka, D. K.

Source: Journal of Wildlife Diseases, 28(4) : 685

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-28.4.685>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

## RESPONSE TO LETTER TO THE EDITOR . . .

### On the Proper Use of Succinylcholine in Elk

We appreciate the opportunity to respond to the letter of Cooper et al. regarding the use of succinylcholine as reported in our paper published in the *Journal of Wildlife Diseases*. Please be assured that we were cognizant of our actions at all times. The elk on which we used this drug were all in hard antler at the time (mid-October) and not in velvet. As such it was not necessary to use an analgesic or anaesthetic since there is no invasion of living tissue during the procedure.

In addition, Parks Canada policy has been amended to discontinue use of succinylcholine in any handling procedures in order to avoid having it used inadvertently in a manner contrary to accepted procedures.

We resent the suggestion that our program may have used succinylcholine as a drug of choice due to "cost or ease of use." The paper states clearly the time frame when antlers were removed. In addition, as also stated in the paper (and noted by Cooper et al.), when the procedures involved invasive or prolonged handling, appropriate means of chemical restraint were used.

We trust this matter has been laid to rest.

---

**M. J. Pybus<sup>1</sup> and D. K. Onderka,<sup>2</sup>** <sup>1</sup> Alberta Forestry, Lands and Wildlife, Fish and Wildlife Division, 7th Floor, O. S. Longman Building, 6909-116 Street, Edmonton, Alberta, Canada T6H 4P2; <sup>2</sup> Alberta Agriculture, 1st Floor, O. S. Longman Building, 6909-116 Street, Edmonton, Alberta, Canada T6H 4P2.