

CHARACTERIZATION OF PASTEURELLA MULTOCIDA ASSOCIATED WITH PNEUMONIA IN BIGHORN SHEEP

Authors: Weiser, Glen C., DeLong, Walter J., Paz, Julia L., Shafii, Bahman, Price, William J., et al.

Source: Journal of Wildlife Diseases, 39(3) : 536-544

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-39.3.536>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

CHARACTERIZATION OF *PASTEURELLA MULTOCIDA* ASSOCIATED WITH PNEUMONIA IN BIGHORN SHEEP

Glen C. Weiser,¹ Walter J. DeLong,² Julia L. Paz,³ Bahman Shafii,⁴ William J. Price,⁴ and Alton C. S. Ward^{1,5}

¹ Caine Veterinary Teaching Center, University of Idaho, Caldwell, Idaho 83607, USA

² Idaho Department of Health and Welfare, Bureau of Laboratories, Boise, Idaho 83712, USA

³ University of Health Sciences, Kansas City, Missouri 64124, USA

⁴ Statistical Programs Unit, University of Idaho, Moscow, Idaho 83843, USA

⁵ Corresponding author (email: award@uidaho.edu)

ABSTRACT: *Pasteurella multocida* is a highly diverse group of bacteria recognized as important pathogens. Although *P. multocida* is not ordinarily associated with disease in Rocky Mountain bighorn sheep (*Ovis canadensis canadensis*), numerous isolates were cultured in high numbers from free-ranging bighorn sheep in the Hells Canyon area of Idaho, Washington, and Oregon (USA) during the winter of 1995–96. Animals captured in Hells Canyon and held in captivity, and their offspring, also harbored *P. multocida*. Biochemical utilization tests on 90 isolates identified three subspecies: *P. multocida multocida a* ($n=54$); *P. multocida multocida b* ($n=13$); and *P. multocida gallicida* ($n=15$); and a non-specified biotype, U⁶ ($n=8$). Genomic DNA digestion with restriction endonuclease *Hha* I separated the isolates into 62 unique restriction fragment length polymorphism profiles. Capsular type A was predominant (72% of isolates). Only one isolate type, which may have been transmitted from a feral goat, was capsular type D, possessed the structural gene, *toxA*, for dermonecrotxin detected by polymerase chain reaction, and produced toxin as determined by monoclonal antibody immunoblot. In conclusion, bighorn sheep in this study carried diverse types of generally non-toxicogenic *P. multocida* associated with epizootic pneumonia.

Key words: Capsular type, dermonecrotxin, *Ovis canadensis canadensis*, *Pasteurella multocida*, pneumonia epizootic, polymerase chain reaction, restriction fragment length polymorphism.

INTRODUCTION

Pasteurella multocida, as defined by Rimler and Rhoades (1989), is a highly diverse group of bacteria recognized as important pathogens of animals. These organisms are widespread, colonizing both terrestrial and aquatic birds and mammals (Biberstein, 1979). Since the official recognition of the type strain as *P. multocida* by Rosenbusch and Merchant (1939), taxonomy of this group of organisms has undergone many changes (Mutters et al., 1989; Biberstein et al., 1991) resulting in assignment of isolates previously identified in the species to other species and non-specified biotypes. Currently the species contains three subspecies designations: *P. multocida multocida*, *P. multocida septica*, and *P. multocida gallicida* (Mutters et al., 1989). *Pasteurella multocida multocida* is further divided into two variants, *a* and *b*. The broad complexity of the species involving antigenic variation, host predilec-

tion, and pathogenesis has been reviewed recently (Hunt et al., 2000) and indicates that effective classification should incorporate molecular tools including restriction enzyme analysis and polymerase chain reaction (PCR) procedures.

Each of the subspecies also consists of heterogeneous groups of organisms, some of which are primary pathogens (Christensen and Bisgaard, 2000). Many strains of *P. multocida* have been implicated as causes of severe outbreaks of respiratory disease in cattle, sheep, goats, pigs, and rabbits, atrophic rhinitis in pigs, and fowl cholera (Timoney et al., 1988). Other strains frequently are detected as normal commensals on the mucosa of the upper respiratory tracts of mammals. Many factors such as viral and mycoplasmal infections, poor nutrition, overcrowding, and shipping are associated with reduced physical and immunologic defenses which permit these commensal organisms to invade

TABLE 1. Distribution of *Pasteurella multocida* subspecies and biotype U⁶ isolated from bighorn sheep in association with free-ranging or captive status and health of the animals.

<i>P. multocida</i> subspecies/biotype	Bighorn sheep characteristics and numbers				
	Healthy free-ranging	Healthy captive	Respiratory distress free-ranging	Dead ^a free-ranging	Dead captive
<i>multocida a</i>	2	1	4	20	8
<i>multocida b</i>	1	0	0	4	1
<i>gallicida</i>	0	0	0	7	1
U ⁶	1	0	0	3	0
Toxigenic <i>multocida a</i>	0	0	0	1	0
<i>multocida a</i> + <i>multocida b</i>	0	1	0	6	0
<i>multocida a</i> + <i>gallicida</i>	0	0	1	4	0
<i>multocida a</i> + U ⁶	1	0	0	2	0
<i>multocida b</i> + U ⁶	0	0	0	1	0
<i>multocida a</i> + <i>multocida b</i> + <i>gallicida</i>	0	0	0	2	0

^a All dead sheep showed evidence of lung consolidation at necropsy.

the lower respiratory tract and cause respiratory disease. Some strains appear to be normal commensals of the upper respiratory tracts of wild ruminants (Ward et al., 1997; Jaworski et al., 1998) but are rarely reported to be associated with respiratory disease in North American wild ruminants. Exceptions include hemorrhagic septicemia in North American bison (*Bison bison*) (Rimler and Wilson, 1994), septicemic pasteurellosis in elk (*Cervus elaphus nelsoni*; Wilson et al., 1995), and sporadic cases of respiratory disease affecting small numbers of deer (*Odocoileus hemionus*; Thorne, 1982) and bighorn sheep (Jaworski et al., 1998). Some strains producing hemorrhagic septicemia have characteristic restriction endonuclease profiles (Rimler, 2000). Multiple strains, some of which were shared with wildlife, were isolated from a single turkey flock exhibiting fowl cholera (Snipes et al., 1989). Dermonecrotins, proteins of approximately 112 to 160 kD which exhibit cytotoxic, dermonecrotic, and osteolytic activity, are produced by strains which possess *tox*A genes (Chanter and Rutter, 1989; Ghoshal and Niyo, 1993; Lichtensteiger et al., 1996).

A pneumonia epizootic affecting bighorn sheep in Idaho, Oregon, and Wash-

ington (USA) occurred in Hells Canyon and resulted in the death of 50–75% of the animals in four herds during the winter of 1995–96 (Cassirer et al., 1996). Samples collected by wildlife biologists from dead or dying bighorns were submitted to the Caine Veterinary Teaching Center (Caldwell, Idaho) for bacteriologic analysis. *Pasteurella haemolytica* and *P. trehalosi*, common commensals in bighorn sheep, were isolated from most animals. In addition, multiple strains of *P. multocida*, which are reportedly less common in bighorn sheep, were isolated from a high percentage of the animals. This study was conducted to determine if morbidity and mortality were associated with a single or multiple strains of *P. multocida*, their ability to produce dermonecrotin, or their particular capsular type.

MATERIALS AND METHODS

Pasteurella multocida isolates

Ninety *P. multocida* isolates from oropharyngeal or lung tissue samples of 72 bighorn sheep were evaluated. The condition of the animals (healthy, respiratory distress, or dead) at the time of sample collection was recorded (Table 1). Some of the isolates were from samples collected from live, free-ranging bighorn sheep when they were first noted to be exhibiting signs of respiratory distress and from animals

which had died in their native habitat in Hells Canyon National Recreation Area (46°10'N, 117°00'W) (Cassirer et al., 1996). Other isolates were from live sheep that were captured in Hells Canyon and moved to pens at the Idaho Department of Fish and Game Wildlife Health Laboratory near Caldwell, Idaho during the later stages of the epizootic for monitoring and antibiotic treatment and some of those which subsequently died. Three isolates, 97-1205, 98-879, and 98-1230 were cultured from one of the surviving bighorn ewes and two lambs born in captivity at 22 and 31 mo after first detection of disease in the Hells Canyon population.

Samples from each animal were inoculated onto Columbia blood (5% ovine) agar (CBA) and Columbia blood (5% bovine) agar with antibiotics selective for Pasteurellaceae as previously described (Jaworski et al., 1993). Gram-negative isolates which were oxidase-, indole- and nitrate-positive and urease-negative were tentatively identified as *P. multocida*. Additional biochemical utilization tests were performed to confirm identification and differentiate isolates into subspecies and biotypes using the system of Biberstein et al. (1991). Isolates were subsequently maintained frozen at -70 C in a 60:40 (v/v) phosphate buffered saline (pH 7.2): glycerol cryoprotectant until they were propagated on fresh CBA for evaluation in this study.

Genomic DNA isolation

Bacteria grown on CBA were inoculated into 3 ml of brain-heart infusion (BHI) broth which was then incubated at 37 C with shaking. Genomic DNA was isolated from stationary phase (16 hr) BHI broth cultures containing approximately 5×10^8 colony forming units (cfu) ml⁻¹ using the Puregene® DNA Isolation Kit (Gentra Systems, Minneapolis, Minnesota, USA) and quantified with Hoechst 33258 fluorometry (Labarca and Paigen, 1980).

Restriction fragment length polymorphism analysis

Purified DNA (1.5 µg) was digested with 5 U of *Hha* I restriction endonuclease (Promega Corp., Madison, Wisconsin, USA) for 3 hr at 37 C in a final reaction volume of 20 µl according to manufacturer's directions. Replicate digestates were electrophoresed in 1.0% equivalent agarose-Synergel (Diversified Biotech, Boston, Massachusetts, USA) gels for 16 hr at 30V and stained with ethidium bromide. Gels were photographed on a UV transilluminator and imaged with a Gel Doc® 2000 Gel Documentation System (Bio-Rad Laboratories, Hercules, California, USA) for band analysis.

Data analysis

Imaged banding patterns were used to calculate relationship indices (Nei and Lei, 1979) between all isolates within each subspecies and biotype. Dendrograms were constructed using the unweighted pair-group method using arithmetic averages (UPGMA) cluster analysis (Sneath and Sokal, 1973).

Polymerase chain reaction (PCR) for dermonecrotxin *toxA* gene

Isolates were tested for presence of the dermonecrotxin structural gene, *toxA*, by PCR using the primers and a modified protocol originally designed by Lichtensteiger et al. (1996). Briefly, each 50 µl reaction contained 10 ng of DNA, in MasterAmp® PCR Premix C (Epicentre Technologies, Madison, Wisconsin), 1.5 U RedTaq® DNA polymerase (Sigma Chemical Co., St. Louis, Missouri, USA), and 0.75 mM each primer. Forty cycles, with each cycle consisting of denaturing at 94 C, annealing at 55 C and extension at 72 C for 30 sec each were performed in a thermal cycler (Perkin-Elmer Cetus Corp., Emeryville, California). The resulting PCR products were electrophoresed in 1.1% agarose gels for 45 min at 90V, stained with ethidium bromide (50 ppm) and imaged with a Gel Doc® 2000 Gel Documentation System (Bio-Rad Laboratories) for band analysis.

Immunoblot analysis for dermonecrotxin activity

Crude toxin preparations were made from selected isolates of *P. multocida* as described by Lichtensteiger et al. (1996). Each crude toxin preparation (200 µl) was vacuum filtered onto a nitrocellulose membrane using a Bio-dot apparatus (Bio-Rad Laboratories). Following filtration, the membrane was removed and treated with a blocking buffer (Tris buffered saline [TBS] with 3% gelatin) for 45 min at room temperature. The membrane was washed twice in TBS with 0.05% Tween-20 (TTBS) and incubated at room temperature overnight in a 1:50 dilution in TBS of anti-*P. multocida* dermonecrotxin monoclonal antibody (Hybridoma α-DNT-1B2A3; American Type Culture Collection No. CRL-1965, Manassas, Virginia, USA). Following two additional 5 min washes in TTBS, the membrane was incubated 1 hr in a 1:2,000 dilution of peroxidase-conjugated goat anti-mouse immunoglobulins (Cappel, Organon Teknika, Durham, North Carolina). Subsequently, two washings were performed as above, followed by a third brief wash in TBS. The membrane was then incubated for 15 min in a chromogenic substrate for peroxidase

(3mM 4-chloro-1-naphthol with 0.015% H₂O₂). Finally, the membrane was rinsed in distilled H₂O to stop the reaction and the membrane image recorded digitally (Gel Doc 2000; Bio-Rad Laboratories).

Polymerase chain reaction for capsular type genes

The capsular type of each isolate was determined with multiplex PCR using primers designed to amplify capsule determinant genes by Townsend et al. (2001). Each 50 µl reaction contained 10 ng of DNA, in MasterAmp[™] PCR Premix B (Epicentre Technologies), 1.5 U RedTaq[™] DNA polymerase (Sigma Chemical Co.) and 3.2 µM of each primer. Forty cycles, with each cycle consisting of denaturing at 94 C, annealing at 55 C and extension at 72 C for 30 sec each, followed by a final extension at 72 C for 5 min were performed in a thermal cycler (PerkinElmer Cetus Corp.). The resulting PCR products were electrophoresed in 1.1% agarose gels for 45 min at 90V, stained with ethidium bromide (50 ppm), and imaged with a computer coupled device (Gel Doc 2000, Bio-Rad Laboratories).

RESULTS

Pasteurella multocida multocida a was the predominant subspecies (54/90) isolated from 33 animals including three healthy, four with respiratory distress, and 26 dead bighorn sheep (Table 1). This subspecies was also found in 17 animals in combination with the other subspecies and biotype U⁶. Only three of the animals carrying these combinations were alive at sampling. The 54 isolates presented 29 unique restriction fragment length polymorphism (RFLP) patterns (Fig. 1). Eighty-two percent (44/54) of *P. multocida multocida a* isolates, one of which was dermonecrotogenic, were from dead animals. This dermonecrotogenic isolate (95-1519) was identical to an isolate cultured from a feral goat found near the bighorn sheep in Hells Canyon (K. Rudolph, pers. comm.) and clustered with a dermonecrototoxin-negative *P. multocida multocida a* isolate (95-1508) from an animal with respiratory distress with a similarity coefficient (SC) of 0.78. However, these two isolates were not closely related to the remaining 52 *P. multocida multocida a*




FIGURE 1. Genetic relationships of *Pasteurella multocida multocida a* associated with bighorn sheep in Hells Canyon, 1995–96.

da a isolates based on RFLP profiles. Another cluster of five identical organisms, 4/5 isolated from dead animals, was distantly related to the remaining 47 *P. multocida multocida a* isolates with a similarity coefficient of 0.05. With only two exceptions (96-1103 and 96-997), the isolates from captive animals had identical RFLP patterns or clustered together. Only five isolates were obtained from healthy animals and none of these clustered together.

In contrast, *P. multocida multocida b* was isolated alone from only five dead, free-ranging, or captive animals and only one live animal, and in combination with




FIGURE 2. Genetic relationships of *Pasteurella multocida multocida b* associated with bighorn sheep in Hells Canyon, 1995–96.

the other subspecies or U⁶ biotype in one healthy and nine dead, free-ranging animals, from which *P. multocida multocida a* was also isolated (Table 1). Each of the 13 isolates gave a unique RFLP pattern with SCs ranging from 0.04–0.86 (Fig. 2). Only two *P. multocida multocida b* isolates were from healthy animals (96-432 and 96-997) and these clustered differently. The two isolates from captive animals (96-997 and 97-1100) were unrelated.

All but one (96-75) of the 15 *P. multocida gallicida* isolates was from dead animals and two (96-444 and 96-847) shared identical RFLP patterns (Fig. 3). Only one isolate (97-823) was carried by a captive animal and this had a SC of 0.78 with isolate 96-75 from a live animal showing signs of respiratory distress.

Eight *P. multocida* biotype U⁶ isolates




FIGURE 3. Genetic relationships of *Pasteurella multocida gallicida* associated with bighorn sheep in Hells Canyon, 1995–96.

were found in this study and three (38%) shared identical RFLP patterns (Fig. 4). Two (25%) of these isolates were cultured from apparently healthy animals and these, especially 95-1490, were only distantly related to the rest of the U⁶ isolates. As a group, the *P. multocida* biotype U⁶ isolates were the most similar with seven of eight having SCs of 0.44 or greater.

Overall, 74% (67/90) of isolates carried the capsular type A gene and the remainder were acapsular, with one exception, 95-1519. This dermonecrotizing toxin producing isolate carried the capsular type D gene and was confirmed type D by Dr. Richard Rimler, USDA, Agricultural Research Service (Ames, Iowa, USA). Capsular types (A and acapsular) appeared to be distributed randomly throughout the taxonomic groups (data not shown).

DISCUSSION

The *P. multocida* isolates from lung tissue in this study were cultured in high numbers, suggesting that these organisms




FIGURE 4. Genetic relationships of *Pasteurella multocida* biotype U⁶ associated with bighorn sheep in Hells Canyon, 1995–96.

may be more pathogenic for bighorn sheep than previously recognized. Pasteurellaceae are commonly found among the normal flora of healthy-appearing wild ruminants, with *P. trehalosi* and *Mannheimia haemolytica* being isolated most frequently. In contrast, *P. multocida* is less frequently isolated from healthy wild ruminants (Ward et al., 1997; Jawarski et al., 1998). In the present study, Pasteurellaceae isolated from samples of individual animals were identified as *M. haemolytica*, *P. trehalosi*, and *P. multocida* (15%); *M. haemolytica* and *P. multocida* (1%); *P. trehalosi* and *P. multocida* (28%); or *P. multocida* only (55%), representing extremely high prevalences of the latter species. While respiratory disease in wild ruminant populations is most likely the culmination of several biotic and abiotic factors involved in pathogenesis (Rimler, 2000), *P. multocida* strains have been implicated previously in pneumonia in bighorn sheep and elk, (Thorne, 1982) and hemorrhagic

septicemia in fallow deer (*Dama dama*; Eriksen et al., 1999) and bison.

Isolates in each of the *P. multocida* subspecies and biotype U⁶ comprised multiple RFLP profile groups. Strains of *P. multocida multocida a* were isolated from more bighorn sheep than those in other subspecies and the U⁶ biotype; however, a clearly-defined association with disease was not indicated for any subspecies or RFLP profile group. In addition, no synergistic association between organisms in any of the subspecies or biotype U⁶ appeared to be associated with morbidity or mortality in these animals.

Strains of *P. multocida* that possess the *tox A* gene produce a dermonecrotxin (Lichtensteiger et al., 1996). Infection with toxigenic strains results in progressive atrophic rhinitis in pigs (Chanter and Rutter, 1989; Goshal and Niyo, 1993; Gardner et al., 1994). Goats in Norway have been shown to carry dermonecrotogenic *P. multocida* and develop atrophic rhinitis (Baalsrud, 1987). The role of dermonecrotxin in wild ruminants or its occurrence in *P. multocida* isolated from them has not been well-defined, although our results indicate such occurrences may be rare.

Presence of a capsule is an important virulence factor in *P. multocida* commonly found in domestic animals (Boyce and Adler, 2000). Capsular type A, predominant in the present study, was not particularly associated with any group of bighorn sheep, e.g., dead, live, or captive. Because type A strains are often found associated with avian hosts, these may serve as a reservoir of *P. multocida* in the Hells Canyon area.

Other studies on the genomic RFLP patterns of *P. multocida* (Zhao et al., 1992; Gardner et al., 1994; Donnio et al., 1999; Rimler, 2000) have shown sensitive levels of discrimination and remarkable genetic heterogeneity and are considered especially valuable in epidemiologic studies of pasteurellosis outbreaks (Hunt et al., 2000). Strains selected for analysis in those studies have been mostly derived from

swine herds because of the economic impact of atrophic rhinitis. One study (Donnio et al., 1999) compared isolates from humans and swine, concluding that no apparent differences were discernible and that transmission may occur from pigs to humans working with them on production farms. Few studies on the genetic or phenotypic variability between strains of *P. multocida* isolated from wild ruminant populations have been published. Only one RFLP pattern was found with 70 isolates, all in the same serotype, implicated in pasteurellosis in an elk herd located at the National Elk Refuge (Jackson, Wyoming, USA; Wilson et al., 1995). In another study using six strains of the same serotype isolated from pronghorn (*Antilocapra americana*), moose (*Alces alces*), cattle, and swan each gave a unique RFLP pattern (Dunbar et al., 2000). Results from the present study indicate many unique RFLP patterns from bighorn sheep in the Hells Canyon area of Idaho, Washington, and Oregon.

The diversity of *P. multocida* strains isolated from bighorn sheep in this study is indicative of multiple origins rather than recent transmission from a single source. This is apparent even with *P. multocida multocida a* isolates which were cultured from the majority of the bighorn sheep. Although clusters of *P. multocida multocida a* isolates with identical SC values are suggestive of transmission between bighorn sheep, SC values below 0.8 appear to rule out recent transmission from a single source. The most notable suggestive evidence of transmission includes that of the dermonecrotogenic strain isolated from the feral goat and one of the bighorn sheep found in close proximity. Because this strain was not isolated from any of the other bighorn sheep in the study, it appears that the goat was the most probable source. In another group of animals, identical organisms were isolated from a ewe (97-1205) at the time of her death and the lungs of her two lambs which were euthanized while suffering from acute pneu-

monia 8 (98-879) and 10 (98-1230) months later.

High prevalence of *P. multocida* in bighorn sheep populations during a period of high morbidity and mortality suggests that one or more of the strains may be pathogenic for these animals. Polymorphic DNA patterns and RFLP patterns of PCR amplicons from virulence genes of these isolates are being characterized further for possible use in predicting virulence potential of these organisms for bighorn sheep. Characterizations conducted on strains isolated from animals during health monitoring programs may be useful in averting *P. multocida* epizootics. Wildlife biologists should be aware of the presence of *P. multocida* strains in bighorns and the possible deadly consequences of unchecked transmission within and between bighorn populations.

ACKNOWLEDGMENTS

This research was funded in part by the Idaho Department of Fish and Game, the University of Idaho, Idaho Agricultural Experiment Station Western Regional Research Project W-112 (BGR971) and USDA-CSREES Animal Health and Disease Project IDA-04-AHD98. Idaho Agricultural Experiment Station publication no. 02A05.

We thank D. Hunter and K. Rudolph, technical support personnel L. Cowan and J. Bulgin, plus numerous Idaho Fish and Game Department Biologists for initial sample collection and bacterial isolation.

LITERATURE CITED

- BAALSRUD, K. J. 1987. Atrophic rhinitis in goats in Norway. *The Veterinary Record* 121: 350-353.
- BIBERSTEIN, E. L. 1979. The pasteurelloses. In *Handbook series in zoonoses*, H. Stoenner, M. Torten and W. Kaplan (eds.). CRC Press, Inc., Boca Raton, Florida, pp. 495-514.
- , S. S. JANG, P. H. KASS, AND D. C. HIRSH. 1991. Distribution of indole-producing urease-negative pasteurellas in animals. *Journal of Veterinary Diagnostic Investigation* 3: 319-323.
- BOYCE, J. D., AND B. ADLER. 2000. The capsule is a virulence determinant in the pathogenesis of *Pasteurella multocida* M1404 (B:2). *Infection and Immunity* 68: 3463-3468.
- CASSIRER, E. F., L. E. OLDENBURG, V. L. COGGINS, P. FOWLER, K. RUDOLPH, D. L. HUNTER, AND W. J. FOREYT. 1996. Overview and preliminary anal-

- ysis of a bighorn sheep dieoff, Hells Canyon, 1995–96. Biennial Symposium Northern Wild Sheep and Goat Council 10: 78–86.
- CHANTER, N., AND J. M. RUTTER. 1989. Pasteurellosis in pigs and the determinants of virulence of *Pasteurella multocida*. In *Pasteurella* and pasteurellosis, C. Adlam and J. M. Rutter (eds.). Academic Press, New York, New York, pp. 161–195.
- CHRISTENSEN, J. P., AND M. BISGAARD. 2000. Fowl cholera. Scientific and Technical Review 19: 626–637.
- DONNIO, P.-Y., A. ALLARDET-SERVENT, M. PERRIN, F. ESCANDE, AND J.-L. AVRIL. 1999. Characterisation of dermonecrotic toxin-producing strains of *Pasteurella multocida* subsp. *multocida* isolated from man and swine. Journal of Medical Microbiology 48: 125–131.
- DUNBAR, M. R., M. J. WOLCOTT, R. B. RIMLER, AND B. M. BERLOWSKI. 2000. Systemic pasteurellosis in free-ranging neonatal pronghorn in Oregon. Journal of Wildlife Diseases 36: 383–388.
- ERIKSEN, L., B. AALBAEK, P. S. LEIFSSON, A. BASSE, T. CHRISTIANSEN, E. ERIKSEN, AND R. B. RIMLER. 1999. Hemorrhagic septicemia in fallow deer (*Dama dama*) caused by *Pasteurella multocida multocida*. Journal of Zoo and Wildlife Medicine 30: 285–292.
- GARDNER, I. A., R. KASTEN, G. J. EAMENS, K. P. SNIPES, AND R. J. ANDERSON. 1994. Molecular fingerprinting of *Pasteurella multocida* associated with progressive atrophic rhinitis in swine herds. Journal of Veterinary Diagnostic Investigation 6: 442–447.
- GHOSHAL, N. G., AND Y. NIYO. 1993. Histomorphologic features of the nasal cavity of pigs exposed to *Pasteurella multocida* type-D dermonecrotic toxin. American Journal of Veterinary Research 54: 738–742.
- HUNT, M. L., B. ADLER, AND K. M. TOWNSEND. 2000. The molecular biology of *Pasteurella multocida*. Veterinary Microbiology 72: 3–25.
- JAWORSKI, M. D., A. C. S. WARD, D. L. HUNTER, AND I. V. WESLEY. 1993. Use of DNA analysis of *Pasteurella haemolytica* biotype T isolates to monitor transmission in bighorn sheep (*Ovis canadensis canadensis*). Journal of Clinical Microbiology 31: 831–835.
- , D. L. HUNTER, AND A. C. S. WARD. 1998. Biovariants of isolates of *Pasteurella* from domestic and wild ruminants. Journal of Veterinary Diagnostic Investigation 10: 49–55.
- LABARCA, C., AND K. PAIGEN. 1980. A simple, rapid, and sensitive DNA assay procedure. Analytical Biochemistry 102: 344–352.
- LICHTENSTEIGER, C. A., S. M. STEENBERGEN, R. M. LEE, D. D. POLSON, AND E. R. VIMR. 1996. Direct PCR analysis for toxigenic *Pasteurella multocida*. Journal of Clinical Microbiology 34: 3035–3039.
- MUTTERS, R., W. MANNHEIM, AND M. BISGAARD. 1989. Taxonomy of the group. In *Pasteurella* and pasteurellosis, C. Adlam and J. M. Rutter (eds.). Academic Press, New York, New York, pp. 3–34.
- NEI, M., AND W. -H. LI. 1979. Mathematical model for studying genetic variation in terms of restriction endonucleases. Proceedings of the National Academy of Science (USA) 76: 5269–5273.
- RIMLER, R. B. 2000. Restriction endonuclease analysis using *Hha* I and *Hpa* II to discriminate among group B *Pasteurella multocida* associated with haemorrhagic septicemia. Journal of Medical Microbiology 49: 81–87.
- , AND K. R. RHOADES. 1989. *Pasteurella multocida*. In *Pasteurella* and pasteurellosis, C. Adlam and J. M. Rutter (eds.). Academic Press, New York, New York, pp. 37–73.
- , AND M. A. WILSON. 1994. Re-examination of *Pasteurella multocida* serotypes that caused haemorrhagic septicemia in North America. The Veterinary Record 134: 10.
- ROSENBUSCH, C. T., AND I. A. MERCHANT. 1939. A study of the hemorrhagic septicemia *Pasteurella*. Journal of Bacteriology 37: 69–89.
- SNEATH, P. H. A., AND R. R. SOKAL. 1973. Numerical taxonomy. W. H. Freeman & Co., San Francisco, California, 573 pp.
- SNIPES, K. P., D. C. HIRSH, R. W. KASTEN, L. M. HANSEN, D. W. HIRD, T. W. CARPENTER, AND R. H. MC CAPES. 1989. Use of an rRNA probe and restriction endonuclease analysis to fingerprint *Pasteurella multocida* isolated from turkeys and wildlife. Journal of Clinical Microbiology 27: 1847–1853.
- THORNE, E. T. 1982. Miscellaneous bacteria. In Diseases of wildlife in Wyoming, 2nd Edition, E. T. Thorne, N. Kingston, W. R. Jolley and R. C. Bergstrom (eds.). Wyoming Game and Fish Department, Cheyenne, Wyoming, pp. 30–45.
- TIMONEY, J. F., J. H. GILLESPIE, F. W. SCOTT, AND J. E. BARLOUGH. 1988. The Genus *Pasteurella*. In Hagan and Bruner's microbiology and infectious diseases of domestic animals, 8th Edition, Comstock Publishing Associates, Cornell University Press, Ithaca, New York, pp. 104–116.
- TOWNSEND, K. M., J. D. BOYCE, J. Y. CHUNG, A. J. FROST, AND B. ADLER. 2001. Genetic organization of *Pasteurella multocida cap* loci and development of a multiplex capsular PCR typing system. Journal of Clinical Microbiology 39: 924–929.
- WARD, A. C. S. 1990. Isolation of *Pasteurellaceae* from bovine abortions. Journal of Veterinary Diagnostic Investigation 2: 59–62.
- , D. L. HUNTER, M. D. JAWORSKI, P. J. BEN-OLKIN, M. P. DOBEL, J. B. JEFFRESS, AND G. A. TANNER. 1997. *Pasteurella* spp. in sympatric bighorn and domestic sheep. Journal of Wildlife Diseases 33: 544–557.
- WILSON, M. A., R. M. DUNCAN, T. J. ROFFE, G. E. NORDHOLM, AND B. M. BERLOWSKI. 1995. Pas-

- teurellosis in elk (*Cervus elaphus*): DNA fingerprinting of isolates. *The Veterinary Record* 137: 195–196.
- ZHAO, G., C. PIJOAN, M. P. MURTAUGH, AND T. W. MOLITOR. 1992. Use of restriction endonuclease analysis and ribotyping to study epidemiology of *Pasteurella multocida* in closed swine herds. *Infection and Immunity* 60: 1401–1405.
- Received for publication 1 November 2002.*