

Influenza A Virus in Birds during Spring Migration in the Camargue, France

Authors: Lebarbenchon, Camille, Chang, Chung-Ming, van der Werf, Sylvie, Aubin, Jean-Thierry, Kayser, Yves, et al.

Source: Journal of Wildlife Diseases, 43(4) : 789-793

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-43.4.789>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Influenza A Virus in Birds during Spring Migration in the Camargue, France

Camille Lebarbenchon,^{1,2,4} Chung-Ming Chang,^{2,3} Sylvie van der Werf,³ Jean-Thierry Aubin,³ Yves Kayser,¹ Manuel Ballesteros,¹ François Renaud,² Frédéric Thomas,² and Michel Gauthier-Clerc¹

¹Station Biologique de la Tour du Valat, Le Sambuc, 13200 Arles, France; ²GEMI, UMR CNRS/IRD 2724, IRD, 911 Avenue, Agropolis BP 64501, 34394 Montpellier Cedex 5, France; ³Unité de Génétique Moléculaire des Virus Respiratoires, URA1966 CNRS, EA302 Université Paris 7, Institut Pasteur, 25-28 Rue du Dr Roux, 75724 Paris Cedex 15, France; ⁴Corresponding author (email: lebarbenchon@tourduvalat.org)

ABSTRACT: Wild aquatic birds are considered to be the natural reservoir for influenza A viruses, and previous studies have focused mainly on species in the orders Anseriformes and Charadriiformes. In this study, we surveyed a larger spectrum of potential hosts belonging to 10 avian orders. Cloacal swabs ($n=1,044$) from 72 free-living bird species, were analysed by reverse transcription-polymerase chain reaction for the presence of avian influenza virus. Only two Mediterranean Gulls (*Larus melanocephalus*) tested positive; one of these viruses was identified as an H9N2 subtype. The absence of infection among passerine birds supports the idea that the prevalence of avian influenza virus infection in terrestrial species is low.

Key words: Avian influenza, H9N2, *Larus melanocephalus*, Mediterranean Gulls, passerine birds.

Wild aquatic birds in the orders Anseriformes (ducks, geese, and swans) and Charadriiformes (gulls, terns, and waders) are traditionally considered as natural hosts for most avian influenza viruses (AIV; Webster et al., 1992). Although AIV have been isolated from other wild bird species (Olsen et al., 2006), less attention has been devoted to species in other avian orders. Consequently, the role of many avian species as potential hosts for AIV is unknown.

The Camargue area is an alluvial wetland covering some 140,000 ha in the Rhône delta in the south of France, and it is at the crossroads of numerous migratory routes of Palaearctic birds (Blondel and Isenmann, 1981; Berthold, 2001). Therefore, the Camargue is considered as a high-risk area for the introduction and transmission of diseases transmitted by wild avian species, such as West Nile virus

(Jourdain et al., 2007b) and AIV. The risk of virus introduction by birds from sub-Saharan Africa into Western Europe is highest during spring migration between March and June (Jourdain et al., 2007a). The aim of this study was to investigate the prevalence of AIV infection in a large diversity of avian hosts during spring migration.

From mid-March to late June 2006, different bird species were trapped using mist nets. Nets were placed in bushes located a few hundred meters behind the Piémanson beach, southeast of Salins de Giraud (Arles, France), in an attempt to capture migratory birds immediately after their crossing of the Mediterranean Sea. Cloacal swabs were used to collect fecal samples, except for passerines; due to their small size, we collected samples of fresh droppings to avoid injury. For Ciconiiformes (herons and egrets) and Phoenicopteriformes (greater flamingo [*Phoenicopterus ruber*]), cloacal swabs were collected from chicks in breeding colonies. For gulls, fresh dropping samples were collected near nests.

Cloacal swabs were collected using the Viral Pack kit (Biomedics, S.L., Madrid, Spain) and kept at 4 C until they were transported back to the laboratory and kept at –80 C until RNA extraction was performed. Automatic RNA extraction was performed using the BioRobot MDx workstation and QIAamp Virus BioRobot MDX kit (QIAGEN GmbH, Hilden, Germany) according to the manufacturer's instructions. The presence of AIV was first detected by reverse transcription-polymerase chain reaction (RT-PCR) targeting

TABLE 1. Bird orders, families, species, number of samples analyzed, and number of infected birds.

Order	Family	Species	Common name	<i>n</i>	NIB ^a
Accipitriformes	Accipitridae	<i>Accipiter nisus</i>	Eurasian Sparrowhawk	1	
		<i>Buteo buteo</i>	Common Buzzard	2	
Caprimulgiformes	Caprimulgidae	<i>Caprimulgus europaeus</i>	European Nightjar	2	2
Charadriiformes	Laridae	<i>Larus melanocephalus</i>	Mediterranean Gull	67	
		<i>Larus michahellis</i>	Yellow-legged Gull	29	
	Recurvirostridae	<i>Recurvirostra avosetta</i>	Pied Avocet	4	
	Scolopacidae	<i>Actitis hypoleucos</i>	Common Sandpiper	1	
	Sternidae	<i>Sterna albifrons</i>	Little Tern	1	
	Ardeidae	<i>Ardea cinerea</i>	Grey Heron	1	
		<i>Ardeola ralloides</i>	Squacco Heron	33	
		<i>Bubulcus ibis</i>	Cattle Egret	46	
		<i>Egretta garzetta</i>	Little Egret	31	
		<i>Ixobrychus minutus</i>	Little Bittern	3	
		<i>Nycticorax nycticorax</i>	Night Heron	12	
		<i>Ciconia ciconia</i>	White Stork	19	
	Threskiornithidae	<i>Plegadis falcinellus</i>	Glossy Ibis	30	
Columbiformes	Columbidae	<i>Streptopelia orientalis</i>	Oriental Turtle Dove	1	
		<i>Streptopelia turtur</i>	European Turtle Dove	2	
Coraciiformes	Meropidae	<i>Merops apiaster</i>	European Bee-eater	6	
	Upupidae	<i>Upupa epops</i>	Hoopoe	13	
Passeriformes	Corvidae	<i>Corvus monedula</i>	Jackdaw	8	
	Emberizidae	<i>Emberiza cirius</i>	Cirl Bunting	2	
	Fringillidae	<i>Carduelis cannabina</i>	Linnet	5	
<i>Carduelis spinus</i>		Siskin	2		
<i>Coccothraustes cocco-thraustes</i>		Hawfinch	1		
	<i>Fringilla coelebs</i>	Chaffinch	4		
	<i>Serinus serinus</i>	European Serin	1		
	Hirundinidae	<i>Delichon urbica</i>	House Martin	10	
	Laniidae	<i>Lanius senator</i>	Woodchat Shrike	3	
Motacillidae	<i>Anthus campestris</i>	Tawny Pipit	2		
	<i>Anthus pratensis</i>	Meadow Pipit	1		
	<i>Motacilla flava</i>	Yellow Wagtail	2		
Muscicapidae	<i>Ficedula albicollis</i>	Collared Flycatcher	1		
	<i>Ficedula hypoleuca</i>	Pied Flycatcher	24		
	<i>Muscicapa striata</i>	Spotted Flycatcher	9		
Oriolidae	<i>Oriolus oriolus</i>	Eurasian Golden oriole	1		
Paridae	<i>Parus major</i>	Great Tit	2		
Passeridae	<i>Passer domesticus</i>	House Sparrow	17		
	<i>Passer montanus</i>	Tree Sparrow	4		
Prunellidae	<i>Prunella modularis</i>	Dunnock	2		
Sturnidae	<i>Sturnus vulgaris</i>	Common Starling	1		
Sylviidae	<i>Acrocephalus scirpaceus</i>	Eurasian Reed-warbler	2		
	<i>Cettia cetti</i>	Cetti's Warbler	1		
	<i>Hippolais icterina</i>	Icterine Warbler	4		
	<i>Hippolais polyglotta</i>	Melodious Warbler	3		
	<i>Locustella naevia</i>	Grasshopper Warbler	1		
	<i>Phylloscopus bonelli</i>	Bonelli's Warbler	3		
	<i>Phylloscopus collybita</i>	Chiffchaff	29		
	<i>Phylloscopus sibilatrix</i>	Wood Warbler	2		
	<i>Phylloscopus trochilus</i>	Willow Warbler	93		
	<i>Phylloscopus trochilus acredula</i>		5		
	<i>Regulus regulus</i>	Goldcrest	3		
	<i>Sylvia atricapilla</i>	Blackcap	31		

TABLE 1. Continued.

Order	Family	Species	Common name	n	NIB ^a
		<i>Sylvia borin</i>	Garden Warbler	18	
		<i>Sylvia cantillans</i>	Subalpine Warbler	26	
		<i>Sylvia communis</i>	Whitethroat	13	
		<i>Sylvia conspicillata</i>	Spectacled Warbler	3	
		<i>Sylvia melanocephala</i>	Sardinian Warbler	5	
		<i>Sylvia undata</i>	Dartford Warbler	1	
	Troglodytidae	<i>Troglodytes troglodytes</i>	Wren	3	
	Turdidae	<i>Erithacus rubecula</i>	Eurasian Robin	145	
		<i>Luscinia megarhynchos</i>	Common Nightingale	14	
		<i>Oenanthe hispanica</i>	Black-eared Wheatear	2	
		<i>Oenanthe oenanthe</i>	Northern Wheatear	1	
		<i>Phoenicurus ochruros</i>	Black Redstart	9	
		<i>Phoenicurus phoenicurus</i>	Common Redstart	65	
		<i>Saxicola rubetra</i>	Whinchat	1	
		<i>Turdus merula</i>	Blackbird	7	
		<i>Turdus philomelos</i>	Song Thrush	29	
Phoenicopteriformes	Phoenicopteridae	<i>Phoenicopterus ruber</i>	Greater Flamingo	113	
Piciformes	Picidae	<i>Jynx torquilla</i>	Wryneck	3	
Strigiformes	Strigidae	<i>Asio otus</i>	Long-eared Owl	1	
		<i>Otus scops</i>	Eurasian Scops Owl	2	

^a NIB = number of infected birds.

the conserved matrix gene segment. Superscript II kit (Invitrogen, Carlsbad, California, USA) was used for RT-PCR in a final volume of 25 µl containing 5 µl total RNA in the presence of 0.4 µM final concentration of each primer: M52C, 5'-CTT CTA ACC GAG GTC GAA ACG-3' and M253R, 5'-AGG GCA TTT TGG ACA AAK CGT CTA-3' (Fouchier et al., 2000). Cycling conditions included a reverse transcription step for 30 min at 45 C, 15 min at 55 C, and 2 min at 94 C; PCR reaction was performed during five cycles, including 15 sec at 94 C, 30 sec at 45 C, and 30 sec at 72 C and during 35 cycles for 15 sec at 94 C, 30 sec at 55 C, and 30 sec (plus 2 sec per cycle) at 72 C. The RT-PCR amplification products were analyzed by 1% agarose gel electrophoresis with ethidium bromide staining.

To confirm RT-PCR results, positive and weakly positive samples were tested again by real-time quantitative RT-PCR (RT-qPCR) using the same primers as for RT-PCR with a specific hydrolysis probe, 5'-(Fam)GCT AAA GAC AAG ACC AAT

CCT GTC ACC TCT G(Tamra)-3' (Sigma-Proligo, Saint-Quentin, Fallavier, France), for the AIV matrix gene. LightCycler RT-qPCR was carried out using the LightCycler RNA amplification kit for probe hybridization (Roche Biochemicals, Basel, Switzerland). Amplification and detection were performed on a LightCycler 1.5 (Roche Biochemicals) after one cycle of reverse transcription (55 C, 15 min; 95 C, 2 min) and 45 cycles of amplification (95 C, 10 sec; 58 C, 30 sec). The RT-qPCR results were analyzed by LightCycler 3.0 software (Roche Biochemicals). Positive samples were subtyped at the Institut Pasteur (Paris, France), using a SYBR Green-based RT-qPCR technique. For the H5, an RT-qPCR was performed on a LightCycler 1.5 using primers H5+/1544, 5'-CCG CAG TAT TCA GAA GAA GC-3' and H5-/1683, 5'-AGA CCA GCT ACC ATG ATT GC-3' and the specific probe H5/probe/1638-1662, 5'-(Fam) AGT GCT AGG GAA CTC GCC ACT GTA G (Tamra)-3' following reaction conditions mentioned above.

Fecal samples were collected and tested

from 1,044 free-living birds representing 72 species, 30 families, and 10 orders (Table 1). Passerine birds ($n=621$) represent 59.5% of our total sampling. Of the 25 samples found positive by the initial RT-PCR, only two fecal samples from Mediterranean Gulls (*Larus melanocephalus*) were confirmed positive by RT-qPCR. Prevalence for this species reached 3% ($n=67$); 2% when all Charadriiformes' samples ($n=102$) or gull samples ($n=96$) were considered.

The prevalence of AIV detected in gulls is comparable with recent data found in the literature for Eurasian species (Fouchier et al., 2003) and to prevalence estimates from numerous studies reporting AIV isolations from gulls and terns from 1974 to 1984 from Eurasia, North America, and Australia (Stallknecht and Shane, 1988). Based on a summary from all published reports of AIV isolations from gulls worldwide, a prevalence of 1.4% was reported (Olsen et al., 2006). In gulls, AIVs are divided into American and Eurasian lineages, and some specific subtypes, such as the H13 and H16, are thought to be associated with a gull reservoir (Fouchier et al., 2005; Olsen et al., 2006).

Molecular subtyping of hemagglutinin and neuraminidase revealed that the virus from one of the two positive gull samples is H9N2, whereas the H5N1 and H5N2 subtypes were formally excluded for the other sample. The H9 AIV has been isolated in wild ducks throughout the world. However, for gulls, the only available report of H9 viruses is from the Americas (Obenauer et al., 2006), but these were not of the N2 subtype. Influenza A H9N2 viruses have been detected worldwide in poultry, and they currently are endemic in poultry in Asia (Li et al., 2005; Xu et al., 2007). Cases of H9N2 influenza virus infection in humans in this area have also been reported since 1999 (Peiris et al., 1999). Our results support a circulation of H9N2 AIV in Mediterranean Charadriiformes, although more investigations are

required to better understand the AIV infection level for Palearctic species.

Gulls breed in colonies located on small islands in salt marsh areas. Although it is possible that these habitats may be less favorable for environmental transmission via water than freshwater habitats (Stallknecht et al., 1990), this may be offset by a high contact rate. Like gulls, Ciconiiformes, and Phoenicopteriformes breed in colonies, with adults and juveniles crowded in small areas, this creates good opportunities for viral transmission. The absence of positive detection of AIV is of particular value, because very few data are available for these orders.

Prevalence of AIV infections in passerine birds is known to be particularly low (Fouchier et al., 2003; Morishita et al., 1999; Schnebel et al., 2005). Although our results support these observations, further data should be collected on a larger number of species and individuals, and at other periods of the year to more fully understand the ecology of AIV in Passeriformes.

We are grateful to Antoine Arnaud, Leire Paz, Philippe Perret, Christophe Pin, and Nicolas Vincent-Martin for help on birds sampling. We acknowledge the technical help of Frédérique Cuvelier, Patricia Jeannin, Vanessa Roca, and Philippe Thebault for analysis of the samples. Camille Lebarbenchon is supported by a "Tour du Valat/Région Languedoc-Roussillon" PhD grant. This work was funded by the French "Agence Nationale de la Recherche" (ANR) "Santé-Environnement."

LITERATURE CITED

- BERTHOLD, P. 2001. Bird migration: A general survey. Oxford University Press, Oxford, UK, 253 pp.
- BLONDEL, J., AND P. ISENMANN. 1981. Guide des oiseaux de Camargue. Delachaux & Niestlé, Neuchâtel, Switzerland.
- FOUCHIER, R. A. M., T. M. BESTEBOER, S. HERFST, L. VAN DER KEMP, G. S. RIMMELZWANN, AND A. D. M. E. OSTERHAUS. 2000. Detection of influenza A viruses from different species by PCR amplification of conserved sequences in the matrix

- gene. *Journal of Clinical Microbiology* 38: 4096–4101.
- , B. OLSEN, T. M. BESTEBROER, S. HERFST, L. VAN DER KEMP, G. F. RIMMELZWAAN, AND A. D. M. E. OSTERHAUS. 2003. Influenza A virus surveillance in wild birds in Northern Europe in 1999 and 2000. *Avian Diseases* 47: 857–860.
- , V. MUNSTER, A. WALLENSTEN, T. M. BESTEBROER, S. HERFST, D. SMITH, G. F. RIMMELZWAAN, B. OLSEN, AND A. D. M. E. OSTERHAUS. 2005. Characterization of a novel influenza A virus hemagglutinin subtype (H16) obtained from black-headed gulls. *Journal of Virology* 79: 2814–2822.
- JOURDAIN, E., M. GAUTHIER-CLERC, D. BICOUT, AND P. SABATIER. 2007a. Bird migration routes and risk for pathogen dispersion into western Mediterranean wetlands. *Emerging Infectious Diseases* 13: 365–372.
- , Y. TOUSSAINT, A. LEBLOND, D. BICOUT, P. SABATIER, AND M. GAUTHIER-CLERC. 2007b. Bird species potentially involved in introduction, amplification and spread of West Nile Virus in a Mediterranean wetland, the Camargue (southern France). *Vector-Borne and Zoonotic Diseases* 7: 15–33.
- MORISHITA, T. Y., P. P. AYE, E. C. LEY, AND B. S. HARR. 1999. Survey of pathogens and blood parasites in free-living passerines. *Avian Diseases* 43: 549–552.
- LI, C., K. YU, G. TIAN, D. YU, L. LIU, B. JING, J. PING, AND H. CHEN. 2005. Evolution of H9N2 influenza viruses from domestic poultry in mainland China. *Virology* 340: 70–83.
- OBENAUER, J. C., J. DENSON, P. K. MEHTA, X. SU, S. MUKATIRA, D. B. FINKELSTEIN, X. XU, J. WANG, J. MA, F. YIPING, K. M. RAKESTRAW, R. G. WEBSTER, E. HOFFMANN, S. KRAUSS, J. ZHENG, Z. ZHANG, AND C. W. NAEVE. 2006. Large-scale analysis of avian influenza isolates. *Science* 311: 1576–1580.
- OLSEN, B., V. J. MUNSTER, A. WALLENSTEN, J. WALDENSTRÖM, A. D. M. E. OSTERHAUS, AND R. A. M. FOUCHIER. 2006. Global patterns of influenza A virus in wild birds. *Science* 312: 384–388.
- PEIRIS, M., K. Y. YUEN, C. W. LEUNG, K. H. CHAN, P. L. IP, R. W. LAI, W. K. ORR, AND K. F. SHORTRIDGE. 1999. Human infection with influenza H9N2. *Lancet* 354: 916–917.
- SCHNEBEL, B., DIERSCHKE, V., RAUTENSCHLEIN, S., AND M. RYLL. 2005. No detection of avian influenza A viruses of the subtypes H5 and H7 and isolation of lentogenic avian paramyxovirus serotype 1 in passerine birds during stopover in the year 2001 on the island Helgoland (North Sea). *Deutsche Tierärztliche Wochenschrift* 112: 456–460.
- STALLKNECHT, D. E., AND S. M. SHANE. 1988. Host range of avian influenza virus in free-living birds. *Veterinary Research Communications* 12: 125–141.
- , M. T. KEARNEY, S. M. SHANE, AND P. J. ZWANK. 1990. Effects of pH, temperature, and salinity on persistence of avian influenza viruses in water. *Avian Diseases* 34: 412–428.
- WEBSTER, R. G., W. J. BEAN, O. T. GORMAN, T. M. CHAMBERS, AND Y. KAWAOKA. 1992. Evolution and ecology of influenza A viruses. *Microbiological Reviews* 56: 152–179.
- XU, K. M., K. S. LI, G. J. D. SMITH, J. W. LI, H. TAI, J. X. ZHANG, R. G. WEBSTER, J. S. M. PEIRIS, H. CHEN, AND Y. GUAN. 2007. Evolution and molecular epidemiology of H9N2 influenza viruses from quail in southern China, 2000 to 2005. *Journal of Virology* 81: 2635–2645.

Received for publication 7 November 2006.