

HEMATOLOGICAL VALUES OF CAPTIVE ROCKY MOUNTAIN BIGHORNS

Authors: WOOLF, ALAN, and KRADEL, DAVID C.

Source: Journal of Wildlife Diseases, 6(1) : 67-68

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-6.1.67>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

HEMATOLOGICAL VALUES OF CAPTIVE ROCKY MOUNTAIN BIGHORNS

Reports of blood values for Rocky Mountain Bighorn Sheep (*Ovis canadensis canadensis* Shaw) have not been published to our knowledge and we needed to know normal values for our studies of the physiology and diseases of bighorns in captivity. In April 1969, blood samples were collected from 10 captive bighorns in apparent good health. The only known health problem affecting the animals at the time of sampling were light to moderate endoparasitic infestations in all sheep determined by Baerman extraction and sugar-flotation analysis of fecal samples. However, the parasitism was not severe enough to produce signs of illness or discomfort.

Erythrocytes and leucocytes were counted with a hemacytometer manufactured by Celloscope Particle Data Inc. Hematocrit was determined by the microhematocrit method, spinning for five minutes at 14,300 RCF. Hemoglobin was measured by the cyanomethemoglo-

bin method. The differential WBC was determined by standard methods.

Total serum protein, albumin, and globulin were measured by the Biuret Ferro-Ham modification reported in the Dade Manual (Anom. 1965. *Dade Manual of Clinical Chemistry Procedures*. Dade Reagents, Inc. Miami, Florida, 194 p.). Serum phosphorous was determined by the Fiske and Subba Row-Gomori modification (op. cit.); magnesium by the Orange-Rhein-modified (Titan Yellow) method (op. cit.); and calcium by the Murexide method — modified (Virginia Marshall, Iowa Veterinary Diagnostic Lab, pers. comm.).

Since these are preliminary data, the results are presented without interpretation. The hematological values are shown in Tables 1, 2, and 3. Additional sampling will continue to represent all seasons, to obtain a range of values for each animal, and to compare "normal" values with those determined when animals are displaying visual signs of disease.

TABLE 1. *Erythrocyte and leukocyte values for captive bighorn sheep.*

| Sex/Age | Test Number | WBC* | RBC ($\times 10^6$) | PCV (%) | Hemoglobin (g/100ml) |
|---------|-------------|-------|-----------------------|---------|----------------------|
| M—2 yrs | # 5 | 6,494 | 9.17 | 48 | 16.4 |
| M—6 yrs | # 2 | 7,814 | 10.42 | 55 | 23+ |
| F—1 yr | # 2 | 6,821 | 10.23 | 53 | 18.5 |
| F—2 yrs | # 8 | 5,677 | 10.43 | 56 | 19.0 |
| F—2 yrs | #11 | 6,009 | 9.70 | 50 | 17.4 |
| F—3 yrs | # 3 | 6,716 | 10.07 | 54 | 19.0 |
| F—4 yrs | # 1 | 5,723 | 10.51 | 53 | 18.0 |
| F—5 yrs | # 4 | 7,719 | 9.96 | 54 | 18.0 |
| F—5 yrs | # 9 | 6,245 | 10.28 | 56 | 19.0 |
| F—8 yrs | #10 | 7,458 | 10.05 | 52 | 18.0 |

* Abbreviations stand for leukocytes (WBC), erythrocytes (RBC), and hematocrit (PCV).

TABLE 2. *Differential white blood cell counts for captive bighorns.*

| Sex/Age | Test Number | Polymorphs | Percent of Count | | |
|---------|-------------|------------|------------------|-----------|--------------|
| | | | Lymphocytes | Monocytes | Eosinophiles |
| M—2 yrs | # 5 | 61 | 32 | 3 | 4 |
| M—6 yrs | # 2 | 53 | 35 | 4 | 8 |
| F—1 yr | # 6 | 38 | 58 | 1 | 3 |
| F—2 yrs | # 8 | 39 | 53 | 2 | 6 |
| F—2 yrs | #11 | 42 | 56 | 1 | 1 |
| F—3 yrs | # 3 | 55 | 42 | 1 | 2 |
| F—4 yrs | # 1 | 56 | 39 | 2 | 3 |
| F—5 yrs | # 4 | 55 | 40 | 2 | 3 |
| F—5 yrs | # 9 | 36 | 53 | 3 | 7 |
| F—8 yrs | #10 | 41 | 39 | 3 | 7 |

TABLE 3. *Blood serum proteins and minerals for captive bighorns.*

| Sex/Age | Test Number | Total protein | | | | Ca | P (mg/100 ml) | Mg |
|---------|-------------|---------------|----------|-----|--------------------|------|------------------|-----|
| | | Albumin | Globulin | A/G | (grams per 100 ml) | | | |
| M—2 yrs | # 5 | 6.8 | 4.0 | 2.8 | 1.4 | 11.3 | 10.8 | 3.5 |
| M—6 yrs | # 2 | 7.9 | 4.0 | 3.9 | 1 | 11.3 | 9.2 | 3.0 |
| F—1 yr | # 6 | 8.2 | 4.5 | 3.7 | 1.2 | 12.4 | 7.6 | 3.5 |
| F—2 yrs | # 8 | 7.9 | 5.5 | 2.4 | 2.2 | 13.2 | 8.2 | — |
| F—2 yrs | #11 | 7.0 | 4.4 | 2.6 | 1.7 | 9.6 | 7.3 | — |
| F—3 yrs | # 3 | 7.5 | 4.5 | 3.0 | 1.5 | 10.9 | 5.8 | 3.0 |
| F—4 yrs | # 1 | 7.2 | 4.5 | 2.7 | 1.6 | 12.4 | 6.0 | — |
| F—5 yrs | # 4 | 7.7 | 4.4 | 3.3 | 1.3 | 11.0 | 5.6 | — |
| F—5 yrs | # 9 | 8.2 | 4.5 | 3.7 | 1.2 | 12.0 | 4.7 | 3.0 |
| F—8 yrs | #10 | 7.7 | 4.5 | 3.2 | 1.4 | 11.7 | 4.5 | 2.7 |

ALAN WOOLF

*Rachelwood Wildlife Research Preserve
New Florence, Pennsylvania*

DAVID C. KRADEL

*Animal Disease Laboratory
Pennsylvania State University
University Park, Pennsylvania*

July 31, 1969