



THE OCCURRENCE OF DRACUNULUS INSIGNIS (LEIDY, 1858) CHANDLER, 1942 IN A SKUNK FROM ONTARIO, CANADA

Authors: WEBSTER, W. A., and CASEY, G. A.

Source: Journal of Wildlife Diseases, 6(1) : 71

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-6.1.71>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

THE OCCURRENCE OF *DRACUNULUS INSIGNIS* (LEIDY, 1858) CHANDLER, 1942 IN A SKUNK FROM ONTARIO, CANADA

There have been numerous reports of guinea worm infection in wild carnivores in North America (see Ewing and Hibbs, 1966, Amer. Midl. Nat. 76: 515-519 for a review of reported infections). However, only one such record lists *Dracunculus* sp. from a Canadian host; Chitwood (1933, J. Amer. Med. Assoc. 100: 802-804) reporting the occurrence in a raccoon from Ontario.

Over a period of six years, we have examined numerous wild carnivores: i.e. raccoons, foxes, mink, and various weasels. In particular, approximately 125 striped skunks, *Mephitis mephitis* have been skinned in connection with various projects. It is only recently that we have found a single gravid female nematode indistinguishable from those described by Chandler (1942, J. Parasitol. 28: 255-268; Amer. J. Trop. Med. 22: 153-157)

as *Dracunculus insignis* (Leidy, 1858). Our specimen was found loosely coiled in the subcutaneous tissue of the front leg of a skunk from Peterborough, Ontario. From the number of skunks examined, both from Ontario and Quebec, and the single finding of this nematode, it would appear that *D. insignis* is not common in skunks in this part of North America. This is in agreement with other observations since *D. insignis* has been reported from North American skunks only three times previously; in *Mephitis nigra* from New York (Cheatum and Cook, 1948, Cornell Vet. 38: 421-423); in *Mephitis mephitis* from Minnesota (Erickson, 1946, Amer. Midl. Nat. 36: 494-509) and Nebraska (Ewing and Hibbs, 1966, Amer. Midl. Nat. 76: 515-519).

W. A. WEBSTER

G. A. CASEY

*Animal Pathology Division,
Health of Animals Branch,
Canada Department of Agriculture,
Animal Diseases Research Institute,
P.O. Box 1400, Hull, Quebec.*

August 6, 1969