



Echimyopus dasypus Fain et al. (Acari: Astigmatina: Echimyopodidae) from a nine-banded armadillo, Dasypus novemcinctus L. (Mammalia: Dasypodidae), in Florida, USA

Authors: Mertins, James W., Hanson, Britta A., and Corn, Joseph L.

Source: Systematic and Applied Acarology, 16(3) : 252-254

Published By: Systematic and Applied Acarology Society

URL: <https://doi.org/10.11158/saa.16.3.10>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

***Echimyopus dasypus* Fain *et al.* (Acari: Astigmatina: Echimyopodidae) from a nine-banded armadillo, *Dasypus novemcinctus* L. (Mammalia: Dasypodidae), in Florida, USA**

JAMES W. MERTINS¹, BRITTA A. HANSON² & JOSEPH L. CORN²

¹U.S. Department of Agriculture, Animal and Plant Health Inspection Service, Veterinary Services, National Veterinary Services Laboratories, 1920 Dayton Avenue, Ames, Iowa 50010, USA. E-mail: James.W.Mertins@aphis.usda.gov

²Southeastern Cooperative Wildlife Disease Study, College of Veterinary Medicine, University of Georgia, Athens, Georgia 30602, USA.

Abstract

We collected parasitic deutonymphs of *Echimyopus dasypus* Fain *et al.* from a wild-caught nine-banded armadillo, *Dasypus novemcinctus* L., for the first time within the United States of America, although this mite was previously found in England on armadillos imported from Florida. The new specimens came from a free-ranging animal captured in Hendry County, Florida, in 2004.

Key words: Deutonymph, hypopus, mange, *Blomia*, new distribution data, *Amblyomma auricularium*

The astigmatine mite family Echimyopodidae (five genera, 31 species) is associated mostly with rodents and marsupials in Australia, South America, and North America (OConnor 2009). Many echimyopodid species are known from only their deutonymphal stages (hypopi), which live as subcutaneous or endofollicular parasites of their mammalian hosts, but presumed, the free-living stages habitually live as detritivores in mammal nests (OConnor 1982). Free-living life stages are described for only a few species in three of the genera, i.e., *Marmosopus* Fain & Lukoschus, *Echimyopus* Fain, and *Blomia* Oudemans, the latter of which have no known deutonymphs and no demonstrated parasitic connections (Mariana *et al.* 1996). Alone in the family, the seven currently recognized *Blomia* species are widely known primarily as synanthropic house dust and stored food mites (Colloff 2009).

At least two species of *Blomia* are documented from the USA (Banks 1917, Fain *et al.* 1977, Fernandez-Caldas *et al.* 1990), but to date, members of only one of the parasitic echimyopodid genera, *Echimyopus*, have been found there. Whitaker *et al.* (1976) first found deutonymphal *E. nyctomys* Fain (later reidentified as *E. orphanus* Fain & Philips by Whitaker *et al.* (2007)) on eastern gray, *Sciurus carolinensis* Gmelin, and fox squirrels, *S. niger* L., in Indiana. Fain and Philips (1977) described deutonymphs of *E. orphanus* from the nest of an eastern screech owl, *Megascops asio* (L.), in New York (Philips & Dindal 1990), but at the time, they surmised that the mites came from an unknown rodent host. Later, Fain and Philips (1981) used additional mites from the same owl nest to describe the protonymph, tritonymph, and adult female of *E. orphanus* and clarify the status of this taxon as a senior synonym of *Blomia angustivulva* Fain & Philips. A second purported US *Echimyopus* species record came only by indirect inference involving the collection of *E. dasypus* deutonymphs in England from eight of 20 newly imported nine-banded armadillos, *D. novemcinctus*, that originated in Florida (locality unspecified) (Baskerville & Francis 1981). Also, Forrester (1992) cited unpublished observations of two many armadillos (of 130 examined) in Alachua County, Florida, with no demonstrable mites. In the present report, we document *E. dasypus* from a wild-caught armadillo at a specified locality in Florida.

As part of a long-term, extensive surveillance effort looking for exotic parasites on wildlife species in the southeastern US and Caribbean region (Hansen *et al.* 2007, Mertins *et al.* 2009, Corn *et al.* 2009, 2011), we periodically encounter and examine nine-banded armadillos. On 4 May 2004, we collected ectoparasites (NVSL Accession Number 317368) from an adult male armadillo in the Okaloacoochee State Forest, Hendry County, Florida, USA. The parasites were preserved in a labeled vial of 70% isopropyl alcohol by field personnel of the Southeastern Cooperative Wildlife Disease Study (SCWDS), and later, they were sorted and identified at the US Department of Agriculture, Animal and Plant Health Inspection Service, National Veterinary Services Laboratories in Ames, Iowa, USA. The most evident, larger parasites on the host animal were 23 adult armadillo ticks, *Amblyomma auricularium* (Conil), but the collection vial also contained 24 relatively minute (~200 μm) mites identified as deutonymphs of *E. dasyopus* by comparison with the original description (Fain *et al.* 1973). Neither field notes nor personal recollections provide any pertinent information on the health and physical condition of the infested host armadillo. In both of the previous known occurrences of *E. dasyopus* (Fain *et al.* 1973, Baskerville & Francis 1981), infested armadillos showed dry, crumbly mange over extensive ventral areas of their necks, thoraces, and abdomens, and each individual mite was attached to the host epidermis within the lacuna of a separate, gall-like cyst in these areas.

Armadillos may not be native residents of the USA, although by the early 1850s, they were present in the southern tip of Texas, probably having crossed the Rio Grande from Mexico (Lever 1985). Through their ongoing dispersal movements, their US range has since expanded northwards and eastwards from South Texas to encompass most of the Southeast south of 38 degrees north latitude (Taulman & Robbins 1996, Hofmann 2005). However, the first armadillos in peninsular Florida were deliberately introduced, beginning during World War I (Bailey 1924, Lever 1985), although they evidently did not establish viable feral populations until about 1924 in Brevard County, after releases from a private zoo in Titusville (Taulman & Robbins 1996). By direct implication, *E. dasyopus*—a species-specific associate of nine-banded armadillos—is likewise not a native resident of Florida or the USA, entering on either naturally dispersing or actively introduced host animals.

The type specimens for *E. dasyopus* came from two armadillos at two different localities in the Republic of Panama (Fain *et al.* 1973), but until our collection in Hendry County, Florida, those were the only documented, detailed distribution records for this mite. The exact origins in Florida of the armadillos and their associated *E. dasyopus* found by Baskerville and Francis (1981) in England are unknown. We do, however, suspect that this mite may be a widespread, though possibly uncommon, associate of *D. novemcinctus* throughout Florida and in the rest of its New World geographic range.

Acknowledgements. We gratefully acknowledge Craig Okraska for assistance in the field and the Okaloacoochee State Forest for permitting the collections. Funding for this project was provided through Cooperative Agreements 0491130808CA and 1091130808CA, Veterinary Services, Animal and Plant Health Inspection Service, U.S. Department of Agriculture. Additional funds were provided through sponsorship from the fish and wildlife agencies of Alabama, Arkansas, Florida, Georgia, Kansas, Kentucky, Louisiana, Maryland, Mississippi, Missouri, North Carolina, Puerto Rico, South Carolina, Tennessee, Virginia, and West Virginia; through the Federal Aid to Wildlife Restoration Act (50 Stat. 917) and Grant Agreement 06ERAG0005, Biological Resources Division, U.S. Geological Survey, U.S. Department of the Interior; and through Cooperative Agreements 0496130032CA and 1096130032CA, Veterinary Services, Animal and Plant Health Inspection Service, U.S. Department of Agriculture.

References

- Bailey, H.H. (1924) The armadillo in Florida and how it reached there. *Journal of Mammalogy*, 5, 264–265.
- Banks, N. (1917) New mites, mostly economic (Arachn., Acar.). *Entomological News*, 28, 193–199.
- Baskerville, A. & Francis, L. (1981) Mange in newly-imported armadillos (*Dasyurus novemcinctus*). *Laboratory Animals*, 15, 305–307.
- Colloff, M.J. (2009) *Dust Mites*. Dordrecht, Springer Science and Collingwood, CSIRO Publishing. 583 pp.
- Corn, J.L., Berger, P. & Mertins, J.W. (2009) Surveys for ectoparasites on wildlife associated with *Amblyomma variegatum* (Acari: Ixodidae)-infested livestock in St. Croix, United States Virgin Islands. *Journal of Medical Entomology*, 46, 1483–1489.
- Corn, J.L., Mertins, J.W., Hanson, B. & Snow, S. (2011) First reports of ectoparasites collected from wild-caught exotic reptiles in Florida. *Journal of Medical Entomology*, 48, 94–100.
- Fain, A. & Philips, J.R. (1977) Astigmatic mites from nests of birds of prey in U.S.A. I. Description of four new species of Glycyphagidae. *International Journal of Acarology*, 3, 105–114.
- Fain, A. & Philips, J.R. (1981) Astigmatic mites from nests of birds of prey in U.S.A. VI. The adult forms of *Echimyopus orphanus* Fain & Philips, 1977 and of *Dermacarus pilarsus* Fain & Philips, 1977. *International Journal of Acarology*, 7, 235–237.
- Fain, A., Hyland, K. & Tadmowski, T. (1977) *Blomia gracilipes* (= *Chortoglyphus gracilipes*): redescription & status (Acarina: Glycyphagidae). *Entomological News*, 88, 267–269.
- Fain, A., Lukoschus, F.S., Louppen, J.M.W. & Méndez, E. (1973) *Echimyopus dasyopus* n. sp., a hypopus from *Dasyurus novemcinctus* in Panama (Glycyphagidae, Echimyopinae, Sarcoptiformes). *Journal of Medical Entomology*, 10, 552–555.
- Fernandez-Caldas, E., Fox, R.W., Bucholz, G.A., Trudeau, W.L., Ledford, D.K. & Lockey, R.F. (1990) House dust mite allergy in Florida. Mite survey in households of mite-sensitive individuals in Tampa, Florida. *Allergy Proceedings*, 11, 263–267.
- Forrester, D.J. (1992) *Parasites and Diseases of Wild Mammals in Florida*. Gainesville, Florida, University of Florida Press. 459 pp.
- Hanson, B.A., Frank, P.A., Mertins, J.W. & Corn, J.L. (2007) Tick paralysis of a snake caused by *Amblyomma rotundatum* (Acari: Ixodidae). *Journal of Medical Entomology*, 44, 155–157.
- Hofmann, J.E. (2005) A survey for the nine-banded armadillo (*Dasyurus novemcinctus*) in Illinois. Illinois Department of Natural Resources, Center for Biodiversity, *Technical Report 2005*, 16, 1–29.
- Lever, C. (1985) *Naturalized Mammals of the World*. London, Longman Group Ltd. 487 pp.
- Mariana, A., Ho, T.M. & Heah, S.K. (1996) Life-cycle, longevity and fecundity of *Blomia tropicalis* (Acari: Glycyphagidae) in a tropical laboratory. *Southeast Asian Journal of Tropical Medicine and Public Health*, 27, 392–395.
- Mertins, J.W., Hanson, B.A., & Corn, J.L. (2009) *Whartonacarus floridensis* sp.n. (Acarina: Trombiculidae), with a generic review and the first record of *Whartonacarus* chiggers from the continental United States. *Journal of Medical Entomology*, 46, 1260–1268.
- OConnor, B.M. (1982) Evolutionary ecology of astigmatid mites. *Annual Review of Entomology*, 27, 385–409.
- OConnor, B.M. (2009) Cohort Astigmatina. In: Krantz, G. W. & Walter, D. E. (eds.) *A Manual of Acarology*, 3rd edition. Lubbock, Texas Tech University Press. pp. 565–657.
- Philips, J.R. & Dindal, D.L. (1990) Invertebrate populations in the nests of a screech owl (*Otus asio*) and an American kestrel (*Falco sparverius*) in central New York. *Entomological News*, 101, 170–192.
- Taulman, J.F. & Robbins, L.W. (1996) Recent range expansion and distributional limits of the nine-banded armadillo (*Dasyurus novemcinctus*) in the United States. *Journal of Biogeography*, 23, 635–648.
- Whitaker, J.O. Jr., Spicka, E.J. & Schmeltz, L.L. (1976) Ectoparasites of squirrels of the genus *Sciurus* from Indiana. *Proceedings of the Indiana Academy of Science*, 85, 431–436.
- Whitaker, J.O. Jr., Walters, B.L., Castor, L.K., Ritzi, C.M. & Wilson, N. (2007) Host and distribution lists of mites (Acari), parasitic and phoretic, in the hair or on the skin of North America wild mammals north of Mexico: records since 1974. *Faculty Publications from the Harold W. Manter Laboratory of Parasitology* 1. Lincoln, Nebraska, University of Nebraska. 173 pp. Available from: <http://digitalcommons.unl.edu/parasitologyfacpubs/1/> (26 April, 2011).

Accepted by Anne Baker: 6 May 2011; published 14 Oct. 2011