

Chromium, Cadmium, Lead, and Arsenic Concentrations in Water, Vegetables, and Seafood Consumed in a Coastal Area in Northern Vietnam

Authors: Ngoc, Nguyen Thi Minh, Chuyen, Nguyen Van, Thao, Nguyen Thi Thu, Duc, Nguyen Quang, Trang, Nguyen Thi Thu, et al.

Source: Environmental Health Insights, 14(1)

Published By: SAGE Publishing

URL: <https://doi.org/10.1177/1178630220921410>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.


Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Chromium, Cadmium, Lead, and Arsenic Concentrations in Water, Vegetables, and Seafood Consumed in a Coastal Area in Northern Vietnam

Environmental Health Insights
Volume 14: 1–9
© The Author(s) 2020
Article reuse guidelines:
sagepub.com/journals-permissions
DOI: 10.1177/1178630220921410



Nguyen Thi Minh Ngoc^{1*}, Nguyen Van Chuyen^{2*},
Nguyen Thi Thu Thao^{1,3}, Nguyen Quang Duc¹, Nguyen Thi Thu Trang²,
Nguyen Thi Thanh Binh¹, Hoang Cao Sa⁴, Nguyen Bao Tran⁵,
Nguyen Van Ba⁶, Nguyen Van Khai¹, Ho Anh Son⁷, Pham Van Han¹,
Elizabeth V Wattenberg⁸, Hiroyuki Nakamura³ and Pham Van Thuc⁹

¹Faculty of Public Health, Haiphong University of Medicine and Pharmacy, Haiphong, Vietnam.

²Department of Military Hygiene, Vietnam Military Medical University, Hanoi, Vietnam. ³Department of Environmental and Preventive Medicine, Kanazawa University, Kanazawa, Japan. ⁴Institute of Health Management Training, Hanoi University of Public Health, Hanoi, Vietnam. ⁵Department of Anatomy, Haiphong University of Medicine and Pharmacy, Haiphong, Vietnam. ⁶Oncology Center, 103 Military Hospital, Vietnam Military Medical University, Hanoi, Vietnam. ⁷Institute of Biomedicine and Pharmacy, Department of Pathophysiology, Vietnam Military Medical University, Hanoi, Vietnam. ⁸School of Public Health, University of Minnesota, Minneapolis, MN, USA. ⁹Department of Clinical Allergy-Immunology, Haiphong University of Medicine and Pharmacy, Haiphong, Vietnam.

ABSTRACT

BACKGROUND: Heavy metal contamination and related risks for the environment and human health are matters of increasing concern.

METHODS: The levels of 4 heavy metals (Cr, Cd, Pb, and As) were evaluated in 2 water types (surface and well), 4 types of seafood (tiger shrimp, stuffed snail, snake-head fish, and catfish), and 27 types of vegetables (12 leafy vegetables, 4 pea plants, 4 tuber vegetables, and 7 herbs) that are commonly consumed in northern coastal communes located in Vietnam. Atomic absorption spectrometry was employed for quantification.

RESULTS: The mean concentrations of heavy metals detected in water, seafood, and vegetable samples exceeded the national permitted standards and World Health Organization (WHO) recommendation values by at least 2-fold, 2.5-fold, and 5-fold for surface water, vegetables, and well water, respectively. The concentrations of all 4 heavy metals detected in seafood samples were higher than the standards. The levels of heavy metals decreased with increasing distance between the sample collection point and the pollution source.

CONCLUSIONS: This is the first report of heavy metal contamination of common sources of food and water in the northern coastal area of Vietnam. Significantly, the concentrations of heavy metals detected in study samples exceeded the regulatory limits. These results underscore the importance of continued monitoring and the development of intervention measures to ensure that the quality of food and water meets established standards and protects the health of the local population.

KEYWORDS: Heavy metals, pollution, water, food contamination, coastal commune, Northern Vietnam

RECEIVED: January 6, 2020. **ACCEPTED:** April 2, 2020.

TYPE: Environmental Health Research and Practice in Vietnam - Original Research

FUNDING: The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: This research was funded by the Vietnam National Key Science and Technology Program period of 2016-2020 under grant number KC10.06/16-20.

DECLARATION OF CONFLICTING INTEREST: The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

CORRESPONDING AUTHOR: Nguyen Thi Minh Ngoc, Hai Phong University of Medicine and Pharmacy, Hai Phong, 04254, Ngo Quyen District, Vietnam.
Email: ntmngoc@hpmu.edu.vn

Introduction

Environmental pollution is one of the major challenges facing modern human society.¹ One of the major threats to ecological and human health is environmental contamination and pollution by heavy metals.^{2,3} Rapid industrialization and urbanization have resulted in contamination of the environment by heavy metals; the rates of mobilization and transport of heavy metals in the environment have greatly accelerated since the 1940s.^{4,5} While some metals are essential for human health at trace concentrations due to their role as coenzymes

(eg, Fe and Cu), others are toxic at any concentration level (eg, Pb and Cd).⁵ Chromium (Cr VI), cadmium (Cd), lead (Pb), and arsenic (As) have been recognized as water, soil, and sediment pollutants in many places in the world.^{6,7} As and other heavy metals have even been detected in soil and sediment from estuarine areas in remote locations.^{7,8} Metals can cause severe toxicity in humans, depending on the concentration, the exposure pathway, and the duration of exposure. It is thus critical to conduct substantial assessment of heavy metal contamination of the environment and to develop strategies to remediate the contamination and protect human and ecological health.⁹

* These authors contributed equally to this work.



Creative Commons Non Commercial CC BY-NC: This article is distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 License (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits non-commercial use, reproduction and distribution of the work without




Figure 1. Map showing the location of the sampling sites.

The toxicity of heavy metals such as Cd, Pb, Cr(VI), and As to humans is widely recognized and documented.^{10,11} For example, the International Agency for Research on Cancer (IARC) classifies As, Cd, and Cr as *Group 1 Carcinogenic to Humans*.¹² Food consumption accounts for up to 80% to 90% of the daily doses of As, Cr, Cd, and Pb to which humans are exposed.¹³ Vegetables, which are an important part of the diet in Vietnam, can take up heavy metals when they are grown in contaminated environments; substantial amounts of heavy metals accumulate in the leaves, and therefore, consumption of vegetables may account for a substantial fraction of total exposures to Cd, Pb, and As.^{14,15} Likewise, seafood, also an important component of the diet in Vietnam, can accumulate high concentrations of heavy metals when aquaculture is conducted in contaminated areas.¹⁴ This indicates that the consumption of vegetables and seafood may be a major source of exposure to heavy metals. Currently, limited data on the levels of heavy metal contamination in food, especially heavy metal contamination in vegetables and seafood in Vietnam, are available.¹⁶

Located on the eastern margin of the Indochinese peninsula, bordering the Gulf of Thailand, and the South China Sea, including Gulf of Tonkin, Vietnam has a long coastline of 3260 km, even excluding its 3000 islands that are dispersed along its entire length.¹⁷ A major environmental disaster occurred in the coastal area of Central Vietnam in 2016 when toxic wastewater containing heavy metals was accidentally, or perhaps illegally, discharged to the coast, causing several billions of dollars of damage.¹⁸ The people and the Government in Vietnam are now well aware of the risk of upstream discharges to the coastal areas, including estuarine areas. This study aims to determine the concentrations of Cd, Cr, Pb, and As in water and measure heavy metals in local vegetables and seafood cultivated in 2 coastal communes in Northern Vietnam.

Methods

Study area

The study was conducted in Tam Hung and Minh Duc communes, 2 coastal sites that belong to the Thủy Nguyên district of Hai Phong city, located in the north of Vietnam, 123 km West of Hanoi capital (Figure 1). The study communes cover an area of 23.12 km²; the population in 2016 was 19036. The per capita income of 2 communes in 2016 was VND43 000 000 (US\$1891).^{19,20}

A program for monitoring metal contamination in water and food (vegetables and seafood) was set up in the Minh Duc and Tam Hung communes, Thủy Nguyên district, Hai Phong city in North Vietnam (Figure 1). Industrial production activities that involve wastewater release into the surrounding environment have been operating in this area since the late 1970s. Use of potentially contaminated stream water may enhance the heavy metal concentrations in vegetables and seafood produced near the industrial production areas. Water, vegetable tissue, and seafood samples were taken from crop fields and aquafarms cultivated by local households at 6 industrial sites including site 1—Trang Kenh stone mining cooperative (20°57'42.9"N 106°44'25.3"E), site 2—MinhDuc chemical company (20°57'39.3"N 106°44'27.9"E), site 3—Chinphon cement plant (20°57'36.0"N 106°45'09.2"E), site 4—HaiPhong thermal power 1 plant (20°56'28.5"N 106°45'34.4"E), site 5—HaiPhong thermal power 2 plant (20°56'21.6"N 106°45'21.9"E), and site 6—NamTrieu shipbuilding company (20°55'59.2"N 106°45'42.6"E).

Sampling

Seafood, vegetable, and water samples were collected in areas within a radius of 500 m, 500 to 1000 m, and 1000 to 1500 m

from factories and enterprises in December 2016. Immediately after the collection, all samples were contained in an airtight insulating box, stored at -4°C , and transported to the Military Hygiene Laboratory, Vietnam Military Academic University for analysis.

Water samples. A total of 54 surface water samples (3 samples were taken at each designated per distance from each of the 6 industrial sources of pollutions) and 222 drinking (well) water samples were collected. The well water samples were collected from households who responded to a survey (administered to 1010 residents) that they use well water. The samples at each position were mixed in a plastic bucket, and a sample of 1 L was stored in a polyethylene bottle. Water samples were acidified with nitric acid to $\text{pH} < 2$ after collecting and transferred on ice to the laboratory for analysis; 10 mL of each sample were filtered through a $0.45\ \mu\text{m}$ Whatman pore-size disposable capsule filter before elemental determination. Surface water and drinking water samples were collected following the regulations for general guidance on water sampling and quality control from the Ministry of Agriculture and Rural Development, Ministry of Health.^{21,22}

Vegetables samples. Samples were collected and processed directly for analysis within 1 to 3 days. Only the edible parts were used for the analysis. Samples included 12 leafy vegetables, 4 pea plants, 4 tuber vegetables, and 7 herbs, which are among the most commonly grown and consumed vegetables in Northern Coastal Vietnam. With 5 replicates per species, a total of 135 samples were collected. Plant samples were collected approximately after 40 to 45 days of sowing.

Seafood samples. A total of 40 seafood samples (10 for each seafood [tiger shrimp, stuffed snails, snake-head fish, and catfish]) were collected at the 2 biggest open markets and the residents' fish ponds in the study area. Immediately after the collection, all samples were placed in an airtight insulating box and transported to the Military Hygiene Laboratory, Military Academic University, Hanoi, Vietnam. Seafood samples were collected by following the regulations on general guidance on food sampling for inspection, quality control, hygiene and safety food published in the Circular No. 14/2011/TT-BYT of the Ministry of Health.²³

Sample preparation

Sample preparation and digestion methods for vegetables and seafood followed Anh TK Bui et al¹⁶ and P Olmedo²⁴ with some modification. Only edible parts of each seafood and vegetable type were used for the analysis. Any spoiled or damaged parts were removed before preparation as a simulation of the real food preparation for human consumption. Fresh samples were weighed and then cut into small portions for drying in an oven at 105°C . Complete drying of the samples was achieved within 72 to 96 hours. Samples are considered dry when they

attained the stable weight. Dry samples were ground in a household coffee grinder and sieved in a 0.8 mm mesh. The fresh weight (fw) and dry weight (dw) were recorded to calculate moisture contents. Trace element concentrations of vegetables and seafood were determined on a dw basis and converted to a fw basis for comparison with the MLs for contaminants and toxins in foods. The samples were then stored in a pre-labeled polyethylene bag for digestion.

After grinding, vegetable and seafood samples (400 mg per each) were digested with 1 mL H_2O_2 (30%, Puriss p.a.; purchased from Merck, Germany), 1 mL HF (40%, Puriss p.a.; purchased from Merck, Germany), and 6 mL HNO_3 (65%, Puriss p.a.; purchased from Merck, Germany) using the Multiwave PRO (Anton Paar) microwave. The microwave was set to 5 minutes for ramping up the temperature to 170°C and then held at this temperature for 10 minutes. The microwave was then set to 15 minutes ramping up to 180°C and then allowed to cool at room temperature. The digested samples were analyzed for As, Cr, Cd, and Pb using atomic absorption spectrophotometry (AAS/Model: AZ3000, Hitachi, Japan). Prepared reagent blanks and digests of the reference material were analyzed in parallel with each type of water, seafood, and vegetables. The AAS system was calibrated using at least 5 standard solutions for external calibration of each metal. Heavy metal concentrations were expressed as milligram per liter (mg/L) for water samples and as milligram per kilogram fresh weight (mg/kg fw.) for vegetable and fish tissues samples. The standard reference material for heavy metals in spinach leaves (SRM 1570a) and in fish (DORM-4) was obtained from the National Institute of Standard and Technology (NIST), USA and the National Research Council of Canada (NRCC), respectively; and standard reference solutions of 1000 mg/L were used for As, Pb, Cr, and Cd from Merck, Germany.

Quality control of the chemical analyses. The blank method was used to implement the quality control of the chemical analyses, performed through the complete preparation and analytical procedure. The blank spike duplicate (BSD) was performed in duplicate to determine to confirm the procedure was working within established control limits.

Great care was taken during sampling and laboratory analysis to prevent sample contamination and to ensure the reliability, quality, and accuracy of the analysis. We avoided the use of metallic tools whenever possible. All glass and plastic ware was washed with soap and rinsed with Sunfua-cromic (50 g $\text{K}_2\text{Cr}_2\text{O}_7$ mix in 500 mL H_2SO_4 98%) and then rinsed thoroughly with distilled and deionized water before use, to ensure there was no contamination of the laboratory supplies and tools; these procedures were conducted in with the guidelines, standards, and protocols of the National Institute of Standards & Technology (NIST).

One calibration blank (HNO_3 1%) was used to establish the analytical curve. Method blanks were used to check possible contamination from the sample preparation procedure. The blank and the standard samples were analyzed. The spike

Table 1. Heavy metal concentrations (mg/L) in water samples (n=276).

METAL	SURFACE WATER (N=54)		WELL WATER (N=222)	
	\bar{X} (MIN, MAX)	TIMES EXCEEDED PERMISSIBLE LIMIT (%)	\bar{X} (MIN, MAX)	TIMES EXCEEDED PERMISSIBLE LIMIT (%)
Cd	0.02 (0.00-0.03)	2.00 (64.81)	0.03 (0.00-0.15)	10 (84.68)
Pb	0.17 (0.03-0.39)	3.40 (87.04)	0.12 (0.01-0.42)	12 (94.59)
As	0.19 (0.02-0.42)	3.80 (68.52)	0.06 (0.01-0.48)	6 (83.33)
Cr	2.56 (0.32-4.32)	5.12 (98.15)	0.25 (0.02-0.82)	5 (74.77)

Abbreviation: WHO, World Health Organization. Allowable limits of Cd, Pb, As, and Cr in surface water recommended by the WHO guideline 2008 and Vietnam Standards 08:2008/BTNMT are 0.01, 0.05, 0.05, and 0.5 mg/L, respectively. Allowable limits of Cd, Pb, As, and Cr in well water recommended by the WHO guideline 2004²⁵ and Vietnam Standards 01:2009/BYT are 0.003, 0.01, 0.01, and 0.05 mg/L, respectively.

method was done in each analysis. Cadmium, lead, arsenic, and chromium levels in the blank samples were lower than limit of detection (LOD). The LOD was calculated by multiplying the SD of minimum values from 5 times repetition running by 3.

Continuing calibration verification was used to determine whether the sample analysis was within control limits (a close linear correlation between the concentration of the standard solution and the absorbance, with a correlation coefficient $r=0.9990$).

The recovery rates were 93.08% to 93.52%, 88.81% to 89.19%, 83.62% to 84.38%, and 93.82% to 94.18% for Cd, Pb, As, and Cr, respectively.

Statistical analysis. Descriptive statistics including mean and standard deviation (SD) were calculated. Differences in the concentration of heavy metals among places were assessed using a one-way analysis of variance (one-way analysis of variance [ANOVA]) with Tukey honestly significant difference (HSD) post hoc test. The statistical significance was set as $P<.05$. All the statistical analyses were done with the IBM SPSS statistics (version 19.0).

Ethical considerations. The protocol of this study was approved by the Scientific and Ethical Committee in Biomedical Research, Hai Phong University of Medicine and Pharmacy. Local commune authorities were asked for their permission and all households were asked for their consent before collecting samples.

Results

Levels of Cd, Pb, As, and Cr in water samples

The levels of Cd, Pb, As, and Cr in the surface water and well water detected in the current study are presented in Table 1. The average concentrations of Cd, Pb, As, and Cr detected in surface water collected from 2 coastal communes in Northern Vietnam ranged between 0.0 and 0.03, 0.03 and 0.39, 0.02 and 0.42, and 0.32 and 4.32 mg/L, respectively, while the concentration of Pb, Cd, As, and Cr in well water ranged from 0.01 to 0.42, 0.0 to 0.15, 0.01 to 0.48, and 0.02 to 0.82 mg/L, respectively (Table 1). In brief, Cr was detected at the highest concentrations, followed by Pb, and then Cd.

The highest mean concentrations of metals detected in both surface and well water were for Cr and the lowest concentrations were observed for Cd. The concentration of As was higher than Pb in surface water, while the adverse result was found in drinking water (well water). The maximum limits (MLs) for Cd, Pb, As, and Cr in surface water according to the Vietnam standards (QCVN 08:2008 BTNMT) are 0.01, 0.05, 0.05, and 0.50 mg/L, respectively. These results indicate that the concentrations of all 4 heavy metals in surface water samples were 2- to 5-fold higher than national standards. The concentration of all 4 heavy metals in well water samples was 5- to 12-fold higher than WHO guidelines and national standards, as the MLs of Cd, Pb, As, and Cr in well water according to the Vietnam standards (QCVN 01:2009 BYT) are 0.003, 0.01, 0.01, and 0.05 mg/L, respectively.

The heavy metal concentrations in water by distance between the sampling sites and the pollution sources are shown in Figure 2. The concentrations of Pb, As, and Cr in surface water samples were statistically reduced in accordance to longer distances between the sampling sites and the pollutants sources ($P<.01$). Cd concentrations in surface water >1000 m from pollution source were statistically lower than the samples taken closer to the pollution source. No difference in Cd concentrations was found in surface water <500 m from the location and 500 to 1000 m distant from pollution sources ($P=.092$).

A statistically significantly lower concentration in As was detected in well water from sampling sites located at longer distances from the pollution source ($P<.001$). The concentration of Pb, Cd, and Cr in well water samples from locations near the pollution source (<500 m) was higher than locations that were 500 to 1000 m from the pollution source ($P<.001$); however, there was no difference in the 3 heavy metals in wells located at 500 to 1000 m and >1000 m from the pollution source ($P=.928$; $P=.998$; $P=.866$ respectively; Figure 2).

Heavy metal in seafood from aquaculture in sampling sites in Thuy Nguyen district, Hai Phong city

Heavy metal concentrations detected in seafood samples are shown in Table 2. The concentration of Cd was higher than




Figure 2. Heavy metal concentration in water by distance between the sampling sites and the pollution sources (n=276). (A) Surface water and (B) Well water. WHO ML indicates World Health Organization maximum limit.

Table 2. Heavy metal concentration (Mean ± SD, mg/kg) in seafood samples.

SEAFOOD	PB	CD	CR	AS
Tiger shrimp	1.24 ± 0.46 ^a	1.62 ± 0.24 ^a	1.46 ± 0.28	0.80 ± 0.01
Stuffed snails	2.13 ± 0.34 ^a	3.26 ± 0.27 ^a	1.69 ± 0.40	1.20 ± 0.01
Snake-head fish	0.08 ± 0.01	2.30 ± 0.26	2.12 ± 0.13	1.18 ± 0.003
Catfish	0.10 ± 0.02	1.06 ± 0.09	2.25 ± 0.15	1.66 ± 0.08

Abbreviation: WHO, World Health Organization; FAO, Food and Agriculture Organization. Allowable limits of Pb, Cd, Cr, and As in seafood recommended by the WHO/FAO, Codex and Vietnam National Technical Regulation are 0.5 mg/kg for Pb and Cd in tiger shrimp; 1.0 and 2.0 mg/kg for stuffed snails, respectively. ^aThe heavy metal concentrations are higher than permitted standards.

those of the other 3 metals in seafood samples except catfish, in which the concentration of Cr exceeded that of the other 3 metals. The concentrations of metals detected in tiger shrimp and stuffed snail samples were the highest for Cd, followed by Pb and Cr.

In contrast to the other types of seafood, in snake-head fish, the order of the heavy metal concentrations was Cd > Cr > As > Pb. The concentrations of Pb and Cd in bottom layer seafood (shrimp and snail) were detected at levels higher than National permissible values and Food and Agriculture Organization (FAO)/(WHO)²⁶ tolerable concentrations, indicating that consuming these seafood products may pose a risk to human health.

Level of heavy metals in vegetables grown in sampling sites in Thuy Nguyen district, Hai Phong city

The concentrations of Pb, Cr, Cd, and As (mg/kg fw) in local vegetables collected from the 6 study sites ranged between 0.11 and 1.96, 0.02 and 1.57, 0.0 and 3.27, 0.17 and 1.70 mg/kg fw, respectively (Table 3). The concentrations of Pb, Cr, and Cd detected in 100% of the vegetable samples were higher than permitted standards. The concentrations of As in the

vegetables samples were lower than the permitted standard except some species of bean (lady’s fingers, green bean, and French bean), red amaranth, and some herb species (rau dang, Laksa leaves, Perilla, and Lettuce). The concentrations of heavy metals across Brassica vegetable samples were the highest for Pb, followed by Cd, As, and then Cr, while the order of heavy metal concentrations in bean vegetable samples was As > Pb > Cr > Cd except lady’s fingers, for which the levels of Cd were higher than Cr. The order of concentration of heavy metals in amaranth was Cd > As > Pb > Cr but in Den tieu Pb was the highest, followed by As and Cd and then Cr.

The highest concentrations of Pb and Cd (mg/kg fw) were found in mustard greens (1.20 and 1.25, respectively; Table 3). The highest concentrations of As (mg/kg fw) were identified in green beans and French beans (1.26).

Figure 3 shows that the average concentrations of As and Pb in vegetables grown further from the sources of pollutant release were statistically lower than vegetables grown nearer to the sources of pollution (P < .001). Cd concentrations in samples from >1000m distance from the pollution sources were lower than those from <500m distance from the pollution sources (P = .024). The Cr concentrations in vegetable grown at <500m from pollution sources were statistically different from vegetables grown farther away, 500 to 1000m (P = .006)

Table 3. Concentration of heavy metal (mg/kg) in grown vegetables (n= 135, 5 samples/species).

LOCAL NAME	ENGLISH NAME	LATIN NAME	PB	CR	CD	AS
Mean (Min-Max)			0.80* (0.11-1.96)	0.51* (0.02-1.57)	0.82 ^a (0.00-3.27)	0.87 (0.17-1.70)
<i>Cải bẹ dung</i>			0.79 ^a	0.09	0.13 ^a	0.72
<i>Cải bẹ xanh</i>	Mustard green	<i>Brassica juncea</i>	1.20 ^a	0.75 ^a	1.25 ^a	0.88 ^a
<i>Cải ngót</i>	Kale	<i>Brassica integrifolia</i>	0.50 ^a	0.19 ^a	0.57 ^a	0.49
<i>Cải xanh</i>	Brown mustard	<i>Brassica juncea</i>	0.80 ^a	0.32 ^a	0.24 ^a	0.89
<i>Dền đỏ</i>	Red amaranth	<i>Amaranthus gangeticus</i>	0.84 ^a	0.50 ^a	1.21 ^a	1.07 ^a
<i>Dền tiểu</i>	Amaranth		0.70 ^a	0.48 ^a	0.65 ^a	0.67
<i>Dền xanh</i>	Green amaranth	<i>Amaranthus viridis</i>	0.66 ^a	0.38 ^a	0.82 ^a	0.71
<i>Mồng tơi</i>	Vine spinach	<i>Basella alba</i>	0.75 ^a	0.40 ^a	0.91 ^a	0.77
<i>Rau đắng</i>			1.00 ^a	0.47 ^a	0.90 ^a	1.07 ^a
<i>Rau đay</i>	Jute mallow	<i>Corchorus olitorius</i>	0.82 ^a	0.53 ^a	0.88 ^a	1.02 ^a
<i>Rau lang</i>	Sweet potato leaf	<i>Ipomoea batatas</i>	0.58 ^a	0.51 ^a	1.01 ^a	0.78
<i>Rau muống</i>	Water spinach	<i>Ipomoea aquatic</i>	0.65 ^a	0.50 ^a	1.21 ^a	0.90
<i>Đậu bắp</i>	Lady's fingers	<i>Abelmoschus esculentus</i>	0.97 ^a	0.58 ^a	0.96 ^a	1.05 ^a
<i>Đậu cô ve</i>	Green bean and French bean	<i>Phaseolus vulgaris</i>	0.96 ^a	0.80 ^a	0.66 ^a	1.24 ^a
<i>Đậu đũa</i>	asparagus bean	<i>Vigna unguiculata</i>	0.73 ^a	0.73 ^a	0.47 ^a	0.90
<i>Đậu rồng</i>	Dragon bean	<i>Psophocarpus tetragonolobus</i>	0.83 ^a	0.84 ^a	0.50 ^a	0.88
<i>Mướp</i>	Sponge gourd	<i>Luffa aegyptiaca</i>	0.63 ^a	0.40 ^a	1.22 ^a	0.69
<i>Cà tím</i>	Long purple eggplants	<i>Solanum melongena</i>	0.70 ^a	0.11 ^a	0.09	0.67
<i>Dưa leo</i>	Cucumber	<i>Cucumis sativus</i>	0.86 ^a	0.67 ^a	1.06 ^a	0.85
<i>Khổ qua</i>	Bitter melon	<i>Momordica charantia</i>	0.83 ^a	0.77 ^a	0.87 ^a	0.86
<i>Diếp cá</i>	Fish mint	<i>Houttuynia cordata</i>	0.67 ^a	0.46 ^a	0.74 ^a	0.75
<i>Húng cây</i>	Indian borage	<i>Plectranthus amboinicus</i>	0.57 ^a	0.44 ^a	1.02 ^a	0.60
<i>Húng quế</i>	Basil	<i>Ocimum basilicum</i>	0.84 ^a	0.42 ^a	0.88 ^a	0.79
<i>Lá lốt</i>	Lolot pepper	<i>Piper lolot</i>	0.72 ^a	0.51 ^a	0.74 ^a	0.80
<i>Rau răm</i>	Laksa leaves	<i>Persicaria odorata</i>	0.87 ^a	0.50 ^a	1.18 ^a	1.02 ^a
<i>Tía tô</i>	Perilla	<i>Perilla frutescens</i>	1.04 ^a	0.80 ^a	0.97 ^a	1.11 ^a
<i>Xà lách</i>	Lettuce	<i>Lactuca sativa</i>	1.10 ^a	0.60 ^a	0.98 ^a	1.16 ^a

Abbreviation: WHO, World Health Organization; FAO, Food and Agriculture Organization. Allowable limits of Pb, Cd, Cr, and As in vegetables recommended by the WHO/FAO, Codex, and Vietnam National Technical Regulation are 0.3, 0.1, 0.1, and 1.0 mg/kg in fw, respectively.

^aThe heavy metal concentrations are higher than permitted standards.

and >1000 m ($P < .001$); however, no difference was seen in vegetables grown 500 to 1000 m and >1000 m from the source of pollution ($P = .545$). The concentrations of Cr, Cd, and Pb in

vegetables were 2- to 10-fold higher than MLs within a 1.5 km radius from pollution sources. Except <500 m distance from pollution source, As was detected below the ML.




Figure 3. Heavy metal contaminations in vegetable samples by distance between sampling sites and pollution sources (n = 135).

Discussion

Peiravi et al²⁷ reported that industrial activities, especially plastic, chemical industries, and metal smelting, are major sources of heavy metals in water. These results led us to measure the levels of 4 heavy metals in water, seafood, and vegetables at a coastal site in North Vietnam where 6 industrial plants are located. Our results indicate that the industrial plants may affect the drinking water quality in this region. Our results also suggest that the stream used as irrigation water for vegetable crop production and seafood aquaculture was contaminated by wastewater from industrial production activities, resulting in elevated heavy metal concentrations in vegetables and seafood from the receiving areas. Altogether, our results indicate that suitable sanitation improvement programs should be used to protect the health of local residents.

The levels of heavy metals in water samples

The detection of heavy metals in water samples from Vietnam has some similarities and differences to the results from studies of heavy metals conducted in other areas. For example, the findings from Table 1 show that in surface water, Cr was detected at the highest concentration, followed by As, and then, Pb and Cd are similar to recent results in Taiwan, which was conducted in surface water in environment near a coastal coal-fired power plant.²⁸ Like surface water, in well water, Cr was detected as the highest concentration and Cd was detected at the lowest concentration. However, the concentration of Pd in well water was higher than the concentration of As. These results are consistent with results of a previous study conducted in the southeast coast of India, which reported a similar trend with Pb and Cd, although the concentrations of Cd and Pb measured in the present study were much higher than those reported in the previous study both pre-monsoon and postmonsoon.²⁹ Likewise, studies conducted in Iran and China also found that Cr was present in the highest concentration, although the concentration of Cr in water samples from Vietnam are higher.^{30,31} By contrast, studies conducted in Thailand, India, Malaysia, and Bangladesh detected Pb at higher concentrations than Cr drinking water.³²⁻³⁵ The As

concentration detected in this study is higher than that detected in HaNam province—a hot spot of arsenic in underground water in Vietnam, which ranged from 8 to 579 ppb (mean 301 ppb), equivalent to 0.008 to 0.579 mg/L, mean 0.301 mg/L.³⁶

Heavy metal in seafood from aquaculture in sampling sites

Heavy metals, which have adverse effects on human health, can accumulate in fish and shrimp tissues, which are generally found in the last zone of the aquatic food chain. Metals are usually taken up from food and water in fish and shrimp, distributed by circulation and eventually accumulate in target organs.³⁷ The lower concentration of Cr detected in shrimp in this study is similar to that observed in the research of Batvari et al who conducted a study in southeast coast of India, whereas the concentrations of Cd was higher than the concentration of Pb in this study; the inverse was observed in studies of other coastal regions in Asia, India, Turkey, and Yemen.^{33,37,38} The As levels in seafood in this study were higher than those presented in other studies, which reported levels ranging from 0.003 to 0.08 µg/g and from 0.021 to 0.048 µg/g in common carp.³⁹

The trend of As levels being higher than Pb levels in fish tissues in our study is consistent with results reported from Bangladesh, which showed the mean concentration of As and Pb to be 1.59 and 1.13 mg/kg in summer and 1.81 and 1.45 mg/kg in winter, respectively.⁴⁰ Interestingly, Peshut et al⁴¹ found that arsenic in marine organisms or seafood is mainly in various organic forms, such as arsenobetaine, arsenoribosides, and arsenocholine, which are effectively nontoxic.

The present study results, which found the order of heavy metal concentrations to be Cr > As > Pb > Cd, are different from the findings from Le Quang Dung et al,⁴² which reported the order of heavy metal concentration in oysters along the Hai Phong-Ha Long coast was As > Cd > Pb > Cr with concentrations of 10.10 to 19.33, 3.53 to 12.74, 0.79 to 6.20, and 0.81 to 4.47 mg/kg dry wt, respectively.

Heavy metal in vegetables in sampling sites

Heavy metal contamination in vegetable samples is presented in Table 3. Levels of Pb in mustard greens in this study were higher than in other studies.^{11,15,16,43,44} Pb-Zn mining nearby results in contamination of soil and irrigation water.¹⁶ Fertilizer and other agrochemicals, atmospheric deposition, and irrigation with contaminated water have also been implicated in Pb contamination of crops.⁴⁴

Similarly high Cd concentrations were found in leafy vegetables sampled at Dabaoshan mine⁴⁵ and at Bac Kan.¹⁶ Other studies have found lower^{6,46} and higher^{47,48} Cd concentrations in vegetables compared to those found in our study.

The highest concentrations of As (mg/kg fw) were identified in green beans and French beans (1.26). There have been very few studies that explored As content in vegetables, and our

results were much higher than those reported in another study conducted in Pearl River Delta, South China⁴³ but much lower than those found in the study carried out in Bac Kan, Vietnam.¹⁶

Figure 3 shows the trace element contamination in vegetable samples by distance between sampling sites and pollution sources. Only the As concentration in vegetables grown at a distance >500 m from the pollution source is lower than ML. It is important to consider the health risks from heavy metal contamination of vegetables to the local community because these vegetables are the most favorite foods consumed in Vietnam, especially in the North.

Limitations of the study

To the best of our knowledge, this is the first study to measure heavy metal contamination in water, seafood, and vegetables collected from coastal communes in Northern Vietnam. However, the current data represented only a cross-sectional snapshot. We did not consider the change of daily weather during sampling period to avoid the possible bias in the analysis. Moreover, this present study did not include a health risk assessment to predict the population health problems related to environmental factors. Further studies are needed to investigate the impact of the heavy metals in water, vegetables, and seafood on human health.

Conclusions

In summary, our results indicate that water sources (*surface, drinking*) and food (*vegetables, seafood*) are contaminated with heavy metals (*arsenic, lead, cadmium, chromium*). The concentrations of heavy metal in water and food samples decreased with the distance between where the samples were collected and the pollution sources, especially for arsenic and lead. The research results show that the use of water and food grown near the pollution sources may increase public health risks due to heavy metals.

Recommendations

Considering all available evidence, we do not recommend field cultivation and consumption of vegetables grown 1 km from the sources of pollution. Further studies need to be conducted not only to calculate the risks, especially the cumulative risk resulting from consumption of contaminated water, food, and seafood, but also to apply appropriate measures to ensure the quality of food and water sources to protect the health of the local community.

Acknowledgements

The authors are grateful to local authorities and grassroots of study sites and colleagues from Vietnam Military Medical University and Hai Phong University of Medicine and Pharmacy for their supports during data collection and measurement.

Author Contributions

NTMN, NVC, PVH, HAS, and PVT conceived and designed the study, agreed with the results and conclusions, and came up with arguments for this manuscript. NTMN, NVC, HCS, NQD, and NBT collected data and samples in the field. NTTThao, NTTB, NTTTrang, and NQD analyzed the data. NTMN, NVC, NTTThao, NQD, EVW, and HN wrote the first draft of the paper. All the authors read, made revisions, and agreed on the final version of manuscript. NTMN, NTTThao, and PVT reviewed and approved the final manuscript for submission.

ORCID iD

Nguyen Thi Minh Ngoc  <https://orcid.org/0000-0002-7652-659X>

REFERENCES

1. Ali H, Khan E. Environmental chemistry in the twenty-first century. *Environ Chem Lett.* 2017;15:329-346. doi:10.1007/s10311-016-0601-3.
2. Ali H, Khan E, Sajad MA. Phytoremediation of heavy metals—concepts and applications. *Chemosphere.* 2013;91:869-881. doi:10.1016/j.chemosphere.2013.01.075.
3. Hashem MA, Nur-A-Tomal MS, Mondal NR, Rahman MA. Hair burning and liming in tanneries is a source of pollution by arsenic, lead, zinc, manganese and iron. *Environ Chem Lett.* 2017;15:501-506. doi:10.1007/s10311-017-0634-2.
4. Khan FU, Rahman AU, Jan A, Riaz M. Toxic and trace metals (Pb, Cd, Zn, Cu, Mn, Ni, Co and Cr) in dust, dustfall/soil. *J Chem Soc Pakistan.* 2004;26:453-456.
5. Merian E. Introduction on environmental chemistry and global cycles of chromium, nickel, cobalt, beryllium, arsenic, cadmium and selenium, and their derivatives. *Toxicol Environ Chem.* 1984;8:9-38. doi:10.1080/02772248409357038.
6. Nations U, Osmá E, Serin M, et al. Arsenic concentrations and speciation in a temperate mangrove ecosystem, NSW, Australia. *Appl Organomet Chem.* 2007;9:192-201. doi:10.1016/s0927-5215(06)09021-7.
7. MacFarlane GR, Pulkownik A, Burchett MD. Accumulation and distribution of heavy metals in the grey mangrove, *Avicennia marina* (Forsk.) Vierh.: biological indication potential. *Environ Pollut.* 2003;123:139-151. doi:10.1016/S0269-7491(02)00342-1.
8. Tue NT, Quy TD, Amano A, et al. Historical profiles of trace element concentrations in Mangrove sediments from the Ba Lat Estuary, Red River, Vietnam. *Water Air Soil Pollut.* 2012;223:1315-1330. doi:10.1007/s11270-011-0947-x.
9. Harmanescu M, Alda LM, Bordean DM, Gogoasa I, Gergen I. Heavy metals health risk assessment for population via consumption of vegetables grown in old mining area; a case study: Banat County, Romania. *Chem Cent J.* 2011;5:64. doi:10.1186/1752-153X-5-64.
10. Järup L. Hazards of heavy metal contamination. *Br Med Bull.* 2003;68:167-182. doi:10.1093/bmb/ldg032.
11. Krejpcio Z, Sionkowski S, Bartela J. Safety of fresh fruits and juices available on the Polish market as determined by heavy metal residues. *Polish J Environ Stud.* 2005;14:877-881.
12. WHO. Lead poisoning and health. <https://www.who.int/news-room/fact-sheets/detail/lead-poisoning-and-health>. Accessed December 25, 2019.
13. Jaishankar M, Tseten T, Anbalagan N, Mathew BB, Beeregowda KN. Toxicity, mechanism and health effects of some heavy metals. *Interdiscip Toxicol.* 2014;7:60-72. doi:10.2478/intox-2014-0009.
14. Hu J, Wu F, Wu S, Cao Z, Lin X, Wong MH. Bioaccessibility, dietary exposure and human risk assessment of heavy metals from market vegetables in Hong Kong revealed with an in vitro gastrointestinal model. *Chemosphere.* 2013;91:455-461. doi:10.1016/j.chemosphere.2012.11.066.
15. Kananke T, Wansapala J, Gunaratne A. Heavy metal contamination in green leafy vegetables collected from selected market sites of Piliyandala Area, Colombo District, Sri Lanka. *Am J Food Sci Technol.* 2014;2:139-144. doi:10.12691/ajfst-2-5-1.
16. Bui ATK, Nguyen HTH, Nguyen MN, et al. Accumulation and potential health risks of cadmium, lead and arsenic in vegetables grown near mining sites in Northern Vietnam. *Environ Monit Assess.* 2016;188:525. doi:10.1007/s10661-016-5535-5.

17. Cong L, Van Cu N, Van Shibayama T. *Assessment of Vietnam Coastal Erosion and Relevant Laws and Policies*. Elsevier; 2014. doi:10.1016/B978-0-12-800007-6.00004-6.
18. Herreros MA, Inigo-Nunez S, Sanchez-Perez E, Encinas T, Gonzalez-Bulnes A. Contribution of fish consumption to heavy metals exposure in women of childbearing age from a Mediterranean country (Spain). *Food Chem Toxicol*. 2008;46:1591-1595. doi:10.1016/j.fct.2007.12.024.
19. Minh Duc Commune People Committee. Review report on the result of implementing tasks in 2016 [in Vietnamese].
20. Tam Hung Commune People Committee. Review report on the result of implementing tasks in 2016 [in Vietnamese].
21. QCVN01-132:2013/BNNPTNT. *National Technical Regulation on Fresh Vegetable, Fruit and Tea—Conditions for Ensuring Food Safety in Production and Packing*. <https://vanbanphapluat.co/qcvn-01-132-2013-bnnptnt-rau-qua-che-bup-tuoi-du-dieu-kien>. Accessed December 13, 2019.
22. VMH. QCVN 01:2009. National technical regulation on drinking water quality. <http://www.fsi.org.vn/pic/files/qcvn-01-2009-byt-vc-nuoc-an-uong.pdf>. Accessed December 13, 2019.
23. QCVN8-2:2011/BYT. 2011 Vietnam national technical regulation on the limits of heavy metals contamination in food. http://www.fsi.org.vn/pic/files/qcvn-8-2_2011-byt-gioi-han-o-nhiem-kim-loai-nang.pdf. Accessed December 13, 2019.
24. Olmedo P, Pla A, Hernández AF, Barbier F, Ayouni L, Gil F. Determination of toxic elements (mercury, cadmium, lead, tin and arsenic) in fish and shellfish samples. Risk assessment for the consumers. *Environ Int*. 2013;59:63-72. doi:10.1016/j.envint.2013.05.005.
25. WHO. *Guidelines for Drinking-Water Quality*. 3rd ed. Geneva, Switzerland: WHO; 2004.
26. FAO/WHO 1995. General standard for contaminants and toxins in food and feed (Codex Stan 193-1995). *J Ekon Malaysia*. 2015;51:39-54. doi:10.1017/CBO9781107415324.004.
27. Peiravi R, Alidadi H, Dehghan AA, Vahedian M. Heavy metals concentrations in Mashhad drinking water network. <https://www.sid.ir/en/Journal/ViewPaper.aspx?ID=337508>. Published 2013. Accessed March 29, 2020.
28. Kok VC, Winn PR, Hsieh Y-J, Chien J-W, Yang J-M, Yeh G-P. A pilot survey of potentially hazardous trace elements in the aquatic environment near a coastal coal-fired power plant in Taiwan. *Environ Health Insights*. 2019;13:1178630219862236. doi:10.1177/1178630219862236.
29. Batvari BPD, Sivakumar S, Shanthi K, et al. Heavy metals accumulation in crab and shrimps from Pulicat lake, north Chennai coastal region, southeast coast of India. *Toxicol Ind Health*. 2016;32:1-6. doi:10.1177/0748233713475500.
30. Rajeshkumar S, Liu Y, Zhang X, Ravikumar B, Bai G, Li X. Studies on seasonal pollution of heavy metals in water, sediment, fish and oyster from the Meiliang Bay of Taihu Lake in China. *Chemosphere*. 2018;191:626-638. doi:10.1016/j.chemosphere.2017.10.078.
31. Alidadi H, Tavakoly Sany SB, Zarif Garaati Oftadeh B, Mohamad T, Shamszade H, Fakhari M. Health risk assessments of arsenic and toxic heavy metal exposure in drinking water in northeast Iran. *Environ Health Prev Med*. 2019;24:59. doi:10.1186/s12199-019-0812-x.
32. Wongsasulak P, Chotpantarat S, Siriwong W, Robson M. Heavy metal contamination and human health risk assessment in drinking water from shallow ground-water wells in an agricultural area in Ubon Ratchathani province, Thailand. *Environ Geochem Health*. 2014;36:169-182. doi:10.1007/s10653-013-9537-8.
33. Bajwa BS, Kumar S, Singh S, Sahoo SK, Tripathi RM. Uranium and other heavy toxic elements distribution in the drinking water samples of SW-Punjab, India. *J Radiat Res Appl Sci*. 2017;10:13-19. doi:10.1016/j.jrras.2015.01.002.
34. Ahmad N, Jaafar MS, Alsaffar MS. Study of radon concentration and toxic elements in drinking and irrigated water and its implications in Sungai Petani, Kedah, Malaysia. *J Radiat Res Appl Sci*. 2015;8:294-299. doi:10.1016/j.jrras.2015.04.003.
35. Saha N, Rahman MS, Ahmed MB, Zhou JL, Ngo HH, Guo W. Industrial metal pollution in water and probabilistic assessment of human health risk. *J Environ Manage*. 2017;185:70-78. doi:10.1016/j.jenvman.2016.10.023.
36. Huy TB, Tuyet-Hanh TT, Johnston R, Nguyen-Viet H. Assessing health risk due to exposure to arsenic in drinking water in Hanam province, Vietnam. *Int J Environ Res Public Health*. 2014;11:7575-7591. doi:10.3390/ijerph110807575.
37. Aytekin T, Kargin D, Temiz O, Sa G, Varkal H, Kargin F. Accumulation and health risk assessment of heavy metals in tissues of the shrimp and fish species from the Yumurtalik coast of Iskenderun Gulf, Turkey. *Heliyon*. 2017;5:e02131. doi:10.1016/j.heliyon.2019.e02131.
38. Saleh YS, Marie MAS. Assessment of metal contamination in water, sediment, and tissues of Arius thalassinus fish from the Red Sea coast of Yemen and the potential human risk assessment. *Environ Sci Pollut Res Int*. 2015;22:5481-5490. doi:10.1007/s11356-014-3780-0.
39. Zrnčić S, Oraić D, Čaleta M, Mihaljević Ž, Zanella D, Bilandžić N. Biomonitoring of heavy metals in fish from the Danube River. *Environ Monit Assess*. 2013;185:1189-1198. doi:10.1007/s10661-012-2625-x.
40. Ali MM, Ali ML, Proshad R, et al. Heavy metal concentrations in commercially valuable fishes with health hazard inference from Karnaphuli River, Bangladesh [published online ahead of print October 25, 2019]. *Hum Ecol Risk Assess*. doi:10.1080/10807039.2019.1676635.
41. Peshut PJ, Morrison RJ, Brooks BA. Arsenic speciation in marine fish and shellfish from American Samoa. *Chemosphere*. 2008;71:484-492. doi:10.1016/j.chemosphere.2007.10.014.
42. Dung LQ, Linh NV, Tu VV. Determination of heavy metal accumulation in rock oyster and green mussel from Haiphong coast. *J Vietnam Sci Technol of Agri and Rural Development*. 2013;6:81-89.
43. Chang CY, Yu HY, Chen JJ, Li FB, Zhang HH, Liu CP. Accumulation of heavy metals in leaf vegetables from agricultural soils and associated potential health risks in the Pearl River Delta, South China. *Environ Monit Assess*. 2014;186:1547-1560. doi:10.1007/s10661-013-3472-0.
44. Oteef MDY, Fawy KF, Abd-Rabboh HSM, Idris AM. Levels of zinc, copper, cadmium, and lead in fruits and vegetables grown and consumed in Aseer Region, Saudi Arabia. *Environ Monit Assess*. 2015;187:676. doi:10.1007/s10661-015-4905-8.
45. Zhuang P, Zou B, Li NY, Li ZA. Heavy metal contamination in soils and food crops around Dabaoshan mine in Guangdong, China: implication for human health. *Environ Geochem Health*. 2009;31:707-715. doi:10.1007/s10653-009-9248-3.
46. Kraja F, Kraja B, Mone I, Harizi I, Babameto A, Burazeri G. Self-reported prevalence and risk factors of non-communicable diseases in the Albanian adult population. *Med Arch*. 2016;70:208-212. doi:10.5455/medarh.2016.70.208-212.
47. Maleki A, Zarasvand MA. Heavy metals in selected edible vegetables and estimation of their daily intake in Sanandaj, Iran. *Southeast Asian J Trop Med Public Health*. 2008;39:335-340. <http://www.ncbi.nlm.nih.gov/pubmed/18564723>. Accessed December 13, 2019.
48. Li N, Kang Y, Pan W, Zeng L, Zhang Q, Luo J. Concentration and transportation of heavy metals in vegetables and risk assessment of human exposure to bioaccessible heavy metals in soil near a waste-incinerator site, South China. *Sci Total Environ*. 2015;521-522:144-151. doi:10.1016/j.scitotenv.2015.03.081.