

Harmonizing Goals for Agricultural Intensification and Human Health Protection in Sub-Saharan Africa

Authors: Trimmer, John T., Bauza, Valerie, Byrne, Diana M., Lardizabal, Amanda, and Guest, Jeremy S.

Source: Tropical Conservation Science, 10(1)

Published By: SAGE Publishing

URL: <https://doi.org/10.1177/1940082917720666>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Harmonizing Goals for Agricultural Intensification and Human Health Protection in Sub-Saharan Africa

Tropical Conservation Science
Volume 10: 1–6
© The Author(s) 2017
Reprints and permissions:
sagepub.com/journalsPermissions.nav
DOI: 10.1177/1940082917720666
journals.sagepub.com/home/trc



John T. Trimmer¹, Valerie Bauza¹, Diana M. Byrne¹,
Amanda Lardizabal¹, and Jeremy S. Guest¹

Abstract

Increased agricultural production will be necessary to feed rapidly growing populations in sub-Saharan Africa, where many households currently practice low-input, subsistence farming. Efforts to expand food production will likely include agricultural intensification to enhance productivity of existing cropland, and holistic frameworks are needed to quantitatively evaluate trade-offs and synergies between intensification and other dimensions of development. Beyond well-documented interactions with environmental and economic issues, intensification's complex relationship with human health should take a position of primary importance in any framework designed to advance food security. While intensification can lead to improvements in nutritional status, neglecting sources of potential adverse health impacts, including water source contamination and direct contact with agricultural inputs or environmental pathogens, may undermine prospective gains. Harmonizing goals will require interdisciplinary teams applying frameworks that integrate tools such as quantitative risk assessment, environmental life cycle assessment, and economic models to comprehensively evaluate potentially dissimilar strategies across common metrics while accounting for interdependencies and uncertainties. With local implementation partners, these teams will be well-equipped to develop holistic interventions that effectively promote food security and protect human health while considering local constraints and opportunities across multiple dimensions of development.

Keywords

intensification, human, health, Africa, agriculture

Placing Agricultural Intensification Within a Broader Framework

To feed the growing global population, current agricultural production must double by 2050 (Godfray et al., 2010). Increased production will be particularly crucial in sub-Saharan Africa (SSA), where populations are projected to grow from 1.0 billion in 2017 to 1.9 to 2.3 billion in 2050 (United Nations, 2015), and where approximately 218 million people are currently undernourished (Food and Agriculture Organization of the United Nations, 2015) with increased risk of morbidity and mortality (Bain et al., 2013; de Onis & Branca, 2016). For example, among Ghanaian children younger than 5 years, 23% were stunted and 57% were afflicted with anemia in 2011 (United States Agency for International Development, 2014). The poorest households in SSA typically rely on smallholder farmers, themselves

hovering precariously on the threshold of food insecurity, who practice subsistence agriculture characterized by low chemical inputs and high labor requirements (Frelat et al., 2016; Goldsmith, 2017; van Ittersum et al., 2016; Tscharnitke et al., 2012). Considering agriculture already represents the largest land use on the planet (Foley et al., 2011), one plausible path to increasing food production involves agricultural intensification (Palm et al., 2017; Spera, 2017), which enhances productivity of existing

¹Department of Civil and Environmental Engineering, University of Illinois at Urbana-Champaign, IL, USA

Received 30 May 2017; Accepted 30 May 2017

Corresponding Author:

Jeremy S. Guest, Department of Civil and Environmental Engineering, University of Illinois at Urbana-Champaign, 205 N. Mathews Ave., 3221 Newmark Civil Engineering Laboratory, Urbana, IL 61801, USA.
Email: jsguest@illinois.edu



Creative Commons Non Commercial CC BY-NC: This article is distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 License (<http://www.creativecommons.org/licenses/by-nc/4.0/>) which permits non-commercial use, reproduction and distribution of the work without further permission provided the original work is attributed as specified on the SAGE and Open Access pages (<https://us.sagepub.com/en-us/nam/open-access-at-sage>).

cropland through increased cropping intensity, irrigation, fertilizer and pesticide use, and monoculture (Foley et al., 2011; Matson, Parton, Power, & Swift, 1997; van Ittersum et al., 2016).

Achieving the greatest benefits through agricultural intensification requires considering its interactions with other dimensions of development (Garnett et al., 2013). Perhaps the most well-established connections concern its negative environmental impacts, including greenhouse gas emissions, water scarcity due to irrigation demands and climate change, soil erosion, pesticide resistance, and highly altered nitrogen and phosphorus cycling (Alliance for a Green Revolution, 2014; Foley et al., 2011; Matson et al., 1997; Kassie, Teklewold, Jaleta, Marenja, & Erenstein, 2015; Steffen et al., 2015; Tilman, Cassman, Matson, Naylor, & Polasky, 2002; West et al., 2014), suggesting that any agricultural transition should focus on resource efficiency and conservation along with improved yields (Mueller et al., 2012; van Ittersum et al., 2016). From an economic standpoint, offering households greater opportunities to diversify and safeguard their livelihoods through improved market access, off-farm employment, and rural infrastructure may be at least as important as agricultural intensification in reducing rural poverty (Frelat et al., 2016; Godfray et al., 2010; Rosegrant & Cline, 2003; van Ittersum et al., 2016). Beyond these considerations, the relationship between agricultural intensification and human health should take a position of primary importance in any framework designed to advance food security. While substantial improvements in nutrition, health status, and economic well-being could be achieved through intensification, neglecting sources of potential adverse health impacts could create cycles that undermine prospective health gains.

Elucidating Connections Between Human Health and Agricultural Intensification

While agricultural intensification has the potential to contribute to increased food security and improved health outcomes, it can also open various risk pathways that may compromise human health in other ways. For example, one aspect of intensification involves farmers using greater quantities and application rates of pesticides, but many widely-used pesticides are known to have negative health effects, both acute (e.g., enzyme inhibition causing vomiting, diarrhea, or respiratory failure) and chronic (e.g., endocrine disruption, neuropsychiatric impairment, heart and liver damage, and multiple cancers; Swiss Federal Institute of Aquatic Science and Technology [EAWAG], 2016). Moreover, the most hazardous pesticides, banned in many high-income countries, continue to be stockpiled and circulated in SSA, where workers who may have limited

access to protective equipment also may not be informed of proper handling procedures or health impacts (EAWAG, 2016; Jepson et al., 2014; Sheahan, Barrett, & Goldvale, 2017). Pesticide exposure can occur through numerous pathways, including through consuming foods retaining pesticide residues and drinking water contaminated by agricultural runoff (Akoto, Andoh, Darko, Eshun, & Osei-Fosu, 2013; Akoto, Gavor, Appah, & Apau, 2015; Bempah, Donkor, Yeboah, Dubey, & Osei-Fosu, 2011; Fosu-Mensah, Okoffo, Darko, & Gordon, 2016). For farmers and their families in SSA, the highest risks may come from direct contact or inhalation during pesticide application, mixing, and in-home storage, due to improper handling and insufficient protective equipment (EAWAG, 2016; Jepson et al., 2014).

Increased fertilizer application can lead to another set of health risks, although their severity likely depends on local soil conditions and nutrient retention (Neill et al., 2017; Palm et al., 2017). Along with increased eutrophication potential (Smith, Tilman, & Nekola, 1999), surface water or groundwater sources contaminated with agricultural runoff or leachate can exhibit elevated nitrate and nitrite concentrations (Matson et al., 1997; Tilman, 1999), which are connected with methemoglobinemia (blue baby syndrome; Fewtrell, 2004; Sadler et al., 2016). Possible connections with birth defects and various cancers have also been reported (Brender et al., 2013; Sadler et al., 2016; Ward et al., 2005), although these links remain uncertain and have not been incorporated into international drinking water guidelines (World Health Organization, 2011). Furthermore, if animal manure or human waste are used as supplementary nutrient sources, the risk of pathogenic exposure through direct contact or drinking water contamination may be significant (Gerba & Smith, 2005; Tyrrel & Quinton, 2003).

Pathogenic exposure and infection, while not always directly related to agriculture, can undermine the benefits of increased food availability by affecting human nutrient absorption. Enteric infections, including diarrheal diseases, soil-transmitted helminths, and environmental enteric dysfunction (EED), can reduce intestinal absorption and divert nutrients away from growth to immune response, contributing to a cycle in which infection causes worsened nutritional status and leading to greater susceptibility for further infection (de Onis & Branca, 2016; Guerrant, Oriá, Moore, Oriá, & Lima, 2008; Korpe & Petri, 2012). Acute exposure to infected feces can cause diarrheal and soil-transmitted helminth infections, while chronic fecal pathogen exposure is hypothesized to be the primary cause of EED (Korpe & Petri, 2012; Trehan, Kelly, Shaikh, & Manary, 2016). Exposure to environmental chemicals, including pesticides, may also contribute to EED (Mapesa, Maxwell, & Ryan, 2016). Enteric infections that reduce nutrient absorption

and impair immune function may play a substantial role in malnutrition, as gold standard nutrition interventions are estimated to reduce child stunting by only one third (Bhutta et al., 2008). Therefore, reducing enteric infections through water, sanitation, and hygiene (WASH) interventions may enable greater realization of the nutritional benefits of increased food security.

Harmonizing Goals and Moving Toward Holistic Interventions

To achieve lasting and consequential gains in nutritional security, efforts toward agricultural intensification should account for interactions with health and other-related dimensions of development (e.g., water and sanitation, education, energy; Sanchez, Denning, & Nziguheba, 2009; Wallington & Cai, 2017). In this effort, planners must identify and understand diverse, locally relevant, and often uncertain variables and decisions, such as those related to hydrology, climate, local soil quality, chemical storage and application practices, the types of fertilizers and pesticides available, knowledge and applicability of organic farming practices, and water sources (Godfray et al., 2010; Michelson, 2017; Mueller et al., 2012). Harmonizing goals for improved human health and increased agricultural productivity constitute a core aspect of this endeavor and can encompass a variety of possibilities. Holistic approaches explicitly integrating improved food security, nutrition, and WASH could generate synergistic benefits by breaking the cycle surrounding enteric infections. The results of current trials (Arnold et al., 2013; Humphrey et al., 2015) testing nutrition and WASH interventions' individual and combined effects may provide strong evidence for future integration across these dimensions of development. In addition, strategies already being explored to address the environmental concerns of intensification could provide simultaneous human health protection. Techniques including precision agriculture, reduced tillage, contour farming, crop rotation, polyculture, agroforestry, and improved farmer education on crop-specific nutrient requirements and pesticide application have the potential to limit fertilizer and pesticide use, reduce environmental contamination, and improve crop resilience to climate change, thereby lessening associated health risks (Godfray et al., 2010; Goldsmith, 2017; Matson et al., 1997; Mueller et al., 2012; Waldron et al., 2017).

Given the broad range and varied effects of possible strategies, maximizing intended benefits and minimizing unintended consequences require structured frameworks capable of identifying and analyzing trade-offs and synergies across dimensions of development (Wallington & Cai, 2017). For example, pesticide use is correlated with improved agricultural output and income, but it can also cause adverse health outcomes that lead to increased

health-care costs and time lost due to illness (Sheahan et al., 2017). As another illustration, intensification may reduce crop diversity on smallholder farms, potentially diminishing dietary diversity and its nutritional benefits (Jones, Shrinivas, & Bezner-Kerr, 2014; Pellegrini & Tasciotti, 2014), although access to new and better markets may counteract these dietary losses (Sibhatu, Krishna, & Qaim, 2015) and enhance food security and rural development (Endres & Endres, 2017). Quantitative risk assessment (QRA) frameworks (Haas, Rose, & Gerba, 2014), applicable to chemical and microbial contaminants and various exposure pathways, may be particularly appropriate in ensuring that human health is protected. QRA provides a proactive approach capable of estimating the effects of potential interventions and suggesting improvements before any health hazards occur. Furthermore, QRA can be integrated with other tools, such as environmental life cycle assessment, to simultaneously evaluate outcomes and consider trade-offs across multiple dimensions of development (Kobayashi, Peters, Ashbolt, Shiels, & Khan, 2015; Kobayashi, Peters, & Khan, 2015). Linking QRA and life cycle assessment with economic models that estimate altered labor availability, health-care expenditures, and income from correlations with crop yields, food security, and health status (Sheahan et al., 2017) could lead to comprehensive frameworks able to compare dissimilar interventions across common metrics, accounting for the interdependencies that connect agriculture, environmental sustainability, economics, and health (Brawn, 2017; Cohn, 2017; Waldron et al., 2017; Wallington & Cai, 2017).

Any multifaceted framework or intervention must additionally consider a wide array of social, political, economic, and historical dimensions while accounting for various spatial scales, value systems, local constraints, and variabilities inherent in relevant variables and decisions (Endres & Endres, 2017; Garnett et al., 2013; Godfray et al., 2010; Palm et al., 2017; Sheahan et al., 2017; Tilman et al., 2002; Waldron et al., 2017). Building analytical models that allow for stochastic variables, randomly generated simulations, stakeholder involvement, and multicriteria decision analysis (Mendoza & Martins, 2006) will be particularly useful in addressing the uncertainties surrounding various parameters (e.g., health outcomes) and comprehensively evaluating context-specific strategies with disparate aspects across multiple scenarios. This level of complexity suggests a pressing need for dynamic interdisciplinary research teams capable of thinking holistically to harmonize multiple, qualitatively distinct goals. As evidenced by the connections and risks outlined above, any such team focused on improving agricultural productivity should include environmental and health specialists, as well as experts on complementary topics such as government policy, market access and employment, rural infrastructure,

sociology, and gender equity. With local implementation partners (Palm et al., 2017), these diverse teams are well-equipped to develop holistic interventions that promote food security, protect health, and produce positive, appropriate, and enduring change.

Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: This research is in part funded by the USAID Feed the Future Lab for Soybean Value Chain Research. The award number is Cooperative Agreement Number AID-OAA-L-14-00001.

References

- Alliance for a Green Revolution. (2014). *Africa agriculture status report: Climate change and smallholder agriculture in sub-Saharan Africa*. Retrieved from <https://agra.org/wp-content/uploads/2016/04/africa-agriculture-status-report-2014.pdf>
- Akoto, O., Andoh, H., Darko, G., Eshun, K., & Osei-Fosu, P. (2013). Health risk assessment of pesticides residue in maize and cowpea from Ejura, Ghana. *Chemosphere*, *92*(1): 67–73. Retrieved from <https://doi.org/10.1016/j.chemosphere.2013.02.057>
- Akoto, O., Gavor, S., Appah, M. K., & Apau, J. (2015). Estimation of human health risk associated with the consumption of pesticide-contaminated vegetables from Kumasi, Ghana. *Environmental Monitoring and Assessment*, *187*(5): 244. Retrieved from <https://doi.org/10.1007/s10661-015-4471-0>
- Arnold, B. F., Null, C., Luby, S. P., Unicomb, L., Stewart, C. P., Dewey, K. G., . . . Colford, J. M. (2013). Cluster-randomised controlled trials of individual and combined water, sanitation, hygiene and nutritional interventions in rural Bangladesh and Kenya: the WASH Benefits study design and rationale. *BMJ Open*, *3*(8): e003476. Retrieved from <https://doi.org/10.1136/bmjopen-2013-003476>
- Bain, L. E., Awah, P. K., Geraldine, N., Kindong, N. P., Siga, Y., Bernard, N., & Tanjeko, A. T. (2013). Malnutrition in Sub-Saharan Africa: Burden, causes and prospects. *Pan African Medical Journal*, *15*(1). Retrieved from <https://www.ajol.info/index.php/pamj/article/view/100127>
- Bempah, C. K., Donkor, A., Yeboah, P. O., Dubey, B., & Osei-Fosu, P. (2011). A preliminary assessment of consumer's exposure to organochlorine pesticides in fruits and vegetables and the potential health risk in Accra Metropolis, Ghana. *Food Chemistry*, *128*(4): 1058–1065. Retrieved from <https://doi.org/10.1016/j.foodchem.2011.04.013>
- Bhutta, Z. A., Ahmed, T., Black, R. E., Cousens, S., Dewey, K., Giugliani, E., . . . Shekar, M. (2008). What works? Interventions for maternal and child undernutrition and survival. *The Lancet*, *371*(9610): 417–440. Retrieved from [https://doi.org/10.1016/S0140-6736\(07\)61693-6](https://doi.org/10.1016/S0140-6736(07)61693-6)
- Brawn, J. D. (2017). Implications of agricultural development for tropical biodiversity. *Tropical Conservation Science*, *10*.
- Breder, J. D., Weyer, P. J., Romitti, P. A., Mohanty, B. P., Shinde, M. U., Vuong, A. M., . . . Canfield, M. A. (2013). Prenatal nitrate intake from drinking water and selected birth defects in offspring of participants in the national birth defects prevention study. *Environmental Health Perspectives*, *121*(9): 1083–1089. Retrieved from <https://doi.org/10.1289/ehp.1206249>
- Cohn, A. (2017). Leveraging climate regulation by ecosystems for agriculture to promote ecosystem stewardship. *Tropical Conservation Science*, *10*.
- de Onis, M., & Branca, F. (2016). Childhood stunting: A global perspective. *Maternal & Child Nutrition*, *12*, 12–26. Retrieved from <https://doi.org/10.1111/mcn.12231>
- Endres, A. B., & Endres, R. (2017). The European Union, agriculture, and the tropics: Public financial incentives to enhance food security and expansion of production contracts. *Tropical Conservation Science*, *10*.
- Food and Agriculture Organization of the United Nations. (2015). *Regional overview of food insecurity: African food security prospects brighter than ever*. Retrieved from <http://www.fao.org/publications/card/en/c/9186338b-3fe6-427c-b805-fe81be23a676/>
- Fewtrell, L. (2004). Drinking-Water nitrate, methemoglobinemia, and global burden of disease: A discussion. *Environmental Health Perspectives*, *112*(14): 1371–1374.
- Foley, J. A., Ramankutty, N., Brauman, K. A., Cassidy, E. S., Gerber, J. S., Johnston, M., . . . Zaks, D. P. M. (2011). Solutions for a cultivated planet. *Nature*, *478*(7369): 337–342. Retrieved from <https://doi.org/10.1038/nature10452>
- Fosu-Mensah, B. Y., Okoffo, E. D., Darko, G., & Gordon, C. (2016). Assessment of organochlorine pesticide residues in soils and drinking water sources from cocoa farms in Ghana. *SpringerPlus*, *5*(1): 869. Retrieved from <https://doi.org/10.1186/s40064-016-2352-9>
- Frelat, R., Lopez-Ridaura, S., Giller, K. E., Herrero, M., Douxchamps, S., Djurfeldt, A. A., Wijk, M. T., . . . van (2016). Drivers of household food availability in sub-Saharan Africa based on big data from small farms. *Proceedings of the National Academy of Sciences*, *113*(2): 458–463. Retrieved from <https://doi.org/10.1073/pnas.1518384112>
- Garnett, T., Appleby, M. C., Balmford, A., Bateman, I. J., Benton, T. G., Bloomer, P., . . . Godfray, H. C. J. (2013). Sustainable intensification in agriculture: Premises and policies. *Science*, *341*(6141): 33–34. Retrieved from <https://doi.org/10.1126/science.1234485>
- Gerba, C. P., & Smith, J. E. (2005). Sources of pathogenic microorganisms and their fate during land application of wastes. *Journal of Environmental Quality*, *34*(1): 42–48. Retrieved from <https://doi.org/10.2134/jeq2005.0042>
- Godfray, H. C. J., Beddington, J. R., Crute, I. R., Haddad, L., Lawrence, D., Muir, J. F., . . . Toulmin, C. (2010). Food security: The challenge of feeding 9 billion people. *Science*, *327*(5967): 812–818. Retrieved from <https://doi.org/10.1126/science.1185383>
- Goldsmith, P. (2017). The Faustian bargain of tropical grain production. *Tropical Conservation Science*, *10*.
- Guerrant, R. L., Oriá, R. B., Moore, S. R., Oriá, M. O., & Lima, A. A. (2008). Malnutrition as an enteric infectious disease with long-term effects on child development. *Nutrition Reviews*, *66*(9): 487–505. Retrieved from <https://doi.org/10.1111/j.1753-4887.2008.00082.x>

- Haas, C. N., Rose, J. B., & Gerba, C. P. (2014). *Quantitative microbial risk assessment*. Hoboken, NJ: John Wiley & Sons.
- Humphrey, J. H., Jones, A. D., Manges, A., Mangwadu, G., Maluccio, J. A., Mbuya, M. N. N., . . . Mujuru, H. A. (2015). The sanitation hygiene infant nutrition efficacy (SHINE) trial: Rationale, design, and methods. *Clinical Infectious Diseases*, *61*(suppl_7): S685–S702. Retrieved from <https://doi.org/10.1093/cid/civ844>
- Jepson, P. C., Guzy, M., Blaustein, K., Sow, M., Sarr, M., Mineau, P., & Kegley, S. (2014). Measuring pesticide ecological and health risks in West African agriculture to establish an enabling environment for sustainable intensification. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences*, *369*(1639): 20130491. Retrieved from <https://doi.org/10.1098/rstb.2013.0491>
- Jones, A. D., Shrinivas, A., & Bezner-Kerr, R. (2014). Farm production diversity is associated with greater household dietary diversity in Malawi: Findings from nationally representative data. *Food Policy*, *46*, 1–12. Retrieved from <https://doi.org/10.1016/j.foodpol.2014.02.001>
- Kassie, M., Teklewold, H., Jaleta, M., Marennya, P., & Erenstein, O. (2015). Understanding the adoption of a portfolio of sustainable intensification practices in eastern and southern Africa. *Land Use Policy*, *42*, 400–411. Retrieved from <https://doi.org/10.1016/j.landusepol.2014.08.016>
- Kobayashi, Y., Peters, G. M., Ashbolt, N. J., Shiels, S., & Khan, S. J. (2015). Assessing burden of disease as disability adjusted life years in life cycle assessment. *Science of the Total Environment*, *530–531*, 120–128. Retrieved from <https://doi.org/10.1016/j.scitotenv.2015.05.017>
- Kobayashi, Y., Peters, G. M., & Khan, S. J. (2015). Towards more holistic environmental impact assessment: Hybridisation of life cycle assessment and quantitative risk assessment. *Procedia CIRP*, *29*, 378–383. Retrieved from <https://doi.org/10.1016/j.procir.2015.01.064>
- Korpe, P. S., & Petri, W. A. Jr (2012). Environmental enteropathy: Critical implications of a poorly understood condition. *Trends in Molecular Medicine*, *18*(6): 328–336. Retrieved from <https://doi.org/10.1016/j.molmed.2012.04.007>
- Mapesa, J. O., Maxwell, A. L., & Ryan, E. P. (2016). An exposome perspective on environmental enteric dysfunction. *Environmental Health Perspectives*, *124*(8): 1121–1126. Retrieved from <https://doi.org/10.1289/ehp.1510459>
- Matson, P. A., Parton, W. J., Power, A. G., & Swift, M. J. (1997). Agricultural intensification and ecosystem properties. *Science*, *277*(5325): 504–509. Retrieved from <https://doi.org/10.1126/science.277.5325.504>
- Mendoza, G. A., & Martins, H. (2006). Multi-criteria decision analysis in natural resource management: A critical review of methods and new modelling paradigms. *Forest Ecology and Management*, *230*(1–3): 1–22. Retrieved from <https://doi.org/10.1016/j.foreco.2006.03.023>
- Michelson, H. (2017). Variable soils, variable fertilizer quality, variable prospects. *Tropical Conservation Science*, *10*.
- Mueller, N. D., Gerber, J. S., Johnston, M., Ray, D. K., Ramankutty, N., & Foley, J. A. (2012). Closing yield gaps through nutrient and water management. *Nature*, *490*(7419): 254–257. Retrieved from <https://doi.org/10.1038/nature11420>
- Neill, C., Jankowski, J., Brando, P. M., Coe, M. T., Deegan, L. A., Macedo, M. N., . . . Krusche, A. V. (2017). Surprisingly modest water quality impacts from expansion and intensification of large-scale commercial agriculture in the Brazilian Amazon-Cerrado region. *Tropical Conservation Science*, *10*.
- Palm, C., Neill, C., Lefebvre, P., & Tully, K. (2017). Targeting sustainable intensification of maize-based agriculture in East Africa. *Tropical Conservation Science*, *10*.
- Pellegrini, L., & Tasciotti, L. (2014). Crop diversification, dietary diversity and agricultural income: Empirical evidence from eight developing countries. *Canadian Journal of Development Studies/Revue Canadienne D'études Du Développement*, *35*(2): 211–227. Retrieved from <https://doi.org/10.1080/02255189.2014.898580>
- Rosegrant, M. W., & Cline, S. A. (2003). Global food security: Challenges and policies. *Science*, *302*(5652): 1917–1919. Retrieved from <https://doi.org/10.1126/science.1092958>
- Sadler, R., Maetam, B., Edokpolo, B., Connell, D., Yu, J., Stewart, D., . . . Laksono, B. (2016). Health risk assessment for exposure to nitrate in drinking water from village wells in Semarang, Indonesia. *Environmental Pollution*. Retrieved from <https://doi.org/10.1016/j.envpol.2016.06.041>
- Sanchez, P. A., Denning, G. L., & Nziguheba, G. (2009). The African Green Revolution moves forward. *Food Security*, *1*(1): 37–44. Retrieved from <https://doi.org/10.1007/s12571-009-0011-5>
- Sheahan, M., Barrett, C. B., & Goldvale, C. (2017). Human health and pesticide use in sub-Saharan Africa. *Agricultural Economics*. Retrieved from <http://barrett.dyson.cornell.edu/files/papers/Sheahan%20Barrett%20Goldvale%20-%20SSA%20pesticide%20and%20human%20health%20paper%20Mar%202017%20final.pdf>
- Sibhatu, K. T., Krishna, V. V., & Qaim, M. (2015). Production diversity and dietary diversity in smallholder farm households. *Proceedings of the National Academy of Sciences*, *112*(34): 10657–10662. Retrieved from <https://doi.org/10.1073/pnas.1510982112>
- Smith, V. H., Tilman, G. D., & Nekola, J. C. (1999). Eutrophication: Impacts of excess nutrient inputs on freshwater, marine, and terrestrial ecosystems. *Environmental Pollution*, *100*(1–3): 179–196. Retrieved from [https://doi.org/10.1016/S0269-7491\(99\)00091-3](https://doi.org/10.1016/S0269-7491(99)00091-3)
- Spera, S. (2017). Agricultural intensification can preserve the Brazilian Cerrado: Applying lessons from Mato Grosso and Goiás to Brazil's last agricultural frontier. *Tropical Conservation Science*, *10*.
- Steffen, W., Richardson, K., Rockström, J., Cornell, S. E., Fetzer, I., Bennett, E. M., . . . Sörlin, S. (2015). Planetary boundaries: Guiding human development on a changing planet. *Science*, *347*(6223): 1259855. Retrieved from <https://doi.org/10.1126/science.1259855>
- Swiss Federal Institute of Aquatic Science and Technology. (2016). *Chemical pollution in low- and middle-income countries*. Retrieved from <http://www.eawag.ch/en/departement/sandec/publications/chemical-pollution/>
- Tilman, D. (1999). Global environmental impacts of agricultural expansion: The need for sustainable and efficient practices. *Proceedings of the National Academy of Sciences*, *96*(11): 5995–6000. Retrieved from <https://doi.org/10.1073/pnas.96.11.5995>
- Tilman, D., Cassman, K. G., Matson, P. A., Naylor, R., & Polasky, S. (2002). Agricultural sustainability and intensive production practices. *Nature*, *418*(6898): 671–677. Retrieved from <https://doi.org/10.1038/nature01014>

- Trehan, I., Kelly, P., Shaikh, N., & Manary, M. J. (2016). New insights into environmental enteric dysfunction. *Archives of Disease in Childhood*, *101*(8): 741–744. Retrieved from <https://doi.org/10.1136/archdischild-2015-309534>
- Tscharntke, T., Clough, Y., Wanger, T. C., Jackson, L., Motzke, I., Perfecto, I., . . . Whitbread, A. (2012). Global food security, biodiversity conservation and the future of agricultural intensification. *Biological Conservation*, *151*(1): 53–59. Retrieved from <https://doi.org/10.1016/j.biocon.2012.01.068>
- Tyrrel, S. F., & Quinton, J. N. (2003). Overland flow transport of pathogens from agricultural land receiving faecal wastes. *Journal of Applied Microbiology*, *94*, 87–93. Retrieved from <https://doi.org/10.1046/j.1365-2672.94.s1.10.x>
- United Nations. (2015). *World population prospects: The 2015 revision*. United Nations, Department of Economic and Social Affairs, Population Division. Retrieved from <https://esa.un.org/unpd/wpp/>
- United States Agency for International Development. (2014). *Ghana: Nutrition profile*. Retrieved from <https://www.usaid.gov/sites/default/files/documents/1864/USAID-Ghana-Profile.pdf>
- van Ittersum, M. K., van Bussel, L. G. J., Wolf, J., Grassini, P., van Wart, J., Guilpart, N., . . . Cassman, K. G. (2016). Can sub-Saharan Africa feed itself? *Proceedings of the National Academy of Sciences*, *113*(52): 14964–14969. Retrieved from <https://doi.org/10.1073/pnas.1610359113>
- Waldron, A., Garrity, D., Malhi, Y., Girardin, C., Miller, D. C., & Seddon, N. (2017). Agroforestry can enhance food security while meeting other. *Sustainable Development Goals*, *10*.
- Wallington, K., & Cai, X. (2017). The food-energy-water nexus: A framework to address sustainable development in the tropics. *Tropical Conservation Science*, *10*.
- Ward, M. H., deKok, T. M., Levallois, P., Brender, J., Gulis, G., Nolan, B. T., & VanDerslice, J. (2005). Workgroup report: Drinking-water nitrate and health-recent findings and research needs. *Environmental Health Perspectives*, *113*(11): 1607–1614.
- West, P. C., Gerber, J. S., Engstrom, P. M., Mueller, N. D., Brauman, K. A., Carlson, K. M., . . . Siebert, S. (2014). Leverage points for improving global food security and the environment. *Science*, *345*(6194): 325–328. Retrieved from <https://doi.org/10.1126/science.1246067>
- World Health Organization. (2011). *Guidelines for drinking-water quality* (4th ed.). Geneva, Switzerland: Author. Retrieved from http://www.who.int/water_sanitation_health/publications/dwq-guidelines-4/en/