

Santonian Ammonites from the Blossom Sand in Northeast Texas

Authors: KENNEDY, W.J., LANDMAN, N.H., and COBBAN, W.A.

Source: American Museum Novitates, 2001(3332) : 1-9

Published By: American Museum of Natural History

URL: [https://doi.org/10.1206/0003-0082\(2001\)332<0001:SAFTBS>2.0.CO;2](https://doi.org/10.1206/0003-0082(2001)332<0001:SAFTBS>2.0.CO;2)

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

AMERICAN MUSEUM *Novitates*

PUBLISHED BY THE AMERICAN MUSEUM OF NATURAL HISTORY
CENTRAL PARK WEST AT 79TH STREET, NEW YORK, NY 10024
Number 3332, 9 pp., 5 figures April 26, 2001

Santonian Ammonites from the Blossom Sand in Northeast Texas

W.J. KENNEDY,¹ N.H. LANDMAN,² AND W.A. COBBAN³

ABSTRACT

The Blossom Sand is a lateral equivalent of the Austin Chalk Group, developed in Fannin, Lamar, and Red River counties in northeast Texas. Ammonites from this unit are *Placenticer*
syrtale (Morton, 1834), *Baculites* sp. cf. *haresi* Reeside, 1927, and *Glyptoxoceras texanum*, n.
sp. The general aspect of the *Placenticer* assemblage suggests a late Santonian age for this
unit.

INTRODUCTION

The term Blossom Sand was introduced by Gordon (1909) for what Veatch (1906) and others had termed the sub-Clarksville sand. The type locality is a well at Blossom, to the east of Paris, in Lamar County, Texas (fig. 1). The most comprehensive account of the unit remains that of Stephenson (1918). The unit reaches a thickness of up to 76 m (250 ft) in the eastern part of its exposure where it rests on rocks assigned to the Bonham Formation (Barnes, 1966, 1967). Traced westwards, the formation thins and loses its char-

acter, feathering out in Fannin County (Barnes, 1967).

The Blossom Sand was initially regarded as a lateral equivalent of part of the Cenomanian–Turonian Eagle Ford Group, but Stephenson (1918) demonstrated that it was equivalent to the Austin Chalk, and he suggested that the fauna indicated a correlation with the upper part of the Austin Chalk near Austin in Travis County, Texas. Subsequent authors (e.g., Young, 1963) regarded the unit as Santonian, but it remained imprecisely dated. Stephenson (1918) recorded *Hamites*, *Baculites asper* Morton, 1834, *Placenticer*

¹ Curator, Geological Collections, University Museum, Parks Road, Oxford, OX1 3PW, U.K.

² Curator, Division of Paleontology (Invertebrates), American Museum of Natural History.

³ Research Associate, Division of Paleontology (Invertebrates), American Museum of Natural History.




Fig. 1. The outcrop of the Blossom Sand in northeast Texas, showing the localities cited in the text.

sp., and *Prionotropis*?, but he did not illustrate these specimens, and their present whereabouts are unknown. In addition, most of his localities are no longer accessible. We describe below material from two new localities (fig. 1) that have been discovered since the work of Stephenson (1918): (1) USGS Mesozoic locality 18961: Blossom Sand, borrow pit just west of State Highway 37, 6.6 km (4 mi) north of Clarksville, Red River County, Texas. Collected by L. W. Stephenson, 1944. This collection yields abundant *Placenticerias syrtale* (Morton, 1834) as well as rare *Baculites* sp. cf. *harsi* Reeside, 1927. (2) USGS Mesozoic locality 18120: Top of Blossom Sand between Detroit and Bagwell, Red River County, Texas. Collected by D. J. Flesh, 1939. This collection consists of two specimens of *Glyptoxoceras texanum*, n. sp.

The key to dating the Blossom Sand is *Placenticerias syrtale*. Wolleben (1967) analyzed changes in a succession of *P. syrtale* assemblages through the "Lower Campanian" interval in trans-Pecos Texas. Since this work, the International Sub-Commission on Cretaceous Stratigraphy has reached a preliminary decision on the definition of the base of the Campanian (see below), and the *Placenticerias* sequence studied by Wolleben appears to span the Santonian–Campanian

boundary. The material from the Blossom Sand corresponds to the earliest subspecies of *P. syrtale* recognized by Wolleben, and it dates the unit as late Santonian.


BASE OF THE CAMPANIAN

The working party on the base of the Campanian Stage, meeting in Brussels in 1995 (Hancock and Gale, 1996), reached a preliminary decision that the base of the Campanian should be drawn at the extinction point of the widely distributed crinoid *Marsupites*. Gale et al. (1995) provided a detailed discussion of the relationship between this event and the various other datum-levels used to define this boundary, and they demonstrated that the extinction point of *Marsupites* in Texas occurs at the top of, or just within, the *Submortonicerias tequesquitense* Zone of Young (1963). It is in this sense that the base of the Campanian is applied in the present contribution.

Young (1963: 102) described the holotype (and only recorded specimen) of *Submortonicerias vandalianense* from "a Blossom lithology" that "may indicate a Blossom facies at a higher level than heretofore reported." This specimen appears to be from a different unit of a younger, Campanian age.

→

Fig. 2. *Placenticerias syrtale* (Morton, 1834). A–C. USNM 490961; D–F. USNM 490962; G–I. USNM 490963; J, K. USNM 490964; L, M. USNM 490965; N, O. USNM 490966. All specimens are from the Blossom Sand at USGS Mesozoic locality 18961 in Red River County, Texas. Arrows indicate the base of the body chamber. All figures are $\times 1$.



SYSTEMATIC PALEONTOLOGY

SUBORDER AMMONITINA HYATT, 1889

SUPERFAMILY HOPLITACEAE DOUVILLÉ, 1890

FAMILY PLACENTICERATIDAE HYATT, 1900

Genus *Placenticer* Meek, 1876

TYPE SPECIES: *Ammonites placenta* Dekay, 1828: 278, pl. 5, fig. 2 (not 5).

Placenticer *syrtale* (Morton, 1834)

Figures 2, 3I–P

Ammonites syrtalis Morton, 1834: 40, pl. 16, fig. 4. *Stantonoceras pseudocostatum* Johnson. Scott and Cobban, 1964: pl. 8.

Placenticer *syrtale* (Morton, 1834). Wolleben, 1967: 1161 (with additional synonymy).

Placenticer *syrtale syrtale* (Morton, 1834). Wolleben, 1967: 1161, pl. 150, fig. 5; pl. 151, figs. 1, 2, 5–7; text-fig. 7e, f.

Placenticer *syrtale adkinsi* Wolleben, 1967: 1164, pl. 151, figs. 8, 9; pl. 152, figs. 1, 2, 5–8; text-fig. 89.

Placenticer *syrtale rooneyi* Wolleben, 1967: 150, figs. 6, 7; pl. 151, figs. 3, 4; pl. 152, figs. 3, 4; text-fig. 7d, g.

Placenticer (*Stantonoceras*) *guadalupae* Roemer. Cobban, 1976: 125, pl. 2, fig. 7.

Placenticer *syrtale* (Morton, 1834). Kennedy and Cobban, 1991: 176, figs. 7.1–7.4, 11.5.

Placenticer *syrtale* (Morton, 1834). Kennedy and Cobban, 1993: 835, figs. 5.4–5.6.

Placenticer *syrtale* (Morton, 1834). Kennedy et al., 1995: pl. 2, figs. 1, 2.

TYPE: Holotype, by monotypy, is the original of Morton, 1834 (p. 40, pl. 16, fig. 4), from the “older Cretaceous” of Greene County, Alabama. It was refigured by Hyatt (1903: pl. 27, fig. 15; pl. 28, figs. 1, 2) and is no. 282 in the collection of the Academy of Natural Sciences of Philadelphia.


DESCRIPTION: There are 27 specimens, including USNM 490961–490969. Most specimens are juveniles, the majority of which are body chambers, with whorl heights ranging from 18 to 50 mm. Specimens display a broad range of variation. In more compressed

specimens, the ratio of whorl breadth to height is as low as 0.56; the coiling is very involute, with a tiny craterlike umbilicus. The umbilical wall and shoulder slope outward, with the greatest breadth just outside the umbilical shoulder. The inner flanks are feebly convex, the outer flanks are flattened and convergent, and the venter is very narrow and tabulate. Ornament consists of delicate falcoid ribs that form prominent crescentic ridges on the outer flank, and delicate ventral clavi that alternate across the venter, numbering around 20 per half whorl. As the whorl section becomes less compressed, the ornament coarsens. Low primary ribs strengthen across the umbilical wall and shoulder and form umbilicolateral bullae. These give rise to one or two low prorsiradial ribs that are straight on the inner flank and concave on the outer flank, where they are joined by intercalated ribs, all of which form crescentic bullae on the outer flank (fig. 2A–J). The ventral clavi are more numerous than the bullae. In the most robust specimens, the ratio of whorl breadth to height is as high as 0.8; the greatest breadth is at the umbilicolateral bullae. The flanks are convergent and the venter is broad and feebly convex. Coarse umbilicolateral bullae give rise to pairs of ribs, with prominent outer lateral nodes and ventral clavi that are two to three times as numerous as the nodes (figs. 2L–O, 3N–P). The largest specimen has an estimated diameter of 110 mm and appears to be a microconch body chamber. It shows eccentric coiling and a decline in ornament over the last 90% of the body chamber.

DISCUSSION: Kennedy and Cobban (1991: 175) discussed the temporal changes in the Gulf Coast *Placenticer* *syrtale* lineage: “At any one stratigraphic level there is great intraspecific variation, and many names have been introduced for different morphological variants. . . . Good illustrations of variation (and use of many names) are to be found in

→

Fig. 3. **A–H.** *Baculites* sp. cf. *harsi* Reeside, 1927. **A–D.** USNM 490972; **E–H.** USNM 490973; **I–P.** *Placenticer* *syrtale* (Morton, 1834); **I, M.** USNM 490967; **J–L.** USNM 490968; **N–P.** USNM 490969. All specimens are from the Blossom Sand at USGS Mesozoic locality 18961 in Red River County, Texas. Arrows indicate the base of the body chamber. All figures are $\times 1$.






Fig. 4. *Glyptoxoceras texanum*, n. sp. **A, B.** Paratype USNM 490971; **C–E.** holotype, USNM 490970. Both specimens are from the Blossom Sand at USGS Mesozoic locality 18120 in Red River County, Texas. Arrows indicate the base of the body chamber. All figures are $\times 1$.

Hyatt (1903), Reeside (1927) and Young (1963). Wolleben (1967) is the latest to have tackled this group, studying a collection of over 600 specimens from the San Carlos and Ojinaga areas of Presidio County in Trans-Pecos Texas and Chihuahua, Mexico, respectively. . . . He used the unidirectional trend of whorl height–umbilical node distance coefficients to divide what he termed the ‘polymorphic *Placentoceras* chronocline’ into chronologic polymorphic subspecies.” According to Wolleben *Placentoceras syrtale syrtale* (Morton, 1834) (upper Santonian to lower Campanian) is a subspecies that includes specimens corresponding to *P. syrtale*

of previous authors, *P. syrtale adkinsi* Wolleben, 1967 (lower Campanian) is a subspecies that includes specimens corresponding to *Placentoceras pseudosyrtale* of previous authors, and *P. syrtale rooneyi* Wolleben, 1967 (lower Campanian) is a subspecies that includes specimens of *Placentoceras newberryi* of previous authors. Although Wolleben favored this approach to “avoid future nomenclature confusion” (Wolleben, 1967: 1161), it overlooks the principles of priority set out in the *International Code of Zoological Nomenclature*.

In the Blossom Sand assemblage, specimens range from compressed and feebly or-

namented to less compressed and moderately strongly ornamented. They correspond to the earliest subspecies of Wolleben (1967). The incidence of weakly ornamented individuals is lower than that in the assemblage described by Kennedy and Cobban (1991) from the correlatives of the Santonian *Texanites shiloensis* Zone of Young (1963) in Mississippi, suggesting a higher level than this assemblage, but probably still within the Santonian as here defined.

OCCURRENCE: Blossom Sand at USGS Mesozoic locality 18961. The species ranges from upper Santonian to lower Campanian, and is known from northern Mexico, Texas, Alabama, Mississippi, New Mexico, Colorado, Wyoming, Montana, and Utah.

SUBORDER ANCYLOCERATINA WIEDMANN, 1966

SUPERFAMILY TURRILITACEAE GILL, 1871

FAMILY DIPLOMOCERATIDAE SPATH, 1926

SUBFAMILY DIPLOMOCERATINAE SPATH, 1926

Genus *Glyptoxoceras* Spath, 1925

TYPE SPECIES: *Hamites rugatus* Forbes, 1846:117, by the original designation of Spath, 1926:81.

Glyptoxoceras texanum, new species

Figure 4

DERIVATION OF NAME: From the state of Texas.

TYPE: Holotype is USNM 490970 (fig. 4C–E), paratype is USNM 490971 (fig. 4A, B), from the top of the Blossom Sand at USGS Mesozoic locality 18120, between Detroit and Bagwell, Red River County, Texas.

DIAGNOSIS: A large species of *Glyptoxoceras*; body chamber bears rursiradial ribs and periodic constrictions, each of which is flanked by a flared adapertural rib.

DESCRIPTION: The holotype USNM 490970 is a body chamber fragment 120 mm long, with replaced shell (fig. 4C–E). The maximum preserved whorl height is 31 mm. The specimen has suffered some post-mortem crushing, but the whorl section of the uncrushed portion is compressed ovoid with the venter more narrowly rounded than the dorsum; the ratio of whorl breadth to height is 0.75. For most of the length of the fragment,

the ornament consists of coarse, narrow ribs; the rib index is 6. The ribs are much narrower than are the interspaces; ribs are weak and transverse to feebly concave across the dorsum, stronger and convex across the dorsolateral margin, strong, straight, and rursiradial on the flanks, and transverse on the venter, where they attain their maximum height. There are two broad, deep constrictions on the fragment, each of which is flanked by a flared adapertural rib. The adoral of these two constrictions is succeeded by a section of shell 40 mm in length on which the ornament changes markedly, indicating that the specimen is part of the terminal adult body chamber immediately prior to the final aperture. Ribbing is much finer, lower, and weaker, and the rib index increases to 11. There is a weak constriction and flared rib immediately adapical of the aperture.

The paratype USNM 490971 is much smaller, 52 mm long, crushed, with a maximum preserved whorl height of 11 mm and whorl breadth of 13.5 mm (fig. 4A, B). The depressed whorl section is probably an artifact of post-mortem deformation. Ornament consists of narrow, distant ribs that are weak and feebly concave across the dorsum; they strengthen across the dorsolateral margin, where they are sharp and high and much narrower than the interspaces. The ribs are markedly prorsiradial on the flanks and strong and feebly convex across the venter. The sutures are not exposed.

DISCUSSION: The holotype of *Glyptoxoceras ellisoni* Young, 1963 (p. 46, pl. 1, figs. 10–14, 16–20; pl. 73, fig. 9; pl. 78, fig. 6) from the Dessau Chalk in central Texas is a body chamber fragment 125 mm long with a maximum preserved whorl height of 22 mm (fig. 5). It overlaps in size with the holotype of *G. texanum* but it has much coarser, blunter ribbing (even when the differences of preservation are taken into account), with a rib index of around 4; there are no constrictions. *Glyptoxoceras rubeyi* (Reeside, 1927) (p. 14, pl. 3, figs. 8–10; pl. 5, figs. 3–11) has much more widely separated and coarser ribs. The holotype of *Glyptoxoceras novimexicanum* (Reeside, 1927) (p. 8, pl. 4, figs. 1–4), a body chamber fragment that is comparable in size to the holotype of *G. texanum*, lacks constrictions.




Fig. 5. *Glyptoxoceras ellisoni* Young, 1963. Holotype, Texas Memorial Museum Collections no. 182, from the Austin Chalk on Brushy Creek, 800 m (0.5 mi) south of Hutto, Williamson County, Texas. All figures are $\times 1$.

OCCURRENCE: As for type.

FAMILY BACULITIDAE GILL, 1871

Genus *Baculites* Lamarck, 1799

TYPE SPECIES: *Baculites vertebralis* Lamarck, 1801:103, by subsequent designation by Meek, 1876:391.

Baculites sp. cf. *haresi* Reeside, 1927

Figure 3A–H

cf. *Baculites ovatus* Say, 1820 var. *haresi* Reeside, 1927:10, pl. 6, figs. 5–10; pl. 7, figs. 9, 10.

DESCRIPTION: There are two body chamber fragments, USNM 490972 (fig. 3A–D) and USNM 490973 (fig. 3E–H). The latter has a maximum preserved whorl height of 20 mm and a ratio of whorl breadth to height of 0.74; the whorl section is ovoid and the dorsum is more broadly rounded than the venter. Both fragments are virtually smooth, with traces of delicate riblets and striae that are strongly prorsiradiate on the outer flank and cross the venter in a broad convexity (fig. 3H).

DISCUSSION: These fragments belong to a very generalized *Baculites*, and are most comparable to *Baculites haresi* Reeside, 1927 (e.g., pl. 6, figs. 5, 6).

OCCURRENCE: Blossom Sand at USGS Mesozoic locality 18961.

ACKNOWLEDGMENTS

We thank J. M. Hancock (Shaftesbury, U.K.), H.C. Klinger (Cape Town, South Africa), K. Tanabe (Tokyo, Japan), and R. H. Mapes (Athens, Ohio) for critical reviews of an earlier version of this manuscript. W.J.K. acknowledges the financial support of the Natural Environment Research Council (U.K.) and the technical support of the staff of the Geological Collections, University Museum, Oxford, and Department of Earth Sciences, Oxford. N.H.L. thanks B. Worcester and S. Crooms (both AMNH) for help in preparation of the manuscript. The specimens were provided by the U.S. Geological Survey.

REFERENCES

- Barnes, V. E.
 1966. Geologic atlas of Texas, Texarkana sheet. Univ. Texas Bureau of Economic Geology, scale 1:250,000.
 1967. Geologic atlas of Texas, Sherman sheet. Univ. Texas Bureau of Economic Geology, scale 1:250,000.
- Cobban, W. A.
 1976. Ammonite record from the Mancos Shale of the Castle Valley—Woodside area, east-central Utah. Brigham Young Univ. Geol. Stud. 22: 117–126.
- Dekay, J. E.
 1828. Report on several multilocular shells from the State of Delaware: with observations on a second specimen of the

- new genus *Eurypterus*. *Lyceum Nat. Hist.* 2: 273–279.
- Douvillé, H.
1890. Sur la classification des Cératites de la Craie. *Bull. Soc. Géol. France* (3), 18: 275–292.
- Forbes, E.
1846. Report on the fossil Invertebrata from southern India, collected by Mr. Kaye and Mr. Cunliffe. *Trans. Geol. Soc. London* (2), 7: 97–174.
- Gale, A. S., P. Montgomery, W. J. Kennedy, J. M. Hancock, J. A. Burnett, and J. M. McArthur
1995. Definition and global correlation of the Santonian–Campanian boundary. *Terra Nova* 7 (for 1995): 611–622.
- Gill, T.
1871. Arrangement of the families of Mollusks. *Smithson. Misc. Collect.* 227: xvi + 49 pp.
- Gordon, C. H.
1909. The chalk formations of northeast Texas. *Am. J. Sci.* (4), 27: 369–373.
- Hancock, J. M., and A. S. Gale
1996. The Campanian Stage. *Bull. Inst. R. Sci. Nat. Belgique, Sci. Terre* 66 (suppl.): 103–109.
- Hyatt, A.
1889. Genesis of the Arietidae. *Smithson. Contrib. Knowl.* 673: xi + 239 pp.
1900. Cephalopoda. In K. A. Von Zittel (ed.), *Textbook of palaeontology*: 502–604. *Transl. Eastman. London: Macmillan*, 1896–1900.
1903. Pseudoceratites of the Cretaceous. *U.S. Geol. Surv. Monogr.* 44: 351 pp.
- Kennedy, W. J., and W. A. Cobban
1991. Upper Cretaceous (upper Santonian) *Boehmoceras* fauna from the Gulf Coast region of the United States. *Geol. Mag.* 128: 167–189.
1993. Lower Campanian (Upper Cretaceous) ammonites from the Merchantville Formation of New Jersey, Maryland, and Delaware. *J. Paleontol.* 67: 828–849.
- Kennedy, W. J., R. A. Johnson, and W. A. Cobban
1995. Upper Cretaceous ammonite faunas of New Jersey. In J.E.B. Baker (ed.), *Contributions to the paleontology of New Jersey*: 24–55. *Geol. Assoc. New Jersey*.
- Lamarck, J. P. B. A. de M. de
1799. *Prodrome d'une nouvelle classification des coquilles*. *Mém. Soc. Hist. Nat. Paris*. (1799): 63–90.
1801. *Système des animaux sans vertèbres*. Paris: The author, vii + 432 pp.
- Meek, F. B.
1876. A report on the invertebrate Cretaceous and Tertiary fossils of the upper Missouri country. *U.S. Geol. Surv. Territ. Rep.* (Hayden) 9: lxiv + 629 pp.
- Morton, S. G.
1834. Synopsis of the organic remains of the Cretaceous group of the United States. Illustrated by nineteen plates, to which is added an appendix containing a tabular view of the Tertiary fossils discovered in America. Philadelphia: Key and Biddle, 88 pp.
- Reeside, J. B., Jr.
1927. The cephalopods of the Eagle Sandstone and related formations in the western interior of the United States. *U.S. Geol. Surv. Prof. Pap.* 151: 40 pp.
- Say, T.
1820. Fossil zoology. *Am. J. Sci.* 1st Ser. 2(4): 34–45.
- Scott, G. R., and W. A. Cobban
1964. Stratigraphy of the Niobrara Formation at Pueblo, Colorado. *U.S. Geol. Surv. Prof. Pap.* 454-L: L1–L27.
- Spath, L. F.
1925. On Senonian Ammonoidea from Jamaica. *Geol. Mag.* 62: 28–32.
1926. On new ammonites from the English Chalk. *Geol. Mag.* 63: 77–83.
- Stephenson, L. W.
1918. A contribution to the geology of northeastern Texas and southern Oklahoma. *U.S. Geol. Surv. Prof. Pap.* 120-H: 129–163.
- Veatch, A. C.
1906. Geology and underground water resources of northern Louisiana and southern Arkansas. *U.S. Geol. Surv. Prof. Pap.* 46: 422 pp.
- Wiedmann, J.
1966. Stammesgeschichte und System der posttriadischen Ammonoideen; ein Überblick. *Neues Jahrb. Geol. Paleontol. Abh.* 125: 49–79; 127: 13–81.
- Wolleben, J. A.
1967. Senonian (Cretaceous) Mollusca from Trans-Pecos Texas and northeastern Chihuahua, Mexico. *J. Paleontol.* 41: 1150–1165.
- Young, K.
1963. Upper Cretaceous ammonites from the Gulf Coast of the United States. *Univ. Texas Bull.* 6304: ix + 373 pp.

Recent issues of the *Novitates* may be purchased from the Museum. Lists of back issues of the *Novitates* and *Bulletin* published during the last five years are available at World Wide Web site <http://nimidi.amnh.org>. Or address mail orders to: American Museum of Natural History Library, Central Park West at 79th St., New York, NY 10024. TEL: (212) 769-5545. FAX: (212) 769-5009. E-MAIL: scipubs@amnh.org

∞ This paper meets the requirements of ANSI/NISO Z39.48-1992 (Permanence of Paper).