



Typification of *Arisarum Vulgare* O. Targ. Tozz. (Araceae)

Author: Ferrer-Gallego, Pablo

Source: *Candollea*, 69(2) : 115-118

Published By: The Conservatory and Botanical Garden of the City of Geneva (CJBG)

URL: <https://doi.org/10.15553/c2014v692a2>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Typification of *Arisarum vulgare* O. Targ. Tozz. (Araceae)

Pablo Ferrer-Gallego

Abstract

FERRER-GALLEGO, P. (2014). Typification of *Arisarum vulgare* O. Targ. Toz (Araceae). *Candollea* 69: 115-118. In English, English abstract.

The typification of the name *Arisarum vulgare* O. Targ. Tozz. (= *Arum arisarum* L.) (Araceae) is discussed. The designation of the corresponding type is based on the consultation of the Linnaeus's original material and the literature cited in the respective protologue. An illustration from Clusius is designated as the lectotype. An epitype is also designated here in order to avoid any ambiguity in the interpretation of the lectotype specimen.

Key-words

ARACEAE – *Arisarum* – Typification

Address of the author: CIEF, Centro para la Investigación y la Experimentación Forestal, Servicio de Vida Silvestre, Generalitat Valenciana, Avda. Comarques del País Valencià, 114, 46930, Quart de Poblet, Valencia, Spain. Email: flora.cief@gva.es

Submitted on January 6, 2014. Accepted on July 24, 2014.

Edited by P. Bungener

ISSN: 0373-2967 – Online ISSN: 2235-3658 – *Candollea* 69(2): 115-118 (2014)

© CONSERVATOIRE ET JARDIN BOTANIQUES DE GENÈVE 2014

Introduction

The genus *Arisarum* Mill. (Araceae) with three species (*A. proboscideum* (L.) Savi, *A. simorrhinum* Durieu and *A. vulgare* O. Targ. Tozz.) is distributed in southwestern Europe, Mediterranean and Macaronesian area (MAYO & al., 1997; GALÁN & CASTROVIEJO, 2007).

The lectotype of *A. simorrhinum* was designated by GALÁN (2007) from a specimen in P, and the lectotype of *A. proboscideum* was designated by GALÁN & CASTROVIEJO (2004) from an illustration of Barrelier. The purpose of this paper is to contribute to the stability of the nomenclature by lectotypification of the name *Arisarum vulgare* (= *Arum arisarum* L.). The designation of the corresponding type is based on the consultation of Linnaeus's original material and the literature cited in the respective protologue.

Arisarum vulgare is an herbaceous perennial plant 20-30 (-40) cm; the tuber is irregular ovoid; ovate to arrow-shaped sagittate leaves, with a petiole 5.5-37.5(-51) × 0.1-1.1 cm; the inflorescence is an erect spadix, with a peduncule 9-36(-43.5) × 0.1-0.5 cm, spathe with purplish-brown or olive green striped tube, 2.9-7.5 × 0.6-4 cm, cucullate, tubular, at the top the spadix is club-shaped and curved. Inhabits in damp, fertile sheltered sites in maquis, garrigues and rocky steppes (BOYCE, 1990; MAYO & al., 1997; GALÁN & CASTROVIEJO, 2007). This species hybridizes with *A. simorrhinum*.

Lectotypification of *A. vulgare*

The protologue of *Arum arisarum* L. (LINNAEUS, 1753: 966) consists of a short morphological diagnosis, taken from LINNAEUS (1738: 435) and VAN ROYEN (1740: 7), with three synonyms: two are cited from BAUHIN (1623: 196): “*Arisarum latifolium majus*”, “*Arisarum latifolium alterum*”, and one referred to CLUSIUS (1601: LXXIII, Fig. 1) and LOBELIUS (1581: 598) “*Arisarum latifolium*”, as well as locality indication “Habitat in Mauritaniae, Italiae, Lusitaniae, Hispaniae, Galloprovinciale nemoribus”. The latter two authors provided the illustrations that can be considered original material. In LINN, the specimen no. 1079.10 (image available at <http://linnean-online.org/11408/>) includes the annotation “15 *Arisarum*”, explicitly referring to the number of the species account of *Arum arisarum* L. in LINNAEUS (1753: 966). The sheet bears four specimens certainly to be referred to a single gathering.

BOYCE (1990: 20) cited Brander's specimen in LINN (no. 1079.10) as “holotype”, a typification accepted by GALÁN (2007: 62), but not indicated by GALÁN & CASTROVIEJO (2007: 305). However, JARVIS (2007: 318) indicates that this designation is incorrect, because this material was not received by Linnaeus until after 1753. This herbarium sheet (no. 1079.10) has been written by Linnaeus “*Algir*”. The manuscript list “*Algerica Branderi*”, was prepared around 1756 by Daniel Solander from Brander's specimens (JARVIS, 2007: 195).




Fig. 1. – Lectotype of *Arum arisarum* L. (CLUSIUS, 1601: tab. LXXIII).

At UPS-BURSER, an herbarium sheet exists with original material (sheet X: 143) (Fig. 2). The sheet bears two leaves and one label: “*Arisarum latifolium alterum* Bauh. / [illegible] in Hort. / 143”. Linnaeus's citing the polynomial from BAUHIN (1623) provides a link to the specimen in UPS-BURSER (Herb. Burser X: 143). Joachim Burser's herbarium was arranged and labelled according to BAUHIN (1623) (STEARN, 1957; JARVIS, 2007) and was cited in the introduction to LINNAEUS (1753) as one of the sources for that work. This herbarium was in Uppsala when it was used by Linnaeus for the interpretation of the names that appear in Bauhin's work. Therefore, this specimen is undoubtedly original material. However, unfortunately this specimen is very incomplete, and contains only two very different leaves (Fig. 2), and this could have led Boyce to designate the sheet in LINN (no. 1079.10) as the holotype. We have been unable to trace any further original material in any of the other Linnaean and Linnaean-linked herbaria (see also JARVIS, 2007: 318).



Fig. 2. – Original material of *Arum arisarum* L. [s.coll., UPS-BURSER] [© Herbarium UPS. Reproduced with permission]

Therefore, as the herbarium sheet in UPS-BURSER (Herb. Burser X: 143) is very incomplete, we select here as the lectotype the element of the protologue which best matches the traditional and current concept of *Arum arisarum*. Among the elements cited in the protologue, the illustrations by CLUSIUS (1601) and LOBELIUS (1581) clearly shows the characters of the leaves, spathe and spadix as indicated by LINNAEUS (1753), “*acaule, foliis cordato-oblongis, spatha bifida, spadice incurvo*”, and they are therefore suitable for typification of *A. arisarum*. These two illustrations are identical, therefore we designated here the Clusius’s illustration as the lectotype (Fig. 1).

However, these images do not show an important diagnostic character for distinguishing this species from *A. simorrhinum*, that is the greater length of the spathe peduncule in respect to the length of the leaf petioles. In addition, an epitype is also designated here in order to avoid any ambiguity in the interpretation of the lectotype specimen.

Arum arisarum L., Sp. Pl.: 966. 1753.

≡ *Arisarum vulgare* O. Targ. Tozz. in Ann. Mus. Imp. Fis. Firenze 2: 67. 1810.

Lectotypus (here designated): “*Arisarum latifolium* I”. (CLUSIUS, 1601: tab. LXXIII) (Fig. 1).

Epitypus (here designated): **SPAIN. Prov. Valencia:** Xàtiva, Castillo de Xàtiva, 30SYJ1518, 140 m, 2.III.2014, fl. & fr., P. Ferrer & I. Ferrando 014001 (VAL [VAL 220580!], arrow) (Fig. 3).



Fig. 3. – Epitype of *Arum arisarum* L. (designated by an arrow). [P. Ferrer & I. Ferrando 014001, VAL] [© Herbarium VAL. Reproduced with permission]

Acknowledgements

We thank Dr Mats Hjertson (Museum of Evolution, Botany Section Uppsala University, Sweden) for his help.

References

- BAUHIN, C. (1623). *Pinax theatri botanici*. Basel, Switzerland.
- BOYCE, P. (1990). Friar’s cowls and mouse plants: the genus *Arisarum*. *Bot. Mag. (Kew Mag.)* 7: 14-21.
- CLUSIUS, C. (1601). *Rariorum plantarum Historia, et Appendix ad Historiam plantarum*. Vol. 2. Antwerpen.
- GALÁN, A. (2007). Typification of *Arisarum simorrhinum* Durieu (Araceae). *Candollea* 62: 61-64.
- GALÁN, A. & S. CASTROVIEJO (2004). Typification of two Linnaean names in Araceae: *Arum proboscideum* L. and *Arum muscivorum* L. f. *Taxon* 53: 1047-1048.
- GALÁN, A. & S. CASTROVIEJO (2007). *Arisarum* Mill. In: CASTROVIEJO, S. & al. (ed.), *Fl. Iber.* 19: 304-308. Real Jardín Botánico, CSIC, Madrid.

- JARVIS, C. E. (2007). *Order out of Chaos. Linnaean plant names and their types*. Linnean Society of London & Natural History Museum, London.
- LINNAEUS, C. (1738). *Hortus Cliffortianus*. Amsterdam.
- LINNAEUS, C. (1753). *Species Plantarum*. Stockholm.
- LOBELIUS, M. (1581). *Plantarum seu Stirpium Icones*. Antwerpen.
- MAYO, S. J., J. BOGNER & P. C. BOYCE (1997). *The genera of Araceae*. Royal Botanic Kew, Kew.
- STEARNS, W. T. (1957). *An introduction to the Species Plantarum and cognate botanical works of Carl Linnaeus*. In: LINNAEUS, C., *Species Plantarum, a facsimile of the first edition, 1753*. Vol. 1. London.
- VAN ROYEN, A. (1740). *Florae Leydensis prodromus, exhibens plantas quae in horto academico Lugduno-Batavo aluntur*. Leiden.