



First record of *Dentella repens* (Rubiaceae) from Madagascar

Authors: & , Sylvain G. Razafimandimbison, and Manjato, Nadiah

Source: *Candollea*, 74(1) : 43-45

Published By: The Conservatory and Botanical Garden of the City of Geneva (CJBG)

URL: <https://doi.org/10.15553/c2019v741a6>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

First record of *Dentella repens* (Rubiaceae) from Madagascar

Sylvain G. Razafimandimbison & Nadiah Manjato

Abstract

RAZAFIMANDIMBISON, S.G. & N. MANJATO (2019). First record of *Dentella repens* (Rubiaceae) from Madagascar. *Candollea* 74: 43–45. In English, English and French abstracts. DOI: <http://dx.doi.org/10.15553/c2019v741a6>

A first record of the genus *Dentella* L. (*Rubiaceae*) from Madagascar is reported here. *Dentella repens* (L.) J.R. Forst. & G. Forst. was collected for the first time in the northwestern part of the island, within Analalava District of the Sofia Region. The material collected possesses all the salient characters of the species (e.g., creeping, prostrate herb, toothed corolla lobes, indehiscent fruits covered by transparent, flattened, bulbous-tipped trichomes). We postulate that the species has most likely recently reached Madagascar via long-distance dispersal either from Sri Lanka, where it is native, or Mauritius, where it has become naturalized.

Résumé

RAZAFIMANDIMBISON, S.G. & N. MANJATO (2019). Première mention de *Dentella repens* (Rubiaceae) à Madagascar. *Candollea* 74: 43–45. En anglais, résumés anglais et français. DOI: <http://dx.doi.org/10.15553/c2019v741a6>

Le genre *Dentella* L. (*Rubiaceae*) est signalé pour la première fois à Madagascar. *Dentella repens* (L.) J.R. Forst. & G. Forst. a été récoltée au nord-ouest de l'île, dans le District d'Analalava de la Région Sofia. Le matériel collecté possède tous les caractères essentiels de l'espèce (port herbacé, rampant et prostré, lobes de la corolle dentés, fruits indéhiscent couverts de trichomes transparents et aplatis à extrémité bulbeuse). Nous supposons que l'espèce a récemment atteint Madagascar par dispersion à longue distance, soit à partir du Sri Lanka où elle est indigène, soit de l'île Maurice où elle s'est naturalisée.

Keywords

RUBIACEAE – *Dentella* – Madagascar – Biodiversity

Addresses of the authors:

SGR: Swedish Museum of National History, Department of Botany, Box 50007, SE-10405 Stockholm, Sweden. E-Mail: sylvain.razafimandimbison@nrm.se

NM: Missouri Botanical Garden, B.P. 3319, Antananarivo 101, Madagascar.

First published online on March 28, 2019.

Introduction

Our knowledge of the Malagasy *Rubiaceae* has greatly been improved, as a result of extensive field collecting in Madagascar coupled with the phylogenetic and taxonomic studies during the last 20 years or so (e.g., ROBBRECHT & MANEN, 2006; RAZAFIMANDIMBISON et al., 2008; BREMER & ERIKSSON, 2009). In Madagascar, the coffee family (*Rubiaceae*) is the second largest flowering plant family and the largest woody plant family, with no less than 900 species (DAVIS & BRIDSON, 2003), including introduced species. Members of this family are important components of various habitats in the country, ranging from evergreen rainforest, sclerophyllous forest, deciduous dry forest and thicket forest, to savanna woodland (DAVIS & BRIDSON, 2003). Almost all species of *Rubiaceae* are endemic to Madagascar, with the exceptions of a few species, e.g., *Guettarda speciosa* L. (*Guettardeae*), *Morinda citrifolia* L. (*Morindeae*), *Mussaenda arcuata* Poir. (*Mussaendeae*), *Oldenlandia capensis* L. f., *O. herbacea* (L.) Roxb., *Richardia brasiliensis* Gomes (*Spermacoceae*), that have paleotropical or pantropical distributions. Many herbaceous *Rubiaceae* (e.g., *Oldenlandia* spp., *Spermacoce* spp., all *Spermacoceae*) commonly occur on roadsides and in grasslands or cultivated areas (DAVIS & BRIDSON, 2003). However, it is not known whether these are introduced species that have been naturalized.

We report here the first record of the rubiaceous genus *Dentella* L. from Madagascar (Fig. 1). *Dentella* is a small genus of eight species of creeping herbs, with six of its species endemic to Australia, one, *D. concinna* Airy Shaw, to Myanmar and one species, *D. repens* (L.) J.R. Forst. & G. Forst., extending from Southeast Asia to subtropical Asia and the southwestern Pacific. The name originated from the Greek word “dentos” (= tooth), referring to its distinct toothed corolla lobes (Fig. 1). The genus is currently classified in the predominantly herbaceous tribe *Spermacoceae* of the subfamily *Rubioideae*, and has been shown to be sister to the African genus *Pentodon* Hochst. (e.g., KÅREHED et al., 2007). The *Dentella*-*Pentodon* lineage is distinct by its five-merous flowers. *Dentella* differs from *Pentodon* by its creeping habit, stipules with short sheaths and numerous narrow fimbriae, lax inflorescences, peltate placentas with a bilobed apex and its dry, indehiscent fruits covered by white, glassy hairs (as opposed to erect herbs, entire stipules, solitary flowers, semi-globose placentas along the septum and glabrous capsules in *Pentodon*).

Dentella repens occurs throughout tropical Asia to Australasia, from Sri Lanka through India, Bangladesh, Myanmar and Nepal to southern China and south through Indochina, Peninsular Malaysia, the Philippines and Indonesia to Australia, Micronesia and Polynesia (e.g., KÅREHED & BREMER, 2007). The species is considered to be an introduced weed on Mauritius (VERDCOURT, 1983a: 44). Likewise, it has been reported to have been introduced to Mexico and the USA. We report here the first record of *D. repens* from Madagascar. The

second author and a co-worker collected a herbaceous plant that has the salient features of this species (Fig. 1). Below, we provide a description of the new collection of *D. repens* based on the fertile material in flower and fruit that was collected in a temporarily inundated area near the Mahavelona Masi-loka Swamp in the Antsanifera Fok. of the Antonibe Com., Analalava Dist., Sofia Region (northwestern Madagascar). The swamp was dominated by *Nymphaea nouchali* Burm. f. (*Nymphaeaceae*) et *Typha domingensis* Pers. (*Typhaceae*).

Description

Dentella repens (L.) J.R. Forst. & G. Forst., Char. Gen. Pl.: 25, tab: 13. 1775.

= *Oldenlandia repens* L., Mant. Pl. 1: 40. 1767.

Lectotypus (designated by VERDCOURT, 1983b): **INDIA** [?]: *sine loco*, s.d., *Anon. s.n.* (LINN n° 155/2).

Small prostrate herb, rooting at nodes. *Stems* and *leaves* fleshy, leaf blades sessile, 4–5 × 1.5–2 mm, obovate to spatulate, base narrowed, apex acute; stipules connate. *Flowers* axillary, solitary, sessile, 5-merous. Calyx tube c. 1 mm long; lobes 5, subulate. Corolla tube funnel-shaped, green-yellowish, glabrous outside and inside; lobes white, toothed. Stamens 5, included, attached in the middle of corolla tube. Ovary 2-celled; style filiform; stigma 2-lobed. *Fruits* globose, indehiscent, densely villous with multicellular, transparent, bulbous-tipped trichomes, becoming glabrescent when fully mature, many-seeded with persistent calyx crown. *Seeds* minute, angular, brown, rugose.

Specimen examined. – MADAGASCAR. Reg. Sofia [Prov. Mahajanga]: Dist. Analalava, Cne. Antonibe, Fkt. Antsanifera, marais de Mahavelona Masiloka, [14°45'47"S 47°25'57"E], 20 m, 12.IX.2016, fl. & fr., Manjato & Avizara 777 (MO, TAN).

Note. – We postulate that *D. repens* may well have reached Madagascar via a long-distance dispersal either from Sri Lanka or Mauritius. On the other hand, it is possible that the species might be present in East Africa and remains unnoticed. In this case, a long-distance dispersal from East Africa to Madagascar cannot be ruled out.

Acknowledgments

We thank Evah Haririsoa, Avizara Clovis and the VERAMA team, in particular Alain Andrianandraina and Mr Roger Razakamampianina, for their help in the field.



Fig. 1. – *Dentella repens* (L.) J.R. Forst. & G. Forst. showing its large corollas with toothed lobes and indehiscent fruits covered by transparent, flattened and bulbous-tipped trichomes.

References

- BREMER, B. & T. ERIKSSON (2009). Time tree of Rubiaceae: phylogeny and dating the family, subfamily, and tribes. *Int. J. Plant Sci.* 170: 766–793.
- DAVIS, A.P. & D. BRIDSON (2003). Introduction to the Rubiaceae: Princess Flowers. In: GOODMAN, S.M. & J.P. BEAMSTEAD (ed.), *The Natural History of Madagascar*: 431–432. University of Chicago Press.
- KÅREHED, J. & B. BREMER (2007). The systematics of Knoxiaceae (Rubiaceae) – molecular data and their taxonomic consequences. *Taxon* 56: 1051–1076.
- RAZAFIMANDIMBISON, S.G., C. RYDIN & B. BREMER (2008). Evolution and trends in the Psychotriaceae alliance (Rubiaceae) – A rarely reported evolutionary change from many-seeded carpels from one-seeded carpels. *Molec. Phylogenet. Evol.* 48: 207–159.
- ROBBRECHT, E. & J.-F. MANEN (2006). The major evolutionary lineages of the coffee family (Rubiaceae, Angiosperms). Combined analysis (nDNA and cpDNA) to infer the position *Coptosapelta* and *Luculia*, and supertree construction based on *rbcL*, *rps16*, *trnL-trnF*, and *atpB-rbcL* data. A new classification in two subfamilies, Cinchonoideae and Rubioideae. *Syst. Geogr. Pl.* 76: 85–146.
- VERDCOURT, B. (1983a). Rubiacées. *Fl. Mascareignes* 108: 43–44.
- VERDCOURT, B. (1983b). Notes on Mascarene Rubiaceae. *Kew Bull.* 34: 521–574.