



## **A Comparison of Hemocytes and Their Phenoloxidase Activity among Botryllid Ascidians**

Authors: Shirae, Maki, and Saito, Yasunori

Source: Zoological Science, 17(7) : 881-891

Published By: Zoological Society of Japan

URL: <https://doi.org/10.2108/zsj.17.881>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

# A Comparison of Hemocytes and Their Phenoloxidase Activity among Botryllid Ascidians

Maki Shirae\* and Yasunori Saito

Shimoda Marine Research Center, University of Tsukuba,  
Shimoda 5-10-1, Shizuoka 415-0025, Japan

---

**ABSTRACT**—The colony specificity of colonial animals involves allorecognition reactions, which are the defense reactions for allogeneic tissues that occur naturally. In colonial botryllid ascidians, all of the species already studied have colony specificity, and their allorecognition modes differ from one another. However, in most of these botryllids, morula cells (MCs) always participate in the allorecognition reactions, and the prophenoloxidase (proPO) system of MCs is considered to contribute to the allorecognition reaction. The present study was performed using five botryllids and *Sympylegma reptans*, which is closely related to botryllids, in an effort to clarify the relationship between the modes of allorecognition and the characteristics of MCs, such as the ratio of MCs to total hemocytes and the phenoloxidase (PO) activity levels in the MCs. The MCs of these six ascidians resembled one another morphologically and the MCs of all species showed PO activity. In *Botryllus scalaris*, PO activity was also found in granular leukocytes, but the level of activity was much lower than that in MCs. The PO of these species resembles one another, at least in terms of their sensitivity to inhibition by common inhibitors of the proPO system. The PO activity per fixed number of hemocytes varied among these ascidians. This variation was due to a difference in the ratio of MCs to total hemocytes and/or a difference in PO activity per MC. In most ascidians, except for *B. scalaris*, the rejection reaction area showed a higher level of PO activity than the fusion area of the syngeneic colonies. These results suggest that the characteristics of MCs including their PO activity are closely correlated with the mode of the allorecognition reaction.

---

## INTRODUCTION

Colony specificity, which is a phenomenon observed in colonial animals from sponges to ascidians, is regarded as a type of allogeneic recognition in multicellular animals. When colonies touch at their growing edges, compatible colonies fuse to become a single mass, but incompatible colonies do not fuse and usually reject each other. In botryllids, a group of colonial ascidians, all of the species already examined show colony specificity (Taneda *et al.*, 1985; Rinkevich, 1992; Saito *et al.*, 1994). In botryllids, the fusion reaction occurs in the same way, but their allorecognition modes differ even though they belong to the same family (Taneda *et al.*, 1985; Rinkevich, 1992; Saito *et al.*, 1994). Therefore, allorecognition reactions in botryllids are very useful objects of study to elucidate the organization and diversity of allorecognition systems in colonial animals.

In botryllids, allorecognition reactions are initiated at different stages on the way to fusion, and the variation in allorecognition reactions has been interpreted as reflecting a

difference in nonself-recognition sites in their allogeneic recognition systems (Taneda *et al.*, 1985; Rinkevich, 1992; Saito *et al.* 1994). However, we recently pointed out that a difference in hemocyte behavior, especially the behavior of morula cells (MCs), induced by nonself-recognition also contributes to the variation in allorecognition reactions (Shirae *et al.*, 1999). In *Botryllus scalaris* (Shirae *et al.*, 1999) and an unidentified species of *Botrylloides* from Israel (Rinkevich *et al.*, 1994), MCs do not have any role in the rejection reactions although they exist in the blood. However, in most other botryllid ascidians, MCs participate actively in their allorecognition reactions. From blood vessels, they infiltrate into the tunic and degenerate there to form a necrotic area. Nevertheless, the degrees of their participation are different among botryllid species. In the allorecognition of *Botrylloides*, called "subcuticular rejection" (Hirose *et al.*, 1997), the number of MCs infiltrating into the tunic is much fewer than that in the case of allorecognition in *Botryllus schlosseri* or *Botryllus primigenus*.

The MCs in some ascidians have phenoloxidase (PO) activity, which is one of the most common enzymatic activities in invertebrate immune responses. In solitary ascidians, it contributes to a cellular defense reaction (Chaga, 1980; Smith and Söderhäll, 1991; Jackson *et al.*, 1993; Akita and Hoshi, 1995; Arizza *et al.*, 1995; Parrinello, 1995; Hata *et al.*, 1998).

---

\* Corresponding author: Tel. +558-23-6358;  
FAX. +558-22-0346.  
E-mail: shirae@kurofune.shimoda.tsukuba.ac.jp

**Table 1.** Characteristics of the allojection reactions in the six colonial ascidians.

	Stages of allorecognition	Hemocyte types which mainly participate in the rejection reaction	Cut surface contact	Reference
<i>S. reptans</i>	partial dissolution of cuticle	morula cells	rejection	Mukai and Watanabe (1974) Shirae <i>et al.</i> (1999)
<i>B. scalaris</i>	ampullar fusion	phagocytes	rejection	Saito and Watababe (1982) Saito <i>et al.</i> (1994) Shirae <i>et al.</i> (1999)
<i>B. primigenus</i>	ampullar penetration	morula cells	rejection	Oka and Watanabe (1975, 1960) Tanaka and Watanabe (1973) Mukai and Watanabe (1974) Taneda and Watanabe (1982a, 1982b)
<i>B. schlosseri</i>	partial dissolution of cuticle	morula cells	rejection	Bancroft (1903) Sabbadin (1962) Mukai and Watanabe (1974) Rinkevich (1992) Rinkevich <i>et al.</i> (1994) Saito <i>et al.</i> (1994)
<i>B. simodensis</i>	partial dissolution of cuticle (subcuticular rejection)	morula cells	rejection	Hirose <i>et al.</i> (1990)
<i>B. fuscus</i>	partial dissolution of cuticle (subcuticular rejection)	morula cells	fusion (surgical fusion)	Hirose <i>et al.</i> (1994) Hirose <i>et al.</i> (1997)

Cammarata *et al.* (1997) showed that PO activity leads to the production of superoxide which contributes to cytotoxicity in the solitary ascidian *Styela plicata*. Furthermore, Ballarin *et al.* (1993, 1994, 1995, 1998) demonstrated that, in *B. schlosseri*, PO of MCs contributes to cytotoxicity and its activity increases both in the allojection area between incompatible colonies and in hemocytes cultured with incompatible blood plasma. Therefore, PO of MCs seems to be a key enzyme in the allojection reactions of botryllids.

In the present study, we tried to demonstrate that differences in the ratio of MCs to total hemocytes and in the levels of PO activity in the MCs are related to the variation in allojection reactions among botryllids. For this study, six colonial ascidians were used: *B. scalaris*, *B. primigenus*, *B. schlosseri*, *Botrylloides simodensis*, *Botrylloides fuscus*, and *Symplegma reptans*. These five botryllids show allojection reactions different from one another (Table 1). *S. reptans* of the family Styelidae, a sister family of Botryllidae (Berrill, 1936), is the species most similar morphologically to botryllids among the styelids. This species shows an allojection reaction similar to that of *B. schlosseri* (Shirae *et al.*, 1999).

## MATERIALS AND METHODS

### Animals

Colonies of *B. scalaris*, *B. primigenus*, *B. schlosseri*, *B. simodensis*, *B. fuscus* and *S. reptans* were collected in the vicinity of Shimoda (Shizuoka Prefecture, Japan) and in Uranouchi Inlet near the Usa Marine Biological Institute of Kochi University. Attached to glass slides, they were reared in culture boxes immersed in Nabeta Bay near the Shimoda Marine Research Center, University of Tsukuba. Colonies which grew well were used for the experiments.

### Histochemical analysis of PO activity in hemocytes

Histochemical analysis of PO activity was performed using the

method developed by Ballarin *et al.* (1995) with some modifications. The ascidian colonies were washed with filtered seawater (FSW) and immersed in FSW containing 10 mM L-cysteine for 5 min to prevent hemocytes from clotting during the collection of blood. After the FSW containing L-cysteine had been wiped up gently from the surfaces of the colonies, their vascular vessels were pricked with tungsten needles and the blood oozing through the prick-holes was collected. The collected blood was mixed with FSW, and hemocytes in the suspension were mounted on a glass slide coated with poly-L-lysine. This sample was then incubated in a moist chamber for 5 min to allow the hemocytes to adhere to the slide. The hemocytes on the slide were fixed with 1% glutaraldehyde in FSW containing 1% sucrose for 30 min at 4°C, and washed with FSW containing 1% sucrose. The fixed hemocytes were pre-incubated for 5 min at room temperature in 1 mM Na-cacodylate buffer (pH 8.5) containing 5 mM CaCl<sub>2</sub> (CAB-Ca), and incubated for 10 min at room temperature in a reaction mixture containing CAB-Ca plus 4% N, N'-dimethylformamide, dihydroxyphenyl-L-alanine (L-DOPA)-saturated CAB-Ca and 20.7 mM 3-methyl-2-benzothiazolinonehydrazone hydrochloride (MBTH) in the ratio of 5: 2: 3. Then, these hemocytes were washed with CAB-Ca and examined under a Nikon optiphot light microscope.

### Assay of PO activity in hemocytes

Colonies were stripped from the glass slides on which they were cultured. After removal of detritus from the colony surface, the colonies were immersed in FSW containing 10 mM L-cysteine for 5 min to prevent aggregation of hemocytes and were cut into strips. Blood oozing from the cut surfaces of these strips was collected and stored in a microcentrifuge tube on ice. The blood was then centrifuged at 780×g for 15 min. The pellet was resuspended in FSW to obtain a hemocyte suspension at a cell concentration of 10<sup>9</sup> cells/ml. A 20 µl portion of the hemocyte suspension was mixed with 20 µl of 0.1% Tween 20 in phosphate buffered saline (PBS: 0.8% NaCl, 0.02% KCl, 0.02% KH<sub>2</sub>PO<sub>4</sub>, 0.115% Na<sub>2</sub>HPO<sub>4</sub>, pH 7.2) and 160 µl of PBS on ice. This specimen was used as a "hemolysate" in the following assays.

To measure the PO activity in the hemolysate, the method used by Winder and Harris (1991) and Ballarin *et al.* (1998) was employed with some modifications. A 20 µl portion of the hemolysate was mixed with a reaction mixture consisting of 490 µl of PBS, 200 µl of L-DOPA-saturated PBS, and 290 µl of 20.7 mM MBTH in PBS containing 4%

of N, N'-dimethylfolnamide at 25°C, and, one minute after mixing, the absorbance at 505 nm was read by means of a Shimadzu UV-1200 spectrophotometer. PO activity in the hemocyte sample from each colony was measured in triplicate. The average was taken to be representative of the hemocyte PO activity in each colony. For each species, six colonies were examined. Data are expressed as means  $\pm$  standard error (SE). Differences among the data for these species were analyzed statistically by the Kruskal- Wallis test.

### Experiments on inhibition of PO activity

The effects of the following reagents on the PO activity of the hemolysate were investigated: the PO inhibitors Na-benzoate (20 mM) and tropolone (2 mM), the antioxidants L-cysteine (2 mM) and ascorbic acid (2 mM), and the scavenger enzymes superoxide dismutase (SOD) (120 U/ml) and catalase (140 U/ml). In order to evaluate the inhibitory effects on PO activity, these reagents were added to the reaction mixture described above. Then, the mixture was incubated for 10 min at 25°C, and its absorbance at 505 nm was measured. As a control, a mixture without any of these reagents added was used. In this assay, for each species, from three to six colonies were examined. The decrease in absorbance was calculated for each colony, and the average of the data for three colonies was taken to be representative of the inhibitory effect of a particular reagent on the PO activity in each species. Data are expressed as means  $\pm$  standard deviation (SD). Significance was determined using the Wilcoxon Matched-Pairs Signed-Ranks Test.

### Ratio of cell types among total hemocytes

Ascidian colonies were washed with FSW and fixed with 10% formaldehyde in FSW, pH 7.4, at 4°C overnight. After that, they were rinsed with FSW. Vascular vessels of the colony were then cut with a razor blade, and the blood oozing through the cut surface was col-

lected with a micropipette and smeared on glass slides. After rinsing with FSW, the hemocytes were counted under a light microscope to obtain the ratio of each hemocyte type to total hemocytes (at least 200 hemocytes were counted per slide). Five colonies were examined for each species.

### PO activity in the allorejection area


Colonies undergoing rejection were washed with FSW and fixed with 2% glutaraldehyde in FSW containing 1% sucrose for 1 hr at 4°C. Then, they were washed again with FSW containing 1% sucrose. These fixed colonies were incubated for 2-3 min at room temperature in a mixture consisting of CAB-Ca plus 4% N, N'-dimethylformamide, L-DOPA-saturated CAB-Ca and 20.7 mM MBTH in the same ratio as that used in the histochemical study of hemocytes. The colony parts, stained dark pink due to accumulation of dopa-quinones, were examined under a Nikon optiphoto light microscope and an Olympus IX-70 inverted microscope. As a control, colonies undergoing fusion were also examined.

## RESULTS

### Morphology and PO activity of the hemocytes in the colonial ascidians

We classified the hemocytes of the six ascidians examined here into seven types principally, on the basis of the previous studies (Goodbody, 1974; Wright, 1981; Rowley *et al.*, 1982; Ballarin *et al.*, 1994), 1) hemoblasts, 2) hyaline amebocytes, 3) signet ring cells, 4) macrophages, 5) granular leukocytes, 6) pigment cells, and 7) MCs (Fig. 1).

The hemoblasts (Fig. 1a) were small spherical cells, 4–5




**Fig. 1.** Morphology of living hemocytes incubated in FSW.

a, hemoblast in *B. simodensis*; b, morula cell in *B. scalaris*; c, granular leukocytes (arrow) in *B. primigenus*; d, pigment cell in *B. fuscus*; e, hyaline amebocyte in *B. simodensis*; f, macrophage in *B. simodensis*; g, signet ring cell (arrow) in *B. fuscus*; h, morula cell in *S. reptans*; i, morula cell and pigment cell in *B. primigenus*; j, granular leukocyte in *B. fuscus*; k, granular leukocyte in *B. scalaris*. M, morula cell; P, pigment cell. Scale bar=10  $\mu$ m.

$\mu\text{m}$  in diameter, with a large nucleus. The hyaline amoebocytes (Fig. 1e) were flat and variable in shape, about 6–20  $\mu\text{m}$  in diameter, and they contained small granules. The macrophages (Fig. 1f) were large spherical cells, 12–15  $\mu\text{m}$  in diameter, with a few large vacuoles filled with ingested materials. The signet ring cells (Fig. 1g) were spherical cells, about 5–10  $\mu\text{m}$  in diameter, and they contained a vacuole which pushed the cytoplasm and the nucleus to the periphery. The granular leukocytes (Fig. 1c) were principally spherical in shape, about 6–8  $\mu\text{m}$  in diameter, and they were filled with many small granules, about 1  $\mu\text{m}$  or less in diameter. In addition to this

common type of granular leukocyte, in the two *Botrylloides* species, *B. simodensis* and *B. fuscus*, a specific type of granular leukocyte, which is larger than the common type in the six ascidians (about 10  $\mu\text{m}$  in diameter) was often seen among their hemocytes (Fig. 1j). In *B. scalaris*, another species-specific type of granular leukocyte was found (Fig. 1k). They were about 12–20  $\mu\text{m}$  in diameter and had small red granules (about 1–2  $\mu\text{m}$  in diameter) and white granules that appeared to be thick and short sticks and these were 5–8  $\mu\text{m}$  long in the major axis in both fixed and fresh samples (data not shown). The pigment cells (Fig. 1d) were large vacuolated cells, 12–




**Fig. 2.** Moving morula cells of two *Botrylloides*. Living morula cells changed their shapes during incubation in FSW, as seen by phase-contrast microscopy. a, morula cell in *B. simodensis*; b, after 20 sec; c, after 40 sec; d, morula cell in *B. fuscus*; e, after 20 sec; f, after 40 sec. Scale bar=10  $\mu\text{m}$ .

15  $\mu\text{m}$  in diameter, with one or a few large vacuoles (6–12  $\mu\text{m}$  in diameter) containing many small granules about 0.5  $\mu\text{m}$  in diameter. The color of these granules determines the color of the colony (Hirose *et al.*, 1998). Only in *B. primigenus*, there was a type of the pigment cells resembling the MC in size and shape, and the vacuoles in these cells were filled with a purple-colored material (Fig. 1i).

The MCs of the three botryllids, *B. schlosseri*, *B. primigenus*, and *B. scalaris*, were morphologically very similar to one another. Generally, the MCs are 6–10  $\mu\text{m}$  in diameter and they were filled with many vacuoles about 2  $\mu\text{m}$  in diameter (Fig.

1b). In *B. simodensis* and *B. fuscus*, the vacuoles of the MCs were irregular in size (2–8  $\mu\text{m}$  in diameter). Especially, the MCs of *B. fuscus* formed many long pseudopodia and showed various shapes during incubation (Fig. 2). The MCs of *S. reptans* contained several red granules, which were less than 0.5  $\mu\text{m}$  in diameter (Fig. 1h).

In all of the ascidians, most MCs showed dark pink coloration in the L-DOPA-MBTH assay (Fig. 3). The degree of coloration dependent on PO activity varied among the MCs in each species. In *B. scalaris*, the species-specific granular leukocytes also showed weak PO activity (Fig.3b). In *B.*




**Fig. 3.** Phenoloxidase activity in hemocytes as determined by the L-DOPA-MBTH assay. Morula cells of all botryllids showed dark-pink coloration. a, *S. reptans*; b, *B. scalaris*. The granular leukocytes of *B. scalaris* with pigmentation also showed such coloration in their cytoplasm. c, *B. primigenus*. In *B. primigenus*, the pigment cells resemble morula cells morphologically and their vacuoles include purple-colored components. In this assay, the pigment cells did not show any increase in dark pink coloration. d, *B. fuscus*. M, morula cell; G, granular leukocyte; P, pigment cell. Scale bar=10  $\mu\text{m}$ .


**Table 2.** Effects various reagents on phenoloxidase activity

	Na-benzoate 20 mM	tropolone 2 mM	ascorbic acid 2 mM	cysteine 2 mM	SOD 120U/ml	catalase 140U/ml
	Inhibition(%)					
<i>S. reptans</i>	97.5±0.8%	99.3±0.8%	98.9± 0.8%	99.1±0.4%	20.9± 8.6%	21.1±5.4%
<i>B. fuscus</i>	95.0±3.8	96.4±3.7	93.6± 5.8	98.0±2.0	8.3± 5.1	–
<i>B. simodensis</i>	97.2±3.1	98.9±0.6	97.2± 2.5	98.4±0.5	36.3±14.6	–
<i>B. schlosseri</i>	98.3±0.0	99.1±0.1	98.4± 0.8	98.7±0.5	24.3±24.0	–
<i>B. primigenus</i>	96.8±2.2	97.3±3.6	98.4± 1.5	97.3±2.6	29.7±14.7	–
<i>B. scalaris</i>	88.9±4.6	91.7±8.1	81.5±23.6	97.9±4.3	35.7±11.5	–

Data are expressed as means  $\pm$  SD. Bars mean no reduction of absorbance in the experiments.



**Fig. 4.** PO activity in the allorejection reaction area in colonial ascidians as determined by the L-DOPA-MBTH assay after fixation in 2% glutaraldehyde. a, *S. reptans* colonies before treatment; b, *S. reptans* colonies after treatment; c, *B. scalaris* colonies before treatment; d, *B. scalaris* colonies after treatment; e, *B. primigenus* colonies before treatment; f, *B. primigenus* colonies after treatment; g, *B. fuscus* colonies before treatment; h, *B. fuscus* colonies after treatment. Arrowheads indicate the allorejection sites. Arrows indicate ampullae not undergoing rejection in *B. scalaris*. Scale bar=1mm.



**Fig. 5.** Allorejection reaction area as determined by the L-DOPA-MBTH assay, at high magnification. a, *S. reptans* ampullae before treatment; b, *S. reptans* ampullae after treatment; c, *B. scalaris* ampullae before treatment; d, *B. scalaris* ampullae after treatment; e, *B. primigenus* ampullae before treatment; f, *B. primigenus* ampullae after treatment; g, *B. fuscus* ampulla before treatment; h, *B. fuscus* ampulla after treatment. Scale bar=50  $\mu$ m.




*primigenus*, the MC-like pigment cells did not show further coloration by dopa-quinones (Fig. 3c).

#### Ratios of MCs to total hemocytes

The ratio of MCs to hemocytes differed among the six ascidians. *B. primigenus* showed the highest MC ratio ( $54.1 \pm 8.1\%$ ), followed by *S. reptans* ( $46.8 \pm 10.0\%$ ), *B. simodensis* ( $39.9 \pm 11.4\%$ ), *B. schlosseri* ( $37.5 \pm 5.2\%$ ), and *B. scalaris* ( $14.6 \pm 8.3\%$ ) in that order, and the lowest MC ratio was that observed in the case of *B. fuscus* ( $8.5 \pm 2.2\%$ ). In *B. scalaris*, the ratio of granular leukocytes, which showed little PO activity, to hemocytes was  $24.0 \pm 11.8\%$ .

#### PO activity of hemocytes

In the six ascidians studied here, their hemolysates, prepared from the same number of hemocytes in each instance, showed significantly different levels of PO activity ( $\Delta A_{505} / \text{min} \times 100$ ) in the first minute after mixing with the reaction mixture ( $p < 0.001$ ). *B. schlosseri* showed the highest level of activity ( $22.0 \pm 2.5$ ) followed by *S. reptans* ( $17.7 \pm 2.3$ ), *B. primigenus* ( $5.6 \pm 2.4$ ), *B. simodensis* ( $3.2 \pm 0.7$ ), and *B. scalaris* ( $0.8 \pm 0.1$ ) in that order. The lowest level of activity was that observed in the case of *B. fuscus* ( $0.3 \pm 0.1$ ).



**Fig. 6.** Fusion reaction area as determined by the L-DOPA-MBTH assay in *B. primigenus*. a, colonies before treatment; b, colonies after treatment. Scale bar=1mm.

### Effects of inhibitors on PO activity

The effects of six reagents on the PO activity in the six ascidians are summarized in Table 2. The PO activity in all six species was almost completely inhibited by Na-benzoate (20 mM) or tropolone (2 mM), and was fully inhibited by L-cysteine (2 mM) or ascorbic acid (2 mM;  $p < 0.05$ ). Upon incubation with SOD (120 U/ml), there was a significant decrease in absorbance at 505nm ( $p < 0.05$ ), whereas the other scavenger enzyme tested, catalase, had no effect on the PO activity in the botryllids even at a high concentration, 140 U/ml. Only in the case of *S. reptans*, catalase (140 U/ml) induced a significant decrease in absorbance at 505nm ( $p < 0.05$ ), similar in degree to that observed with SOD (120 U/ml). The effect of each of these reagents on the PO activity in MCs was similar for all of the botryllids. Histochemical observations showed that the coloration by dopa-quinone in the MCs of all of the ascidians and in the granular leukocytes of *B. scalaris* was inhibited by these PO inhibitors and antioxidants. The coloration in the hemocytes was inhibited by the reagents at concentrations ten times as high as those required in the case of the hemolysate (data not shown).

### PO activity in the allorejection areas

Even in the fixed normal colonies, MCs showed strong PO activity upon incubation with the reaction mixture containing L-DOPA and MBTH. Therefore, in this experiment, the incubation time was set at 2–3 min, at room temperature, to obtain better contrast between the rejection area and the non-rejection area (Fig. 4). In *S. reptans* and the four botryllids other than *B. scalaris*, a higher level of PO activity was observed in the allorejection area than in the non-rejection area or the fusion area between compatible colonies. The dark pink colored area was consistent with the range of the rejection area in each species, like the result obtained for *B. schlosseri* as shown by Ballarin *et al.* (1995). PO activity outside of MCs was also found in the tunic in the rejection area, especially around the MCs that infiltrated into the tunic (Figs. 5b and 5f). In *B. scalaris*, PO activity was observed only inside of the vascular vessels, and the change in coloration of the hemocyte aggregates in the ampullae undergoing rejection was not marked (Figs. 4d and 5d). In *B. fuscus*, although the ampullae undergoing rejection and the infiltrating MCs were dark pink in color, PO activity was not observed in the tunic around the infiltrating MCs (Figs. 4h and 5h). In the fusion reactions between syngeneic colonies in the case of all of the ascidians, an increase in PO activity was not evident in the fusion area (Fig. 6).

## DISCUSSION

This comparative study of hemocyte morphology among botryllid ascidians and *S. reptans* has made clear that their hemocytes consist of the same cell types basically. However, in a few cases, hemocytes with genus- or species-specific characteristics were found and these might serve as marker hemocytes in particular species. In *B. primigenus*, the pig-

ment cells resembled MCs morphologically except for the purple-coloration of their granules. However, these pigment cells did not display any PO activity. As observed under the light microscope, these cells were mostly deposited under the epithelium of the zooids and they contributed to the purple color of the zooids (data not shown). These cells were also seen in the ampullae, although they never infiltrated into the tunic in the allorejection reaction. The pigment cells and the MCs surely have a different function and a different origin in terms of cell lineage. Therefore, we should be careful not to confuse these two types of hemocytes in the sectional specimens of *B. primigenus*.

The MCs of *B. fuscus* do not have sufficient granules to fill up the cytoplasm, and there is more cytoplasmic matrix than in the case of the MCs of other species. The MCs of this species form philopodia actively and move much during incubation in FSW. The MCs of *S. reptans* were very distinctive in terms of having red granules. However, because of the common morphological characteristics, they can be classified into the same hemocyte type. The MCs of the botryllids and *S. reptans* were found to have PO activity in their vacuoles, as in the case of *B. schlosseri* and some solitary ascidians (Ballarin *et al.*, 1993; Jackson *et al.*, 1993; Akita and Hoshi, 1995; Arizza *et al.*, 1995). Therefore, PO activity is regarded as one of the common features of MCs in these ascidians. In *B. scalaris*, the granular leukocytes also showed weak PO activity in their cytoplasm. As the PO inhibitors and antioxidants had the same inhibitory effect on the PO activity of the granular leukocytes as that observed in the case of MCs, the PO enzymes present in these two types of hemocytes may be highly similar. In this species, when a colony is injured and its blood oozes out, the granular leukocytes immediately form aggregates to stop the blood leak (data not shown). The role of these hemocytes, especially in relation to the PO activity, should be further studied in *B. scalaris*.

Both the ratio of MCs to total hemocytes and the PO activity of the hemocytes varied among the ascidians examined, and the differences in these two factors in these ascidians were not parallel. The variation in PO activity among species is more conspicuous when the PO activity per MC is estimated. The level of PO activity in the hemocytes was highest in *B. schlosseri* and *S. reptans* among the ascidians examined. *B. simodensis* and *B. primigenus* showed a lower level of PO activity than *B. schlosseri*, but the ratio of MCs to total hemocytes was almost the same for both, and even higher than that in the case of *B. schlosseri*. In these four species, PO activity was observed only in the MCs. These results suggest that in *B. simodensis* and *B. primigenus*, the PO activity per MC is lower than that in *B. schlosseri*. In contrast, in *B. scalaris* and *B. fuscus*, both the MC ratio and the PO activity per MC were markedly lower than those in *B. schlosseri*. In *B. scalaris*, the granular leukocytes also had PO activity, although the level of activity was lower than that in the MCs as described above. Therefore, without the granular leukocytes taken into account, the level of PO activity in the MCs would be estimated as being much lower. In the other experiment

concerning PO activity, the characteristics of the PO in the six ascidians were found to be similar in terms of the effects of inhibitors, except for the weak inhibition by catalase in the case of *S. reptans*. These results suggest that in the hemocytes of the six ascidians, the difference in PO activity is attributable to the two factors, the ratio of MCs to total hemocytes and the PO activity per MC.

The PO activity might be related to the allojection modes in colony specificity. In the compound ascidians examined here, the morphological processes of fusion and allojection have already been described in detail. Table 1 is a list of the characteristics of the allojection reactions that have been reported in five botryllids and *S. reptans*. A close correlation between the characteristics of the MCs and the allojection reaction is obvious as follows. 1) *B. schlosseri* and *S. reptans*, which have hemocytes with a high level of PO activity, show a restricted allojection reaction. In both species, the allojection reaction is initiated after their ampullae come into contact, and MC infiltration and disintegration of these cells leads to severe necrosis around the ampullae undergoing rejection. 2) In *B. scalaris*, the behavior of hemocytes in the allojection reaction is different from that in the case of the other botryllids and *S. reptans*. The allojection reaction is initiated in the latest stage of the fusion process, after vascular fusion and blood exchange. In the allojection reaction in this case, the MCs do not perform any noteworthy role. The blood exchange between two incompatible colonies, however, does not involve necrosis with disintegration of hemocytes. The low level of PO activity might affect this mild rejection reaction in *B. scalaris*. 3) *B. fuscus* and *B. simodensis* belong to the same genus *Botrylloides* (Monniot, 1988). Recent molecular analysis has demonstrated that these two species are more closely related to each other than to the other botryllids (Cohen *et al.*, 1998). However, *B. fuscus* has a much lower proportion of MCs and also has a much lower level of PO activity than *B. simodensis*. Whereas they show the same mode of allojection in the case of growing edge contact, these two species show opposite reactions in the case of contact between the cut surfaces of two incompatible colonies (Table 1). In *B. fuscus*, two incompatible colonies become fused without infiltration or disintegration of MCs (Hirose *et al.*, 1994). In contrast, in *B. simodensis*, severe necrosis is caused by MC disintegration in the contact area (Hirose *et al.*, 1990). The lower level of PO activity and the lower ratio of MCs to total hemocytes in the case of *B. fuscus* than those in the case of *B. simodensis* might be related to the occurrence of artificial fusion between two incompatible colonies.

Previously we demonstrated that the difference in allojection modes might be due to difference in MC behavior, and in this study we showed that there is a correlation between the allojection mode and the ratio of MCs to total hemocytes and their PO activity among botryllids. To understand the mechanisms responsible for the variation in allojection modes in botryllids, further comparative studies will be necessary, such as studies on various factors related to the proPO system, MC infiltration, MC degranulation, and

the allogeneic recognition system.

## ACKNOWLEDGMENTS

This study was supported in part by a grant-in-aid to Y. S. (#08459005) from the Ministry of Education, Science and Culture of Japan. We are grateful to the staff of the Shimoda Marine Research Center (SMRC), University of Tsukuba, for their assistance and hospitality during this work. We also thank Dr. E. Hirose (University of Ryukyus) and Dr. T. Ishii (Akita University) for their advice, and Dr. K. Kawamura (Kochi University) and the staff of Usa marine institute for collecting *S. reptans*. Finally, special thanks to Dr. A. Bellgrove for her kindness in reading the manuscript for grammatical correctness. This report is contribution No. 647 from SMRC.

## REFERENCES

- Akita N and Hoshi M (1995) Hemocytes release phenoloxidase upon contact reaction, an allogeneic interaction, in the ascidian *Halochynthia roretzi*. *Cell Struct Func* 20: 81–87
- Arizza V, Cammarata M, Tomasino CM and Parrinello N (1995) Phenoloxidase characterization in vacuolar hemocytes from the solitary ascidian *Styela plicata*. *J Invert Pathol* 66: 297–302
- Ballarin L, Cima F and Sabbadin A (1993) Histochemical staining and characterization of the colonial ascidian *Botryllus schlosseri* hemocytes. *Bol Zool* 60: 19–24
- Ballarin L, Cima F and Sabbadin A (1994) Phenoloxidase in the colonial ascidian *Botryllus schlosseri* (Urochordata, Ascidiacea). *Anim Biol* 3: 41–48
- Ballarin L, Cima F and Sabbadin A (1995) Morula cells and histocompatibility in the colonial ascidian *Botryllus schlosseri*. *Zool Sci* 12: 757–764
- Ballarin L, Cima F and Sabbadin A (1998) Phenoloxidase and cytotoxicity in the compound ascidian *Botryllus schlosseri*. *Dev Com Immunol* 22: 479–492
- Bancroft FW (1903) Variation and fusion of colonies in compound ascidians. *Proc Calif Acad Sci* 3: 137–186
- Cammarata M, Arizza V, Parrinello N, Candore G and Caruso C (1997) Phenoloxidase-dependent cytotoxic mechanism in ascidian (*Styela plicata*) hemocytes active against erythrocytes and K562 tumor cells. *Eur J Cell Biol* 74: 302–307
- Chaga OY (1980) Ortho-diphenoloxidase system of Ascidians. *Tsitologia* 22: 619–625
- Cohen SC, Saito Y and Weissman IL (1998) Evolution of allorecognition in botryllid ascidians inferred from molecular phylogeny. *Evolution* 52: 746–756
- Goodbody I (1974) The physiology of ascidians. *Adv Mar Biol* 12: 1–149
- Hata S, Azumi K and Yokosawa H (1998) Ascidian phenoloxidase: its release from hemocytes, isolation, characterization and physiological roles. *Comp Biochem Physiol B* 119: 769–776
- Hirose E, Saito Y and Watanabe H (1988) A new type of the manifestation of colony specificity in the compound ascidian, *Botrylloides violaceus* Oka. *Biol Bull* 175: 240–245
- Hirose E, Saito Y and Watanabe H (1990) Allogeneic rejection induced by cut surface contact in the compound ascidian, *Botrylloides simodensis*. *Invertebr Reprod Dev* 17: 159–164
- Hirose E, Saito Y and Watanabe H (1994) Surgical fusion between incompatible colonies of the compound ascidian, *Botrylloides fuscus*. *Dev Com Immunol* 18: 287–294
- Hirose E, Saito Y and Watanabe H (1997) Subcuticular rejection: An advanced mode of the allogeneic rejection in the compound ascidians, *Botrylloides simodensis* and *B. fuscus*. *Biol Bull* 192: 53–61
- Hirose E, Yoshida T, Akiyama T, Ito S and Iwanami T (1998) Pigment cells representing polychromatic colony color in *Botrylloides*

- simodensis* (Ascidacea, Urochordata): cell morphology and pigment substances. *Zool Sci* 15: 489–497
- Jackson AD, Smith VJ and Peddie CM (1993) In vitro phenoloxidase activity in the blood of *Ciona intestinalis* and other ascidians. *Dev Comp Immunol* 17: 97–108
- Monniot C (1988) Ascidiées de Nouvelle-Calédonie. IV. Styelidae (suite). *Bul Mus Nat Hist Nat Paris* 10:163–196
- Mukai H and Watanabe H (1974) On the occurrence of colony specificity in some compound ascidians. *Biol Bull* 147:411–421
- Oka H and Watanabe H (1957) Colony specificity in compound ascidians as tested by fusion experiments (a preliminary report). *Proc Jpn Acad* 33: 657–659
- Oka H and Watanabe H (1960) Problems of colony specificity in compound ascidians. *Bull Mar Biol Stat Asamushi* 10: 153–155
- Parrinello N (1995) Cytotoxic activity of tunicate hemocytes. In *Cellular, Biochemical, and Molecular Aspects of Invertebrate Immunology*. Progress Mol. Subc. Biol. (Müller WEG and Rinkevich B, Eds.), Wiley, New York. Vol. 15, pp 190–217
- Rinkevich B (1992) Aspects of the incompatibility nature in botryllid ascidians. *Anim Biol* 1: 17–28
- Rinkevich B, Linker-Levav T and Goren M (1994) Allorecognition / xenorecognition responses in *Botrylloides* (Ascidacea) subpopulations from the Mediterranean coast of Israel. *J Exp Zool* 270: 302–313
- Rowley AF (1982) The blood cells of *Ciona intestinalis*: an electron probe X-ray microanalytical study. *J Mar Biol Ass UK* 62: 607–620
- Sabbadin A (1962) Le basi genetiche della capacità di fusione fra colonie in *Botryllus schlosseri* (Ascidacea). *Rend Accad Naz Lincei Ser 8* 32: 1031–1035
- Saito Y and Watanabe H (1982) Colony specificity in the compound ascidian *Botryllus scalaris*. *Proc Jpn Acad B* 58: 105–108
- Saito Y, Hirose E and Watanabe H (1994) Colony specificity in botryllid ascidians. *Int J Dev Biol* 38: 237–247
- Shirae M, Hirose E and Saito Y (1999) Behavior of hemocytes in the allorejection reaction in two compound ascidians, *Botryllus scalaris* and *Symplesma reptans*. *Biol Bull* 197: 188–197
- Smith VJ and Söderhäll K (1991) A comparison of phenoloxidase activity in the blood of marine invertebrates. *Dev Comp Immunol* 15: 251–261
- Tanaka K and Watanabe H (1973) Allogeneic inhibition in a compound ascidian, *Botryllus primigenus* Oka. I. Processes and features of “nonfusion” reaction. *Cell Immunol* 7: 410–426
- Taneda Y and Watanabe H (1982a) Studies on colony specificity in the compound ascidian, *Botryllus primigenus* Oka. I. Initiation of “nonfusion” reaction with special reference to blood cell infiltration. *Dev Comp Immunol* 6: 43–52
- Taneda Y and Watanabe H (1982b) Studies on colony specificity in the compound ascidian, *Botryllus primigenus* Oka. II. In vivo bioassay for analyzing the mechanism of “nonfusion” reaction. *Dev Comp Immunol* 6: 243–252
- Taneda Y, Saito Y and Watanabe H (1985) Self or non-self discrimination in ascidians. *Zool Sci* 2: 433–442
- Winder AJ and Harris H (1991) New assays for tyrosine hydroxylase and dopa oxidase activities of tyrosinase. *Eur J Biochem* 198: 317–326
- Wright RK (1981) Urochordates. In *Invertebrate Blood Cells*. Vol. 2. (Ratcliffe NA and Rowley AF, Eds.), Academic Press, London. pp 565–626

(Received February 9, 2000 / Accepted April, 2000)