



## **Identification and Characterization of Microsatellite Markers in *Pinus kesiya* var. *langbianensis* (Pinaceae)**

Authors: Cai, Nian-Hui, Xu, Yu-Lan, Wang, Da-Wei, Chen, Shi, and Li, Gen-Qian

Source: Applications in Plant Sciences, 5(2)

Published By: Botanical Society of America

URL: <https://doi.org/10.3732/apps.1600126>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](http://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

## IDENTIFICATION AND CHARACTERIZATION OF MICROSATELLITE MARKERS IN *PINUS KESIYA* VAR. *LANGBIANENSIS* (PINACEAE)<sup>1</sup>

NIAN-HUI CAI<sup>2,3</sup>, YU-LAN XU<sup>2,3</sup>, DA-WEI WANG<sup>2</sup>, SHI CHEN<sup>3</sup>, AND GEN-QIAN LI<sup>3,4</sup>

<sup>2</sup>Key Laboratory for Forest Genetic and Tree Improvement and Propagation in Universities of Yunnan Province, Southwest Forestry University, Kunming, Yunnan 650224, People's Republic of China; and <sup>3</sup>Key Laboratory for Forest Resources Conservation and Use in the Southwest Mountains of China, College of Forestry, Southwest Forestry University, Kunming, Yunnan 650224, People's Republic of China

- **Premise of the study:** Microsatellite primers were developed in *Pinus kesiya* var. *langbianensis* (Pinaceae), a species native to southwestern China, to investigate its genetic diversity and population structure in order to provide information for the conservation and management of this species.
- **Methods and Results:** Using next-generation sequencing, a total of 2349 putative simple sequence repeat primer pairs were designed. Eighteen polymorphic markers in 60 individuals belonging to four populations of *P. kesiya* var. *langbianensis* were identified and characterized with two to 11 alleles per locus. The observed and expected heterozygosity ranged from 0.000 to 0.800 and 0.000 to 0.840, respectively. Each of these loci cross-amplified in the closely related species *P. massoniana*, *P. densata*, *P. tabuliformis*, and *P. yunnanensis*, with one to seven alleles per locus.
- **Conclusions:** The new markers are promising tools to study the population genetics of *P. kesiya* var. *langbianensis* and related species.

**Key words:** microsatellite; next-generation sequencing; Pinaceae; *Pinus kesiya* var. *langbianensis*; population genetics.

*Pinus kesiya* Royle ex Gordon var. *langbianensis* (A. Chev.) Gaussen (Pinaceae) is an important forest tree species in Yunnan Province, China. It has been recorded at altitudes from 600–1800 m in the southern, semihumid climate zone of Yunnan (Editorial Committee of Flora of China, 1978; Wu, 1986) and accounts for 11% of the forest area and  $1.0 \times 10^8$  m<sup>3</sup> of the forest volume (Jiang et al., 2007). The wood is extensively used in building, furniture, and the fiber industry. *Pinus kesiya* var. *langbianensis* is also highly valued for its high resin content, with an annual output of 179,100,000 kg (Editorial Committee of Flora of China, 1978; Wu, 1986; Dong et al., 2009). Output of gum turpentine from *P. kesiya* var. *langbianensis* accounted for more than 90% of the total output in Yunnan (Yin et al., 2005). However, germplasm resources of *P. kesiya* var. *langbianensis* have decreased in recent years as a result of overexploitation (Zhao et al., 2016).

Information on genetic diversity and spatial structure in *P. kesiya* var. *langbianensis* is important for its future conservation and can be used to help guide local forest management (Sanchez et al., 2014). No specific conservation strategy is available for this species, in part due to the limited understanding of genetic diversity and structure of the natural populations. As a primary forest

tree species in southern Yunnan Province, resource conservation of *P. kesiya* var. *langbianensis* will benefit the entire ecological system in the region (Li et al., 2015). Therefore, in this study we developed novel microsatellite markers for *P. kesiya* var. *langbianensis* by applying next-generation sequencing to investigate the genetic diversity and population structure of this species at the molecular level.

### METHODS AND RESULTS

Needle samples of 60 individuals from four *P. kesiya* var. *langbianensis* populations located in Yunnan Province, China, and 59 individuals from four related species (*P. massoniana* D. Don, *P. densata* Mast, *P. tabuliformis* Carrière, *P. yunnanensis* Franch.) were collected (Appendix 1). All needle samples were dried and preserved in silica gel. Total genomic DNA was isolated from dried needle samples using the cetyltrimethylammonium bromide (CTAB) method (Doyle and Doyle, 1990). Paired-end libraries were constructed on four individuals sampled from Puer City Institute of Forestry Sciences and sequenced by a customer sequencing service (Beijing Honor Tech Co. Ltd., Beijing, China) using the Illumina HiSeq 2500 Sequencing System (Illumina, San Diego, California, USA). Clean reads were assembled using Trinity version 2.2.0 (<https://github.com/trinityrnaseq/trinityrnaseq/wiki>; Grabherr et al., 2011; Haas et al., 2013). Data quality control was carried out with the software FastQC (Andrews, 2010). The Q30 percentage exceeded 90% and the GC contents were 45.60–46.56%, which suggests that the sequencing was highly reliable. Data filtering was carried out according to the following criteria: (1) removed reads with adapters; (2) removed reads with unknown bases >10%; and (3) removed low-quality reads (defined as reads having >50% bases with quality value ≤5). A total of 104,392 unigenes were obtained with an N50 length of 1349 bp. The data have been deposited in the Short Read Archive (SRA) database of the National Center for Biotechnology Information (NCBI; accession no. SRP093696). Genomic microsatellite loci for *P. kesiya* var. *langbianensis* were detected using the software MicroSatellite Identification Tool (MISA; Thiel et al., 2003). Two

<sup>1</sup>Manuscript received 9 October 2016; revision accepted 20 December 2016.

This work was supported by the National Natural Science Foundation of China (grant no. 31360189, 31260191, 31500536) and the Foundation of Southwest Forestry University (grant no. 01102-111436).

<sup>4</sup>Author for correspondence: [ligenqian0016@163.com](mailto:ligenqian0016@163.com)

doi:10.3732/apps.1600126

*Applications in Plant Sciences* 2017 5(2): 1600126; <http://www.bioone.org/loi/apps> © 2017 Cai et al. Published by the Botanical Society of America. This is an open access article distributed under the terms of the Creative Commons Attribution License (CC-BY-NC-SA 4.0), which permits unrestricted noncommercial use and redistribution provided that the original author and source are credited and the new work is distributed under the same license as the original.

TABLE 1. Characteristics of 18 polymorphic microsatellite loci in *Pinus kesiya* var. *langbianensis*.

Locus	Primer sequences (5′–3′)	Repeat motif	Allele size range (bp)	T <sub>a</sub> (°C)	GenBank accession no.
Pkvl001	F: TTTGCAGTCTGTTGCCCTTTG R: GTGGAGGAAGATGGAAACGA	(GCC) <sub>6</sub>	149–179	60.0	KX519328
Pkvl002	F: GCACGTGGATTTCAGGTTT R: GCAACTCCAGCAACTCCTTC	(GCA) <sub>5</sub>	134–146	60.0	KX519329
Pkvl003	F: CAGGCTGTAACGCTCAATCA R: TTGAAGGATCCCAACTTCCTT	(CAG) <sub>5</sub>	151–163	60.0	KX519330
Pkvl004	F: CCCTGAGTTTGACCCGACAGT R: GGGCATTTTAATGCTGCTGT	(AAT) <sub>5</sub>	265–280	60.1	KX519331
Pkvl005	F: GGGATTGGACTGGATGAGAA R: TTTTGTAAACTGCAGCCGTG	(AGA) <sub>5</sub>	241–256	59.9	KX519332
Pkvl006	F: GCAGTGGCTGACATTTGAGA R: GCTTCACGCCGTTCTTTATC	(CTG) <sub>6</sub>	147–174	59.9	KX519333
Pkvl007	F: GTAGGCACTCGTGAAGCTC R: ATCGGGACCTGTTGTTTCAG	(CAG) <sub>5</sub>	240–255	60.0	KX519334
Pkvl008	F: TGGGAAGTTTGCCATGTCA R: GCATTGTTGGCGTTGTATTG	(TCT) <sub>5</sub>	258–273	60.0	KX519335
Pkvl009	F: GTGGTCTGAAAATACCGCGT R: GCAGCAGTAGCCATCATCAA	(TCT) <sub>5</sub>	174–180	60.0	KX519336
Pkvl010	F: CGAGACGAAGCAATTCCAA R: CGAAGATCAAGAAAGGCAGG	(AT) <sub>6</sub>	244–262	59.9	KX519337
Pkvl011	F: AGCTCGGGATCAGGAGTATG R: TTTGGTGAAGTTTGGATTGCC	(TA) <sub>6</sub>	169–183	59.6	KX519338
Pkvl012	F: GTTCTGTGAGCTTGGGGAAG R: TAAGACCATTGGCTACGG	(AT) <sub>6</sub>	144–166	60.0	KX519339
Pkvl013	F: GATTGGCAGAGGCTACAAGC R: ATGCTTCCGCTGTCAACTT	(TA) <sub>11</sub>	114–150	59.9	KX519340
Pkvl014	F: TGCCACATTTGGGTAAAGAAA R: TCGGAATGATGGATAGGAGC	(AT) <sub>9</sub>	224–252	59.5	KX519341
Pkvl015	F: CCTTTTATGGGGCGGATAAT R: TTGACTTGAACACAAAGCCG	(AT) <sub>9</sub>	208–224	59.9	KX519342
Pkvl016	F: ATCCTTACGCTGCAGAGAA R: CATCGATTGGCTACATCAG	(AT) <sub>7</sub>	152–158	60.2	KX519343
Pkvl017	F: GATGCATTTGGATCAGCAAA R: ACCAATCGCTTGCATCTTTC	(TA) <sub>9</sub>	226–242	59.9	KX519344
Pkvl018	F: CTGGTGCATAGACCCGAGAT R: TTTTCTGCTTCAAGTGGCCT	(TA) <sub>9</sub>	235–257	60.0	KX519345

Note: T<sub>a</sub> = annealing temperature.

thousand three hundred forty-nine simple sequence repeat (SSR) loci were designed with Primer Premier 5.0 (PREMIER Biosoft International, Palo Alto, California, USA). Among them, 192 SSR loci with dinucleotide or trinucleotide SSR motifs were randomly chosen to screen using four individuals from four populations. One hundred fifty-nine out of 192 SSR loci were amplified successfully, and 79 out of 159 SSR loci were polymorphic. Eighteen polymorphic SSR loci (Table 1) were then randomly selected for characterization using 60 individuals from four populations (Appendix 1). The SSR amplifications were multiplexed in a 10-μL reaction containing 30 ng of genomic DNA, 0.15 μM of each primer, 5 μL Mix (0.05 units/μL *Taq* DNA Polymerase, 0.4 mM dNTPs, 4.0 mM MgCl<sub>2</sub>; Beijing Ruibio Biotech Co. Ltd., Beijing, China), and 1× PCR Buffer. The amplification protocol was: 95°C for 5 min; followed by 30 cycles at 95°C for 30 s, the annealing temperature for each primer (Table 1) for 30 s, and 72°C for 30 s; and a final extension step at 72°C for 7 min.

Amplification products were resolved using capillary electrophoresis on an ABI 3730xl DNA Analyzer (Applied Biosystems, Waltham, Massachusetts, USA). The electropherograms were analyzed using GeneMarker version 2.2.0 with GeneScan 500 LIZ as a size standard (Applied Biosystems). CONVERT version 1.3.1 (Glaubitz, 2004) was used to convert input files for analysis in subsequent software. The genetic diversity parameters of polymorphic loci, including the number of alleles, observed heterozygosity, and expected heterozygosity were calculated by GenAlEx 6.4 (Peakall and Smouse, 2006), and Hardy–Weinberg equilibrium (HWE) for each locus and the presence of linkage disequilibria for all pairwise loci in each population were tested with POPGENE 1.32 (Yeh et al., 1997). Among the 60 genotyped individuals, the number of alleles per locus varied from one to 11 (Table 2). The observed heterozygosity and expected heterozygosity ranged from 0.000 to 0.800 and from 0.000 to 0.840, respectively. There were no significant departures from HWE over all loci for any populations, but some populations deviated from HWE at up to 15 loci ( $P < 0.05$ ) (Table 2). A significant linkage disequilibrium ( $P < 0.01$ ) was detected in

three pairwise SSR loci in two (Populations 1 and 2) out of four populations (Population 1: Pkvl007 and Pkvl010, Pkvl008 and Pkvl010; Population 2: Pkvl006 and Pkvl014). Furthermore, 16 out of 18 SSR loci were successfully amplified in 59 individuals of four related species (14–15 individuals for each species; Appendix 1). Most of these loci were polymorphic (Table 3).

## CONCLUSIONS

The set of 18 novel SSR markers reported in this study will be helpful for population genetic analysis in *P. kesiya* var. *langbianensis*, which will offer valuable information for the formulation of the rational utilization and conservation strategies of this species in the future. Furthermore, the successful cross-species amplification of these SSR markers in *Pinus* (Pinaceae) suggests their potential to be used in studies of genetic variation for related pine species.

## LITERATURE CITED

- ANDREWS, S. 2010. FastQC: A quality control tool for high throughput sequence data. Website <http://www.bioinformatics.babraham.ac.uk/projects/fastqc> [accessed 13 January 2017].
- DONG, J. X., H. J. GUO, P. LI, Y. F. ZHAO, AND X. H. YUN. 2009. Oleoresin resources and development capacity in Yunnan. *Journal of Northwest Forestry University* 24: 157–160.
- DOYLE, J. J., AND J. L. DOYLE. 1990. Isolation of plant DNA from fresh tissue. *Focus (San Francisco, Calif.)* 12: 13–15.

TABLE 2. Genetic properties of 18 polymorphic microsatellite markers in four *Pinus kesiya* var. *langbianensis* populations.<sup>a</sup>

Locus	Population 1 (N = 15)				Population 2 (N = 15)				Population 3 (N = 15)				Population 4 (N = 15)			
	A	H <sub>o</sub>	H <sub>e</sub>	HWE <sup>b</sup>	A	H <sub>o</sub>	H <sub>e</sub>	HWE <sup>b</sup>	A	H <sub>o</sub>	H <sub>e</sub>	HWE <sup>b</sup>	A	H <sub>o</sub>	H <sub>e</sub>	HWE <sup>b</sup>
Pkvl001	3	0.267	0.371	ns	6	0.400	0.564	**	3	0.133	0.338	***	4	0.133	0.187	***
Pkvl002	5	0.333	0.658	ns	2	0.000	0.391	***	4	0.200	0.296	*	3	0.267	0.338	**
Pkvl003	3	0.267	0.287	ns	4	0.267	0.293	ns	2	0.067	0.358	**	3	0.067	0.127	***
Pkvl004	4	0.000	0.516	***	5	0.067	0.482	***	4	0.133	0.293	*	3	0.067	0.184	**
Pkvl005	5	0.200	0.396	*	4	0.071	0.365	***	2	0.067	0.064	ns	2	0.067	0.064	ns
Pkvl006	7	0.400	0.491	ns	5	0.267	0.349	***	3	0.067	0.127	***	3	0.067	0.127	***
Pkvl007	3	0.133	0.240	**	3	0.133	0.291	*	1	0.000	0.000	—	2	0.067	0.064	ns
Pkvl008	4	0.200	0.393	***	4	0.133	0.242	*	3	0.133	0.127	ns	2	0.067	0.064	ns
Pkvl009	2	0.733	0.464	*	3	0.400	0.418	**	2	0.667	0.444	ns	2	0.800	0.498	*
Pkvl010	5	0.154	0.393	**	5	0.333	0.549	***	3	0.200	0.331	ns	1	0.000	0.000	—
Pkvl011	4	0.533	0.613	***	3	0.667	0.531	ns	5	0.667	0.638	***	4	0.667	0.544	**
Pkvl012	3	0.133	0.438	*	4	0.333	0.607	***	5	0.467	0.609	**	3	0.133	0.240	**
Pkvl013	6	0.133	0.551	***	11	0.333	0.840	***	4	0.000	0.459	***	6	0.143	0.561	***
Pkvl014	9	0.077	0.837	***	5	0.231	0.615	*	6	0.400	0.487	ns	9	0.333	0.616	**
Pkvl015	1	0.000	0.000	—	3	0.267	0.240	ns	1	0.000	0.000	—	1	0.000	0.000	—
Pkvl016	3	0.000	0.418	***	3	0.133	0.338	***	3	0.133	0.291	*	3	0.200	0.184	ns
Pkvl017	4	0.214	0.625	***	4	0.357	0.574	**	6	0.200	0.744	***	5	0.333	0.629	**
Pkvl018	5	0.600	0.671	*	7	0.533	0.820	***	4	0.467	0.656	***	6	0.800	0.631	***

Note: A = number of alleles; H<sub>e</sub> = expected heterozygosity; H<sub>o</sub> = observed heterozygosity; HWE = Hardy–Weinberg equilibrium; N = number of individuals sampled.

<sup>a</sup> See Appendix 1 for locality and voucher information.

<sup>b</sup> Deviation from Hardy–Weinberg equilibrium using  $\chi^2$  tests: ns = not significant, \*P < 0.05, \*\*P < 0.01, \*\*\*P < 0.001, — = monomorphic.

EDITORIAL COMMITTEE OF FLORA OF CHINA. 1978. *Flora Reipublicae Popularis Sinicae*, Vol. 7, 259–260. Science Press, Beijing, China.

GLAUBITZ, J. C. 2004. CONVERT: A user-friendly program to reformat diploid data for commonly used population genetic software packages. *Molecular Ecology Notes* 4: 309–310.

GRABHERR, M. G., B. J. HAAS, M. YASSOUR, J. Z. LEVIN, D. A. THOMPSON, I. AMIT, X. ADICONIS, ET AL. 2011. Full-length transcriptome assembly from RNA-Seq data without a reference genome. *Nature Biotechnology* 29: 644–652.

HAAS, B. J., A. PAPANICOLAOU, M. YASSOUR, M. GRABHERR, P. D. BLOOD, J. BOWDEN, M. B. COUGER, ET AL. 2013. De novo transcript sequence reconstruction from RNA-seq using the Trinity platform for reference generation and analysis. *Nature Protocols* 8: 1494–1512.

TABLE 3. Cross-amplification results showing the number of alleles detected in 18 loci from *Pinus kesiya* var. *langbianensis* in four related species.<sup>a</sup>

Locus	<i>Pinus massoniana</i>	<i>Pinus densata</i>	<i>Pinus tabuliformis</i>	<i>Pinus yunnanensis</i>
Pkvl001	2	2	2	2
Pkvl002	4	4	2	4
Pkvl003	1	2	2	1
Pkvl004	4	2	2	2
Pkvl005	3	2	1	3
Pkvl006	1	1	1	4
Pkvl007	3	1	1	1
Pkvl008	1	1	2	3
Pkvl009	3	2	2	2
Pkvl010	3	5	2	5
Pkvl011	3	4	3	2
Pkvl012	5	3	3	3
Pkvl013	—	—	—	—
Pkvl014	—	—	—	—
Pkvl015	5	7	5	4
Pkvl016	6	2	1	3
Pkvl017	2	5	6	2
Pkvl018	2	5	3	3

<sup>a</sup> See Appendix 1 for locality and voucher information.

JIANG, Y. B., T. WU, S. Y. CHEN, AND R. L. YUAN. 2007. Optimization of RAPD reaction conditions for *Pinus kesiya* var. *langbianensis*. *Journal of Northeast Forestry University* 35: 16–19.

LI, S. F., J. R. SU, W. D. LIU, X. D. LANG, X. B. HUANG, C. X. Z. JIA, Q. TONG, AND H. Y. TANG. 2015. Changes in soil organic carbon and nitrogen stocks in *Pinus kesiya* var. *langbianensis* plantation. *Forest Research* 28: 810–817.

PEAKALL, R., AND P. E. SMOUSE. 2006. GenAlEx 6: Genetic analysis in Excel. Population genetic software for teaching and research. *Molecular Ecology Notes* 6: 288–295.

SANCHEZ, M., M. J. INGROUILLE, R. S. COWAN, M. A. HAMILTON, AND M. F. FAY. 2014. Spatial structure and genetic diversity of natural populations of the Caribbean pine, *Pinus caribaea* var. *bahamensis* (Pinaceae), in the Bahaman archipelago. *Botanical Journal of the Linnean Society* 174: 359–383.

THIEL, T., W. MICHALEK, R. K. VARSHNEY, AND A. GRANER. 2003. Exploiting EST databases for the development and characterization of gene-derived SSR-markers in barley (*Hordeum vulgare* L.). *Theoretical and Applied Genetics* 106: 411–422.

WU, Z. Y. [ED.] 1986. *Flora Yunnanica*, Vol. 4 (Spermatophyta), 57–58. Science Press, Beijing, China.

YEH, F. C., R. C. YANG, AND T. BOYLE. 1997. POPGENE version 1.31: Microsoft Windows-based freeware for population genetics analysis: Quick user guide. Center for International Forestry Research, University of Alberta, Edmonton, Alberta, Canada.

YIN, X. B., S. X. GENG, H. F. MA, AND W. ZHENG. 2005. Differences of physical and chemical characteristics of gum turpentine from Simao Pine. *Journal of Nanjing Forestry University* 29: 80–84 (Natural Sciences Edition).

ZHAO, N., R. SHI, B. LI, Z. XIONG, AND J. WANG. 2016. Correlation between indexes of resin tapping and chemical components of *Pinus kesiya* var. *langbianensis*. *Journal of West China Forestry Science* 45: 63–68.

APPENDIX 1. Voucher information for *Pinus kesiya* var. *langbianensis* and its four related species used in this study.<sup>a</sup>

Species	Collection locality	Geographic coordinates	<i>N</i>
<i>Pinus kesiya</i> Royle ex Gordon var. <i>langbianensis</i> (A. Chev.) Gaussen	Ninger, Yunnan, China (Population 1)	22°57'N, 101°03'E	15
<i>Pinus kesiya</i> var. <i>langbianensis</i>	Zhenyuan, Yunnan, China (Population 2)	23°51'N, 100°53'E	15
<i>Pinus kesiya</i> var. <i>langbianensis</i>	Jiangcheng, Yunnan, China (Population 3)	22°35'N, 101°51'E	15
<i>Pinus kesiya</i> var. <i>langbianensis</i>	Lanchang, Yunnan, China (Population 4)	22°33'N, 99°50'E	15
<i>Pinus massoniana</i> D. Don	Yuping, Guizhou, China	27°30'N, 109°11'E	14
<i>Pinus densata</i> Mast	Xianggelila, Yunnan, China	28°29'N, 99°37'E	15
<i>Pinus tabuliformis</i> Carrière	Lushi, Henan, China	33°44'N, 110°49'E	15
<i>Pinus yunnanensis</i> Franch.	Yilang, Yunnan, China	24°35'N, 103°08'E	15

Note: *N* = number of individuals sampled.

<sup>a</sup>Silica gel-dried needle samples of all vouchers were deposited in the Key Laboratory for Forest Genetic and Tree Improvement and Propagation in Universities of Yunnan Province, Southwest Forestry University, Kunming, Yunnan, China.